

## Appendix

**Table 1. Intra- and inter-observer agreement**

	Intra-observer agreement	Inter-observer agreement
<b>Native T1</b>	0.99, $p<0.001$	0.99, $p<0.001$
<b>Native T2</b>	0.98, $p<0.001$	0.98, $p<0.001$
<b>ECV</b>	1.00, $p<0.001$	0.91, $p<0.001$
<b>Perfusion rest</b>	1.00, $p<0.001$	1.00, $p<0.001$
<b>Perfusion stress</b>	1.00, $p<0.001$	0.99, $p<0.001$
<b>MPR</b>	0.99, $p<0.001$	0.99, $p<0.001$
<b>LVEF</b>	0.89, $p=0.002$	0.85, $p<0.001$
<b>LV GLS</b>	0.95, $p<0.001$	0.94, $p<0.001$
<b>LV GCS</b>	0.84, $p=0.008$	0.92, $p<0.001$
<b>LV GRS</b>	0.92, $p<0.001$	0.82, $p<0.001$

Intra- and inter-observer agreement presented as intraclass correlation coefficient (ICC) together with p-values. Abbreviations: ECV = extracellular volume; MPR = myocardial perfusion reserve; LVEF = left ventricular ejection fraction; LV GLS = left ventricular global longitudinal strain; LV GCS = left ventricular global circumferential strain; LV GRS = left ventricular global radial strain.

**Table 2. RV volumes and function**

CMR findings	WD patients (n=17)	Volunteers (n=17)	p
RVEDV (ml)	170±41	165±35	0.68
RVEDVi(ml/m <sup>2</sup> )	88±19	89±15	0.93
RVESV (ml)	64±21	58±21	0.43
RVESVi (ml/m <sup>2</sup> )	33±11	31±11	0.63
RVSV (ml)	107±29	107±26	0.97
RVSVi (ml/m <sup>2</sup> )	55±12	57±10	0.55
RVEF (%)	63±9	65±9	0.40
RV GLS (%)	-18±4	-20±3	0.16
RV AVPD (mm)	-19±4	-20±3	0.63†

CMR findings are presented as mean ± standard deviation. P-values denotes the independent t-test, p-values marked † denotes the Mann Whitney U test. Abbreviations: CMR = cardiovascular magnetic resonance imaging; RV = right ventricle; RVEDV = right ventricular end-diastolic volume; RVESV = right ventricular end-systolic volume; RVSV = right ventricular stroke volume; RVEF = right ventricular ejection fraction; RV GLS = right ventricular global longitudinal strain; RV AVPD = right ventricle atrioventricular plane displacement; WD = Wilson disease.