

Supplementary Material

Table S1

Comparison of survivor characteristics with participating ($n = 493$) and non-participating ($n = 734$) parents.

	Survivors with participating parents (n = 493)	Survivors with non- participating parents (n = 734)	
	n (%)	n (%)	p value ^b
<i>Study aim</i>			<0.001
Cancer-related study aim			
Male	296 (60.0)	310 (42.2)	
Female	197 (40.0)	424 (57.8)	
Socio-demographic characteristics			
<i>Survivors' sex</i>			0.636
Male	272 (55.2)	415 (56.5)	
Female	221 (44.8)	319 (43.5)	
<i>Survivors' age at study</i>			0.065
<30 years	209 (42.4)	298 (40.6)	
30-<40 years	218 (44.2)	301 (41.0)	
=>40 years	66 (13.4)	135 (18.4)	
	Mean (SD)	Mean (SD)	p value ^b
<i>Survivors' age at study (years)</i>	32 (6.2)	33 (6.8)	0.113
Cancer-related characteristics			
<i>Age at diagnosis</i>	n (%)	n (%)	p value ^b
Age at diagnosis			0.400
<5 years	188 (38.1)	270 (36.8)	
5-<10 years	143 (29.0)	196 (26.7)	
=>10 years	162 (32.9)	268 (36.5)	
<i>Time since diagnosis</i>			0.232
<20 years	140 (28.4)	190 (25.9)	
20-<25 years	125 (25.4)	210 (28.6)	
25-<30 years	121 (24.5)	155 (21.1)	
=>30 years	107 (21.7)	179 (24.4)	
<i>Diagnosis (ICCC-3)</i>			0.803
Leukemia	167 (33.9)	248 (33.8)	
Lymphoma	84 (17.0)	133 (18.1)	
CNS tumor	68 (13.8)	109 (14.9)	
Neuroblastoma	18 (3.7)	37 (5.0)	

Retinoblastoma	14 (2.8)	13 (1.8)	
Renal tumor	33 (6.7)	50 (6.8)	
Hepatic tumor	6 (1.2)	4 (0.5)	
Bone tumor	29 (5.9)	33 (4.5)	
Soft tissue sarcoma	33 (6.7)	46 (6.3)	
Germ cell tumor	17 (3.4)	27 (3.7)	
LCH	24 (4.9)	34 (4.6)	
<i>Therapy</i> ^a			0.178
Surgery only	64 (13.0)	90 (12.3)	
<i>Chemotherapy</i>	263 (53.3)	358 (48.8)	
Radiotherapy	139 (28.2)	249 (33.9)	
SCT	26 (5.3)	32 (4.4)	
<i>Relapse</i>			0.402
No	432 (87.6)	631 (86.0)	
Yes	61 (12.4)	103 (14.0)	
	Mean (SD)	Mean (SD)	p value ^b
<i>Age at diagnosis (years)</i>	7 (4.5)	7 (4.6)	0.578
<i>Time since diagnosis (years)</i>	24 (6.8)	24 (7.0)	0.292

Note. CNS, central nervous system; LCH, Langerhans cell histiocytosis; SCT, stem cell transplantation; SD, standard deviation

^a Missing values; percentages are based on the total number of participants

^b p value calculated from Chi-square test statistics (categorical variables) or t test statistics (continuous variables) comparing survivors of participating and non-participating parents.

Table S2

Comparison of HRQOL between parents of childhood cancer survivors (CCS parents; n = 751) and parents from the general population (comparison parents; n = 454) and separately for mothers and fathers for the physical and mental HRQOL summary measures and the health domain scales and corresponding p values.

Difference between CCS parents (n = 751) and comparison parents (n = 454) ^a			
	Difference	95% CI difference	p value
Scale			
Physical functioning	1.37	(-1.01, 3.76)	0.259
Physical role functioning	-0.41	(-3.01, 2.18)	0.755
Bodily pain	1.17	(-1.97, 4.30)	0.465
General health perceptions	1.51	(-0.61, 3.62)	0.163
Vitality	1.09	(-1.15, 3.34)	0.340
Social role functioning	-0.26	(-2.64, 2.12)	0.831
Emotional role functioning	-0.85	(-3.18, 1.47)	0.472
Mental health	0.56	(-1.44, 2.55)	0.583
Summary measure			
Physical HRQOL	0.62	(-0.56, 1.81)	0.300
Mental HRQOL	-0.37	(-1.56, 0.82)	0.545

**Difference between CCS mothers (n = 442)
and comparison mothers (n = 263) ^b**

	Difference	95% CI difference	p value
Scale			
Physical functioning	2.96	(0.17, 5.76)	0.038
Physical role functioning	0.84	(-2.48, 4.16)	0.620
Bodily pain	-0.09	(-4.11, 3.92)	0.964
General health perceptions	1.80	(-0.79, 4.39)	0.173
Vitality	0.52	(-2.25, 3.29)	0.711
Social role functioning	-0.10	(-3.15, 2.95)	0.950
Emotional role functioning	-1.27	(-4.35, 1.81)	0.418
Mental health	0.86	(-1.65, 3.36)	0.503
Summary measure			
Physical HRQOL	1.13	(-0.35, 2.60)	0.134
Mental HRQOL	-0.60	(-2.12, 0.92)	0.437

**Difference between CCS fathers (n = 309) and
comparison fathers (n = 191) ^c**

	Difference	95% CI difference	p value
Scale			
Physical functioning	0.86	(-2.96, 4.69)	0.658
Physical role functioning	-0.80	(-5.24, 3.64)	0.724
Bodily pain	4.40	(-0.56, 9.35)	0.082
General health perceptions	1.44	(-2.07, 4.96)	0.420
Vitality	1.90	(-1.38, 5.18)	0.255

Social role functioning	-0.11	(-3.60, 3.38)	0.951
Emotional role functioning	0.38	(-3.07, 3.83)	0.829
Mental health	0.05	(-2.81, 2.91)	0.974
Summary measure			
Physical HRQOL	0.76	(-1.29, 2.80)	0.467
Mental HRQOL	-0.17	(-1.88, 1.53)	0.841

Note. CCS, childhood cancer survivor; CI, confidence interval; HRQOL, health-related quality of life; difference with 95% confidence interval (CI) comparing CCS parents and comparison parents; positive differences indicate better HRQOL in CCS parents and negative differences poorer HRQOL compared to comparison parents; p value for the adjusted difference between the two samples; p values <0.05 are indicated with bold font.

Regression analyses adjusted for

^a study aim mentioned, number of children, migration background, employment status, partnership, and chronic condition or health problem

^b study aim mentioned, number of children, migration background, employment status, and chronic condition or health problem

^c study aim mentioned, number of children, and partnership status

Table S3

Multilevel univariable linear regression analysis for physical and mental health-related quality of life (HRQOL) including parents of childhood cancer survivors (CCS parents).

Characteristics	Physical HRQOL (PCS)			Mental HRQOL (MCS)		
	Coeff	95% CI	p value ^b	Coeff	95% CI	p value ^b
Study aim mentioned				0.253		
No	Ref			Ref		0.036
Yes	0.89	(-0.63, 2.40)		-1.54	(-2.99, -0.10)	
Socio-demographic characteristics						
Sex				0.681		
Fathers	Ref			Ref		0.006
Mothers	-0.29	(-1.65, 1.08)		-1.55	(-2.65, -0.44)	
Age ^a				<0.001		
41-65 years	Ref			Ref		0.476
66-75 years	-4.17	(-5.70, -2.63)		0.53	(-0.92, 1.98)	
Number of children ^a				0.246		
1 child	Ref			Ref		
2 children	2.03	(-2.20, 6.25)		-1.82	(-5.70, 2.06)	
>2 children	0.81	(-3.41, 5.03)		-1.65	(-5.54, 2.24)	
Migration background ^a				0.430	<0.001	
No	Ref			Ref		
Yes	-0.87	(-3.03, 1.29)		-3.87	(-5.82, -1.91)	
Education ^a				0.046		
Compulsory schooling	-0.53	(-3.87, 0.82)		-0.58	(-2.73, 1.58)	
Vocational training	Ref			Ref		
Upper secondary education	1.86	(-0.04, 3.77)		0.78	(-0.93, 2.50)	
University education	1.38	(-0.67, 3.43)		1.43	(-0.47, 3.33)	
Employment ^a				<0.001		
Employed	Ref			Ref		0.171
Unemployed	-3.48	(-6.01, -0.92)		-2.12	(-4.38, 0.14)	

Retired	-3.76	(-5.29, -2.23)		-0.06	(-1.50, 1.37)	
Partnership ^a			0.003			0.108
Yes	Ref			Ref		
No	-3.59	(6.00, -1.19)		-1.81	(-4.01, 0.40)	
Questionnaire language			<0.001			<0.001
German	Ref			Ref		
French or Italian	-3.32	(-5.00, -1.63)		-3.48	(-5.07, -1.89)	
Chronic condition or health problem ^a			<0.001			0.011
No	Ref			Ref		
Yes	-8.61	(-9.91, -7.32)		-1.61	(-2.86, -0.36)	

Cancer-related characteristics

Age at diagnosis		0.093		0.543	
<5 years	Ref		Ref		
5-<10 years	-1.95	(-3.75, -0.159)	-0.11	(-1.84, 1.62)	
>= 10 years	-1.24	(-3.01, 0.53)	-0.90	(-2.59, 0.79)	
Time since diagnosis		0.002		0.921	
<20 years	Ref		Ref		
20-<25 years	-1.80	(-3.78, 0.19)	-0.37	(-2.30, 1.56)	
25-<30 years	-2.72	(-4.74, -0.71)	-0.44	(-2.39, 1.51)	
>=30 years	-3.92	(-6.04, -1.80)	0.19	(-1.85, 2.23)	
Cancer diagnosis		0.147		0.808	
Leukemia	Ref		Ref		
Lymphoma	-2.21	(-4.24, -0.17)	-0.97	(-2.91, 0.97)	
CNS tumour	-0.98	(-3.35, 1.40)	-0.45	(-2.72, 1.81)	
Solid tumour	-0.13	(-1.98, 1.72)	-0.49	(-2.25, 1.28)	
Therapy		0.626		0.086	
Surgery only	Ref		Ref		
Chemotherapy	0.45	(-1.82, 2.71)	1.69	(-0.46, 3.84)	
Radiotherapy	-0.49	(-2.97, 1.99)	-0.24	(-2.59, 2.11)	
SCT	1.49	(-2.38, 5.35)	1.91	(-1.75, 5.56)	
Relapse		0.506		0.470	
No	Ref		Ref		
Yes	0.76	(-1.49, 3.01)	-0.79	(-2.94, 1.35)	
Late effects survivor (self-reported) ^b		0.026		0.006	
No	Ref		Ref		
Yes	-2.23	(-4.13, -0.33)	-2.09	(-3.83, -0.35)	

Not assessed	-2.01	(-3.72, 0.31)	0.69	(-0.92, 2.31)
--------------	-------	---------------	------	---------------

Note. HRQOL, health-related quality of life; PCS, physical component summary; MCS, mental component summary; n, number; Coeff, coefficient; CI, confidence interval; Ref, reference category

p values <0.05 are indicated with bold font.

^a Missing values

^b p values calculated from Wald tests

Table S4

Multilevel multivariable linear regression analyses for physical and mental health-related quality of life (HRQOL) including parents of childhood cancer survivors (CCS parents).

Characteristics	Physical HRQOL (PCS) (n parents = 657, n groups = 453)			Mental HRQOL (MCS) (n parents = 695, n groups = 471)		
	Coeff	95% CI	p value ^c	Coeff	95% CI	p value ^c
Study aim mentioned ^a						
No	-	-	-	-	-	-
Yes	-	-	-	-	-	-
Sex						
Fathers	-	-	-	Ref	-	0.002
Mothers	-	-	-	-1.80	(-2.93, -0.66)	-
Age ^b						
41-65 years	Ref	-	0.160	-	-	-
66-75 years	-1.43	(-3.43, 0.56)	-	-	-	-
Migration background ^b						
No	-	-	-	Ref	-	-
Yes	-	-	-	-3.88	(-5.79, -1.97)	0.000
Education ^b						
Compulsory schooling	-1.23	(-3.38, 0.92)	-	-	-	-
Vocational training	Ref	-	-	-	-	-
Upper secondary education	2.01	(0.32, 3.71)	-	-	-	-
University education	1.79	(-0.04, 3.62)	-	-	-	-
Employment ^b						
Employed	Ref	-	-	-	-	-
Unemployed	-2.23	(-4.57, 0.12)	-	-	-	-
Retired	-1.55	(-3.49, 0.39)	-	-	-	-
Partnership ^b						
Yes	Ref	-	-	-	-	-
No	-1.72	(-3.87, 0.43)	-	-	-	-
Questionnaire language						
			0.000			0.001

German	Ref		Ref	
French or Italian	-2.91	(-4.44, -1.38)	-2.75	(-4.36, -1.15)
Chronic condition or health problem^b		0.000		0.006
No	Ref		Ref	
Yes	-8.26	(-9.56, -6.96)	-1.76	(-3.02, -0.51)
Time since diagnosis		0.585		
<20 years	Ref			
20-<25 years	-1.21	(-2.96, 0.54)		
25-<30 years	-0.68	(-2.56, 1.19)		
>=30 years	-0.41	(-2.48, 1.67)		
Late effects survivor (self-reported)^b		0.057		0.029
No	Ref		Ref	
Yes	-1.09	(-2.83, 0.64)	-1.27	(-3.03, 0.48)
Not assessed	-1.81	(-3.30, -0.33)	1.09	(-0.49, 2.67)

Note. HRQOL, health-related quality of life; PCS, physical component summary; MCS, mental component summary; n, number; Coeff, coefficient; CI, confidence interval; Ref, reference category

p values <0.05 are indicated with bold font.

^a Excluded in multivariable regression analysis due to its correlation with late effects

^b Missing values

^c p values calculated from Wald tests