



# Supplemental Appendix


Supplemental Figure i.



Supplemental Figure ii.



Supplemental Figure iii.



**Supplemental Table i.** Model fit comparisons for the ARAT

Model	n par	AIC	BIC	logLik	deviance	$\chi^2$ Decrease	df	P( $\chi^2$ )
Intercept Only	3	3777.1	3789.8	-1885.6	3771.1			
Linear Time	6	3537.5	3562.7	-1762.8	3525.5	245.634	3	<0.001
Inverse Square	6	3379.9	3405.1	-1684.0	3367.9	157.594	0	na
Inverse Cubic	6	3479.8	3505.1	-1733.9	3467.8	(increased)	0	na
3-Param. Exponential	9	3283.5	3325.5	-1631.8	3263.5	204.3	3	<0.001
Quadratic Time	10	3373.1	3415.1	-1676.5	3353.1	(increased)	1	na
Cubic Time	11	3338.1	3384.3	-1658.0	3316.1	37.016	1	<0.001
<b>2-Knot Spline†</b>	<b>15</b>	<b>3261.9</b>	<b>3325.0</b>	<b>-1615.9</b>	<b>3231.9</b>	<b>84.198</b>	<b>4</b>	<b>&lt;0.001</b>

Note, all models included random slopes for all time parameters. The 2-knot spline shown refers to overall best fitting spline with knots at 5- and 11-weeks. AIC = Akaike's information criterion; BIC = Bayesian Information Criterion; loglik = log(Likelihood); df = degrees of freedom; p( $\chi^2$ ) = p-value from Wald's test for the change in deviance compared to previous model. † = best fitting model based on AIC.

**Supplemental Table ii.** Model fit comparisons for the FM-UE

Model	n par	AIC	BIC	logLik	deviance	$\chi^2$ Decrease	df	P( $\chi^2$ )
Intercept Only	3	2969.9	2981.8	-1481.9	2963.9			
Linear Time	6	2772.2	2796	-1380.1	2760.2	203.696	3	< 2.2e-16
Inverse Square	6	2675.1	2698.9	-1331.5	2663.1	97.12	0	na
Inverse Cubic	6	2759.2	2783	-1373.6	2747.2	(increased)	0	na
3-Param. Exponential	6*	2578.9	2606.8	-1282.5	2564.9	182.2	0	na
Quadratic Time	10	2619.8	2659.5	-1299.9	2599.8	(increased)	4	na
Cubic Time	11	2602.3	2646	-1290.1	2580.3	19.535	1	9.88E-06
<b>2-Knot Spline†</b>	<b>15</b>	<b>2554.1</b>	<b>2613.7</b>	<b>-1262.1</b>	<b>2524.1</b>	<b>56.15</b>	<b>4</b>	<b>1.87E-11</b>

Note, all models included random slopes for all time parameters. The 2-knot spline shown refers to overall best fitting spline with knots at 5- and 11-weeks. AIC = Akaike's information criterion; BIC = Bayesian Information Criterion; loglik = log(Likelihood); df = degrees of freedom; p( $\chi^2$ ) = p-value from Wald's test for the change in deviance compared to previous model. \* = model failed to converge with random-effect of the rate parameter, so was refit with only a fixed rate parameter. † = best fitting model based on AIC.

**Supplemental Table iii.** Model fit comparisons for the Use Ratio

Model	n par	AIC	BIC	logLik	deviance	$\chi^2$ Decrease	df	P( $\chi^2$ )
Intercept Only	3	-459.45	-446.82	232.73	-465.45			
3-Param. Exponential	4*	-548.3	-527.2	279.10	-558.28	92.8	1	< 2.2e-16
Linear Time	6	-533.04	-507.76	272.52	-545.04	(increased)	2	na
Inverse Square	6	-608.13	-582.85	310.06	-620.13	75.0868	0	
Inverse Cubic	6	-587.68	-562.4	299.84	-599.68	(increased)	0	
Quadratic Time	10	-588.96	-546.83	304.48	-608.96	9.2819	4	0.054427
Cubic Time	11	-597.73	-551.39	309.87	-619.73	10.7704	1	1.03E-03
<b>2-Knot Spline†</b>	<b>15</b>	<b>-619.19</b>	<b>-556</b>	<b>324.60</b>	<b>-649.19</b>	<b>29.4616</b>	<b>4</b>	<b>6.30E-06</b>

Note, all models included random slopes for all time parameters. The 2-knot spline shown refers to overall best fitting spline with knots at 5- and 11-weeks. AIC = Akaike's information criterion; BIC = Bayesian Information Criterion; loglik = log(Likelihood); df = degrees of freedom; p( $\chi^2$ ) = p-value from Wald's test for the change in deviance compared to the previous model. \* = model failed to converge with random-effects of the rate and change parameters, so was refit with only a random effect on the asymptote. † = best fitting model based on AIC.

**Supplemental Table iv.** Details of the best fitting spline model for the ARAT.

<b>AIC</b>	<b>BIC</b>	<b>logLik</b>	<b>deviance</b>	<b>df.resid</b>
3261.9	3325.0	-1615.9	3231.9	481

Scaled residuals:

Min	1Q	Median	3Q	Max
-3.4678	-0.3429	0.0152	0.2958	3.0871

Random effects:

Groups	Name	Variance	Std.Dev.	Corr
SubIDName	(Intercept)	553.605	23.529	
	WeeksPostStroke	13.226	3.637	-0.56
	knot05	11.695	3.420	0.43 -0.93
	knot11	1.645	1.283	0.43 -0.47 0.13
	Residual	7.419	2.724	

Number of obs: 496, groups: SubIDName, 98

Fixed effects:

	Estimate	Std. Error	df	t value	Pr(> t )
(Intercept)	12.7206	2.8857	85.6438	4.408	3.01e-05 ***
WeeksPostStroke	3.1852	0.5159	68.0570	6.174	4.19e-08 ***
knot05	-1.8513	0.5306	67.2708	-3.489	0.00086 ***
knot11	-1.2056	0.1836	81.1598	-6.566	4.54e-09 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1

**Supplemental Table v.** Details of the best fitting spline model for the FM-UE.

<b>AIC</b>	<b>BIC</b>	<b>logLik</b>	<b>deviance</b>	<b>df.resid</b>
2554.1	2613.7	-1262.1	2524.1	378

**Scaled residuals:**

<b>Min</b>	<b>1Q</b>	<b>Median</b>	<b>3Q</b>	<b>Max</b>
-3.1768	-0.3573	0.0209	0.3821	3.3408

**Random effects:**

<b>Groups</b>	<b>Name</b>	<b>Variance</b>	<b>Std.Dev.</b>	<b>Corr</b>
SubIDName	(Intercept)	622.746	24.955	
	WeeksPostStroke	5.707	2.389	-0.54
	knot05	6.314	2.513	0.22 -0.73
	knot11	3.833	1.958	0.30 -0.25 -0.47
	Residual	7.847	2.801	

Number of obs: 393, groups: SubIDName, 68

**Fixed effects:**

	<b>Estimate</b>	<b>Std. Error</b>	<b>df</b>	<b>t value</b>	<b>Pr(&gt; t )</b>
(Intercept)	24.7254	3.2574	66.1576	7.591	1.41e-10 ***
WeeksPostStroke	2.7116	0.4118	57.3172	6.585	1.52e-08 ***
knot05	-1.4258	0.4737	52.5736	-3.010	0.004008 **
knot11	-1.1319	0.2939	57.4449	-3.851	0.000298 ***

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1



**Supplemental Table vi.** Details of the best fitting spline model for the Use Ratio.

<b>AIC</b>	<b>BIC</b>	<b>logLik</b>	<b>deviance</b>	<b>df.resid</b>
-619.2	-556.0	324.6	-649.2	484

**Scaled residuals:**

<b>Min</b>	<b>1Q</b>	<b>Median</b>	<b>3Q</b>	<b>Max</b>
-3.6433	-0.4583	-0.0067	0.3897	4.0693

**Random effects:**

<b>Groups</b>	<b>Name</b>	<b>Variance</b>	<b>Std.Dev.</b>	<b>Corr</b>
SubIDName	(Intercept)	0.0808610	0.28436	
	WeeksPostStroke	0.0019462	0.04412	-0.43
	knot05	0.0024237	0.04923	0.31 -0.90
	knot11	0.0006486	0.02547	-0.02 0.18 -0.59
	Residual	0.0054398	0.07375	

Number of obs: 499, groups: SubIDName, 98

**Fixed effects:**

	<b>Estimate</b>	<b>Std. Error</b>	<b>df</b>	<b>t value</b>	<b>Pr(&gt; t )</b>	
(Intercept)	0.395352	0.042263	74.221063	9.355	3.47e-14	***
WeeksPostStroke	0.045277	0.008431	68.480103	5.370	1.01e-06	***
knot05	-0.038468	0.009880	72.821508	-3.893	0.000217	***
knot11	-0.005270	0.004160	84.278495	-1.267	0.208719	

---

Signif. codes: 0 '\*\*\*' 0.001 '\*\*' 0.01 '\*' 0.05 '.' 0.1 ' ' 1