



Supplemental Figure 1. Overview of ENIGMA Suicidal Thoughts and Behaviours consortium cohorts included in the analysis.

Table S1. Main diagnosis of participants and study exclusion criteria in the different sites

Site	Main diagnosis participants	Inclusion criteria	Exclusion criteria
BiDirect	MDD	For patients: The recruitment of outpatients was limited to those who had been hospitalised due to depression at least once during the 12 months period prior to inclusion into the study. Inclusion criteria were (i) age ( $\geq 35$ and $< 66$ years) and (ii) current in- or outpatient treatment due to acute depression. For healthy controls: The participants ( $\geq 35$ and $< 66$ years) had been randomly sampled from the population register of the city of Münster.	Exclusion criteria for both patients and controls were (i) compulsory admission, (ii) comorbid dementia, and (iii) comorbid drug abuse (including alcohol).
Brazilian High Risk Cohort Porto Alegre City	Various	Children and adolescents 6 to 14 years of age at baseline were attending one of the 57 public schools in two Brazilian cities (São Paulo and Porto Alegre). From the total cohort (N = 2511), 1004 children were invited for MRI scanning, and the images from 741 were obtained at cohort baseline.	Contraindications to MRI assessments (metal, braces, etc.)
Chiba University	OCD	For patients: OCD patients who are willing to receive CBT	For patients: Suicide attempt, Antisocial Personality Disorder; For healthy controls: History of a psychiatric disorder
CHU Montpellier BICS	MDD	For patients : right-handed, Caucasian adults aged between 18 and 70 years. current depressive episode according to the DSM-5 criteria. For healthy controls: right-handed, Caucasian adults aged between 18 and 70 years. No history of psychiatric disorders according to the DSM-5 criteria . No lifetime history of suicide attempt	Current neurologic or inflammatory disease; CRP $>10$ mg/L; current antibiotic or anti-inflammatory treatment; history of head trauma with loss of consciousness; current substance misuse, excluding tobacco; lifetime history of manic or hypomanic episode, schizophrenia, or schizoaffective disorder according to the DSM-5; pregnancy and breastfeeding; contraindication to MRI.
CHU Montpellier IMPACT	MDD	Aged between 18 and 65 years, having a history of suicide attempt within the last 12 months	Lifetime history of schizophrenia, a current alcohol/ illicit drug use disorder, a current manic or hypomanic episode, a lifetime history of severe brain injury or neurologic disease and pregnancy;contraindication to MRI.
CHU Montpellier UF8555	MDD/BD	Healthy controls : Caucasian right-handed females, $>18$ yo, no lifetime history of psychiatric disorder	Lifetime history of severe head trauma, CNS disorder, schizophrenia, and a history of alcohol or drug abuse or

		Patients : Caucasian right-handed females, euthymic ( Hamilton Depression Rating Scale score <7 and Young Mania Rating Scale score < 7, history of major past depressive episode ; >18 yo	dependence within the past 12 months, pregnancy and breastfeeding ; contraindication to MRI.
DCHS	MDD/PTSD	For patients: enrolled in DCHS study; mothers between ages of 18 and 50; could have PTSD, MDD, or both; For healthy controls: enrolled in DCHS study; mothers between ages of 18 and 50; no psychopathology	For patients: no psychopathology; brain injury or loss of consciousness for longer than 30 minutes; MRI exclusion criteria; For healthy controls: brain injury or loss of consciousness for longer than 30 minutes; MRI exclusion criteria
MOODS team DEP-ARREST- CLIN	MDD	For patients: 18-65 years, current MDD diagnosis, minimum HDRS scores of 18, antidepressant-free at least one month prior to study inclusion For healthy controls: absence of current or past psychiatric disorders or somatic conditions (including nasal polyposis and acute sinusitis or rhinitis)	For patients: diagnosis of bipolar disorder, psychotic disorder, eating disorder or addiction according to the DSM-V For all participants: pregnancy or breastfeeding, nasal polyposis, acute or chronic sinusitis or rhinitis.
EPISCA (Leiden adolescents)	MDD, PTSD	For patients: aged between 12 and 19, estimated full scale IQ>80, right handed, normal or corrected to normal vision, sufficient understanding of the Dutch language, no history of neurological impairments and no contraindications for MRI. Having a depressive or anxiety disorder as classified by DSM-IV criteria (including PTSD), no current or prior use of antidepressants. For healthy controls: aged between 12 and 19, estimated full scale IQ>80, right handed, normal or corrected to normal vision, sufficient understanding of the Dutch language, no history of neurological impairments and no contraindications for MRI. No current or passed DSM-IV classifications, no clinical scores on validated mood and anxiety questionnaires, no history of traumatic experiences, and no current psychotherapeutic and/or psychopharmacological intervention of any kind	For both patients and healthy controls: primary DSM-IV clinical diagnosis of ODD, CD, pervasive developmental disorder, Tourette's syndrome, OCD, bipolar disorder, and psychotic disorder, current substance use, history of neurological disorders or severe head injury, and pregnancy
ETPB-STB	MDD	For patients: Eligible participants included men and women, ages 18–65 years. Subjects with MDD had been diagnosed with recurrent MDD without psychotic features using the Structured Clinical Interview for Axis I DSM-IV Disorders (SCID)-Patient Version [40]. Subjects were required to have a score ≥20 on the Montgomery-Åsberg Depression Rating	For patients: psychotic features; For healthy controls: family history of Axis I disorders in first degree relatives

Scale (MADRS). Subjects had to have not responded to at least one adequate antidepressant trial during their current episode, as assessed using the Anti-depressant Treatment History Form, and the current episode had to have lasted at least four weeks. Subjects were free from psychotropic medications in the two weeks before randomization (5 weeks for fluoxetine, 3 weeks for aripiprazole);

For healthy controls: Healthy control subjects consisted of males and females, 18–65 years old with no Axis I disorder as determined by SCID-NP, and no family history of Axis I disorders in first degree relatives. Healthy control subjects were free of medications affecting neuronal function or cerebral blood flow or metabolism. Subjects in both groups were in good physical health as determined by medical history, physical exam, blood labs, electrocardiogram, chest x-ray, urinalysis, and toxicology.

FSL_NPL- OCD sample	OCD	For patients: DSM-V diagnosis For all participants: age between 18-65 years	Exclusion criteria for all participants: history of alcohol or drug abuse within past 2 years, lifetime drug dependence, traumatic head injury with loss of consciousness, MRI contraindications. For patients: past or present major medical illness or neurological disorder, mental retardation, brain abnormalities, vascular lesions. For controls: current or lifetime DSM-V psychiatric or personality disorder according to SCID, first-degree relative with psychosis, mental retardation, minor or major DSM-V major neurocognitive disorder, any brain abnormalities or vascular lesions
FSL_NPL- SZ sample	Psychosis	For patients: DSM-V diagnosis For all participants: age between 18-65 years	Exclusion criteria for all participants: history of alcohol or drug abuse within past 2 years, lifetime drug dependence, traumatic head injury with loss of consciousness, MRI contraindications. For patients: past or present major medical illness or neurological disorder, mental retardation, brain abnormalities, vascular lesions. For controls: current or lifetime DSM-V psychiatric or personality disorder according to SCID, first-degree relative with psychosis,

			mental retardation, minor or major DSM-V major neurocognitive disorder, any brain abnormalities or vascular lesions
FOR2107 Marburg	MDD	For patients: age 18-65 years; patients were diagnosed by SCID-Interview with major depressive disorder (currently depressed or remitted) or with bipolar disorder (currently depressed, (hypo)manic or remitted) For healthy controls: age 18-65 years	Exclusion criteria all: any MRI contraindications; any neurological abnormalities. Exclusion criteria controls: any current or former psychiatric disorder; Exclusion criteria patients: substance dependence or current benzodiazepine treatment (wash out of at least three half-lives before study participation)"
FOR2107 Muenster	MDD	For patients: age 18-65 years; patients were diagnosed by SCID-Interview with major depressive disorder (currently depressed or remitted) or with bipolar disorder (currently depressed, (hypo)manic or remitted) For healthy controls: age 18-65 year	Exclusion criteria all: any MRI contraindications; any neurological abnormalities. Exclusion criteria healthy controls: any current or former psychiatric disorder; Exclusion criteria patients: substance dependence or current benzodiazepine treatment (wash out of at least three half-lives before study participation)
Ghent		For patients: euthymic outpatients with a history of suicide attempts (SAs) and euthymic outpatients without a history of suicide attempt (NA) were included. All patients had a history of depression. They were recruited in different centres in the Flemish part of Belgium; For healthy controls: Healthy controls (HC), without any history of depression or suicidal behavior, were included.	For patients and controls: The MINI was used to rule out lifetime bipolar disorder, lifetime psychotic disorder and alcohol abuse in the past 12 months, as these were exclusion criteria. Possible neurological disorders and current medication use were questioned in detail by the primary researcher. Subjects with a lifetime history of neurological disorders or using psychotropic medication at time of participation, except for selective serotonin reuptake inhibitors and serotonin–norepinephrine reuptake inhibitors, were excluded.
GIPSI BD	BD	Diagnosis of Bipolar Disorder I or II based on DSM-IV-TR criteria using the Diagnostic Interview for Genetics Studies (DIGS), age between 18 and 60 years, education level between 5 and 16 years	Personal history of neurological disorders, mental disability, autism, electroconvulsive therapy and/or traumatic brain injury
GIPSI SZ	Psychosis	Diagnosis of Schizophrenia based on DSM-IV-TR criteria using the Diagnostic Interview for Genetics Studies (DIGS), age between 18 and 60 years, education level between 5 and 16 years	Personal history of neurological disorders, intellectual disability, autism, electroconvulsive therapy and/or traumatic brain injury
Grady Trauma Project Emory University	PTSD	All: 18-65 years of age, English-speaking, endorsed at least 1 criterion A trauma. For PTSD: meet DSM-IV criteria for PTSD	All: Current psychotic symptoms or bipolar disorder, current substance or alcohol dependence, history of head trauma, taking

			any psychoactive medication, current illegal drug use (verified with urine drug screen within 24 hours of scan)
Halifax	BD	<p>Inclusion criteria. The BD patients (both Li and non-Li groups) had to have: (i) a diagnosis of bipolar I or II disorder made by a psychiatrist using the SCID; (ii) at least 10 years of illness; (iii) a history of at least five episodes of illness (including manic, depressive, or mixed episodes); (iv) current Hamilton Depression Rating Scale, 17-item version (HAM-D-17) score &lt; 7; (v) current Young Mania Rating Scale (YMRS) score &lt; 5; (vi) current Clinical Global Impressions Scale–Bipolar (CGI-BP) score &lt; 3; and (vii) a period of euthymia for at least four months prior to scanning, as aside from state- related factors, patients in acute episodes may present with additional difficult to control confounding variables, including recent medication change or substance abuse. The non-Li group had to have less than three months of lifetime Li exposure, more than 24 months prior to the scanning. The Li group had to have a current Li treatment lasting a minimum of 24 months. (Halifax Diabetes Study: The subjects with BD were required to 1) have the diagnosis of bipolar I or II disorder made by a psychiatrist; and 2) be at least 18 years of age. Patients were excluded if they had 1) the diagnosis of organic mood disorder; 2) mood disorder not otherwise specified; or 3) more than one lifetime course of electroconvulsive therapy or electroconvulsive therapy within the last 6 months. Halifax High Risk Study: Families were identified through adult probands with BD, who had participated in 1) previous genetic and high-risk studies for the Halifax sample. Only the offspring from these families, not the probands, were a part of the MRI study. The offspring from BD parents were divided into two subgroups: 1) the Unaffected HR group, which consisted of 50 offspring with no lifetime history of psychiatric disorders. These individuals were at an increased risk for BD because they had one parent affected with a primary mood disorder. 2) The Affected Familial group, which consisted of 36 offspring who met criteria for a lifetime Axis I diagnosis of mood disorders (i.e., a personal history of at least one episode of depression, hypomania, or mania meeting full DSM-IV criteria). When available, we recruited more than one offspring per family. From this study, we provided data only from patients who had a personal history of bipolar disorder.</p>	<p>Exclusion criteria. Individuals from any of the three groups were excluded if they met any of the magnetic resonance imaging (MRI) exclusion criteria or had any serious medical illness (e.g., brain injury, Cushings disease, or conditions treated with corticosteroids). Individuals with BD were excluded if they had: (i) more than one lifetime course of electroconvulsive therapy (ECT) or ECT in the previous 12 months; (ii) comorbid psychiatric disorders, and/or personality disorder; (iii) active substance abuse in the previous 12 months; (iv) significant change in their medication in the previous three months; or (v) current psychotic features or acute suicidality. Individuals from the non-Li group were excluded if they had: (i) Li exposure &lt; 2 years before the scanning; or (ii) lifetime Li exposure of more than three months. The neuropsychiatrically healthy individuals were excluded if they had a personal history of psychiatric disorders. The neuropsychiatrically healthy, euglycemic subjects were excluded if they had 1) a personal history of psychiatric disorders; or 2) T2DM. Subjects from any group were excluded if they 1) met any magnetic resonance imaging (MRI) exclusion criteria; 2) suffered from substance abuse in the last 12 months; had a history of 3) neurodegenerative disorders; or 4) cerebrovascular disease/stroke, as we were interested in the more subtle T2DM-related neuronal changes.</p>

Houston BD	BD	<p>Patients: Bipolar I or II disorder diagnosed by SCID-I, according to DSM-IV-R diagnostic criteria; Any current mood state; Any comorbid psychiatric disorder, except for current substance abuse or dependence (see exclusion criteria); Age 18-65 years old.</p> <p>Controls: No lifetime diagnosis of a major psychiatric (mood, anxiety, psychotic, addictive) and neurological disorders according to DSM-IV-R; No family history of any axis I psychiatric disorder and/or hereditary neurological disorders in first-degree relatives; 18-65 years old.</p>	<p>Healthy controls reporting current or past Axis I disorders, suicidal history and a first-degree relative with any Axis I disorder were excluded. Participants with any endocrinological disease, head trauma, neurological disease, and family history of any hereditary neurological disorder or medical conditions such as hypertension, diabetes, active liver disease and kidney problems were excluded from this study.</p>
Jena-SB	MDD	<p>Patients and healthy controls aged between 18–63, right-handed. Patients must meet the criteria of a current major depressive episode according to SCID-I. SA was defined according to Silverman et al. 2007 as a “self-inflicted, potentially injurious behavior with a nonfatal outcome for which there is evidence (either explicit or implicit) of intent to die.” Healthy control subjects must have no current or previous mental disorders, suicidal ideation, or suicide attempts.</p>	<p>Comprised a lifetime history of schizophrenia or bipolar disorder, a history of alcohol/substance abuse or dependence spanning the previous 6 months, a major general medical condition requiring ongoing pharmacological treatment, a lifetime history of severe head trauma or central nervous system disorder, and contraindication to MRI. ■</p>
KASP	Psychosis	<p>Patients who seek health care for psychotic symptoms for the first time were recruited at psychiatric emergency wards and in- or outpatient facilities at the psychiatric clinics located in Stockholm, Sweden. A baseline diagnosis is established based on a structured clinical interview of the Diagnostic and Statistical Manual of Mental Disorders IV, DSM-IV (Structured Clinical Interview for DSMIV-Axis I Disorders, SCID-I) at the time of inclusion. Healthy controls were recruited by advertisement and matched on age and sex.</p>	<p>Exclusion criteria for patients were an ongoing or previous prescription of an antipsychotic for more than 30 days, severe somatic and neurological diseases, current substance abuse (except nicotine use), or autism spectrum disorder. The Mini International Neuropsychiatric Interview (MINI), performed by either a resident or a specialist in psychiatry, was used to exclude previous or current psychiatric illness among the healthy controls. Further exclusion criteria were former or current use of illegal drugs, first-degree relatives with psychotic illness or bipolar disorder, as well as neurologic disease and/or severe somatic disease.</p>
LTS Colorado	HC	<p>Ongoing participation in the Colorado Longitudinal Twin Study (LTS).</p>	<p>Contraindications to MRI scanning or health conditions resulting in an inability to provide informed consent or complete tasks.</p>
McGill University	MDD	<p>Patients: Aged between 25 to 55, right-handed. Current major depressive episode as part of MDD according to SCID-1 and HDRS&gt;19.</p> <p>CLC: no personal or first-degree family history of suicidal behavior.</p> <p>HC: no personal or second-degree family history of</p>	<p>CLC: Lifetime history of schizophrenia, bipolar disorder, and history of alcohol/drug abuse or dependence over the last 6 months; history of major general medical condition that requires current pharmacological treatment; no use of an accepted method of contraception if nonmenopausal woman; pregnancy; a lifetime</p>

		depressive disorders or suicidal behavior and no current psychotropic treatment	history of severe head trauma, or central nervous system disorder; counter-indication to MRI (ferromagnetic body implants, piercings, claustrophobia, inability to lie still during one hour, etc.).
MR-IMPACT	MDD	Patients: age 11-17, current DSM-IV unipolar MDD diagnosis determined using child and parent interviews using the KSADS-PL, score of 27 or higher on the MFQ	Exclusion criteria for all participants: alcohol dependence, drug dependence, pervasive developmental disorder or generalised learning problem, pregnancy or breastfeeding, current medication use that could adversely interaction with SSRIs, MRI contraindications, brain abnormalities, intolerance to the MRI environment
Muenster Neuroimaging Cohort	MDD	"Inclusion criteria: age 17-65 years; patients were diagnosed with major depressive disorder by SCID-Interview, currently depressed or remitted.	For patients: presence of bipolar disorder, schizoaffective disorders and schizophrenia; substance-related disorders or current benzodiazepine treatment (wash out of at least three half-lives before study participation), and former electroconvulsive therapy. For controls: any current or former psychiatric disorder. For patients & controls: any neurological abnormalities, MRI contra-indications
PAFIP	Psychosis	For patients: 15-50 years, DSM-IV diagnosis of schizophrenia, schizophreniform disorder, schizoaffective disorder, delusional disorder, brief reactive psychosis or psychosis not otherwise specified For healthy controls: no history of mental illness treatment and no axis-I disorder	For patients: history of neurological disease, head injury, mental retardation or drug dependence. For healthy controls: current or past psychiatric diagnosis, mental retardation, neurological or general medical illnesses. Psychosis in first-degree relatives
San Raffaele Hospital	BD/MDD	For patients: Age 18-65, a baseline Hamilton Depression Rating Scale (HDRS) score of 17 or higher; For healthy controls: Age 18-65	For patients: other diagnoses on Axis I; mental retardation on Axis II; pregnancy, history of epilepsy, major medical and neurological disorders; treatment with long-acting neuroleptic drugs in the last three months before admission; history of drug or alcohol dependency or abuse within the last six months and for part of the sample inflammation-related symptoms, including fever and infectious or inflammatory disease; uncontrolled systemic disease; uncontrolled metabolic disease or other significant uncontrolled somatic disorder known to affect mood; somatic medications known to affect mood or the immune system, such as corticosteroids, non-steroid anti-inflammatory drugs and statins For healthy controls: mental retardation, pregnancy, major medical/neurological or psychiatric disorders.



Stanford University FAA	MDD	<p>For patients: women aged 18–55 years currently experiencing a diagnosable depressive episode (established using DSM-IV-TR criteria assessed with the Structured Clinical Interview for DSM Axis I)</p> <p>For healthy controls: women aged 18–55 years not meeting criteria for any past or current Axis I disorder (established using DSM-IV-TR criteria assessed with the Structured Clinical Interview for DSM Axis I)</p>	<p>For patients: Cardiovascular or neurological disorders, history of head injury, hyper- or hypothyroidism, lifetime diagnosis of bipolar disorder or psychosis; substance abuse/dependence in the past six months</p> <p>For healthy controls: Cardiovascular or neurological disorders, history of head injury, hyper- or hypothyroidism, history of any psychiatric disorder or substance abuse/dependence</p>
STRADL	MDD	<p>For patients: MDD diagnosis</p> <p>For healthy controls: No psychological disorders</p>	<p>For patients: neurological disorders</p> <p>For healthy controls: psychological or neurological disorders</p>
Sydney Bipolar Kids and Siblings	BD	<p>For patients: Any DSM-IV diagnosis</p>	<p>For healthy controls: parent or sibling with bipolar I or II, recurrent MDD, schizoaffective disorder, schizophrenia, recurrent substance abuse, psychiatric hospitalisation. Parent with a first degree relative who had a past mood disorder hospitalisation or history of psychosis</p>
Sydney Brain and Mind Centre	MDD + OCD + BD + Psychosis + GAD + SAD	<p>For patients: Age 12 - 25 and presented with an affective, psychotic or developmental/behavioural syndrome.</p>	<p>Exclusion criteria for all subjects included medical instability (as determined by a psychiatrist), history of neurological disease (e.g. tumour, head trauma, epilepsy), medical illness known to impact cognitive and brain function (e.g. cancer), intellectual and/or developmental disability and insufficient English for neuropsychological assessment. All subjects were asked to abstain from drug or alcohol use for 48 hours prior to testing and informed about a drug screen protocol.</p>
TIPS-Jena	MDD	<p>For individuals with MDD: presence of a DSM-IV diagnosis using the German version of the Structured Clinical Interview for DSM-IV Axis I Disorders (SKID-I) that were confirmed by two psychiatrists.</p> <p>For healthy controls: participants did not have current or history of a DSM-IV axis I psychiatric disorder, a first-degree relative with a known psychiatric disorder or use of psychotropic medication.</p>	<p>For both individuals with MDD and healthy controls: exclusion criteria were history of major neurological and untreated major general medical conditions, and IQ lower than 80.</p>
UCSF Adolescent MDD	MDD	<p>For patients: presence of a primary diagnosis of MDD as determined via Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (Kaufman et al., 1997). For healthy control</p>	<p>Exclusion criteria for all participants included: 1) use of pharmacotherapeutics for treating psychiatric conditions within the past 6 months, 2) misuse of drugs within two months prior to</p>

		<p>participants absence of any psychiatric diagnosis as determined by the Diagnostic Interview Schedule for Children (Shaffer et al., 2000) and Diagnostic Predictive Scale (Lucas et al., 2001) instruments.</p>	<p>MRI scanning; 3) two or more alcoholic drinks per week within the previous month (as assessed by the Customary Drinking and Drug Use Record; CDDR) (Brown et al, 1998); 4) a full scale IQ score of less than 75 (as assessed by the Wechsler Abbreviated Scale of Intelligence; WASI) (Wechsler, 1999); 5) contraindications for MRI including ferromagnetic implants and claustrophobia; 6) pregnancy or the possibility of pregnancy; 7) left-handedness; 8) prepubertal status (as assessed as Tanner stages of 1 or 2) (Tanner, 1962); 9) inability to understand and comply with procedures; 10) neurological disorder (including meningitis, migraine, or HIV); 11) head trauma; 12) learning disability; 13) serious health problems; and 14) complicated or premature birth (i.e., birth before 33 weeks of gestation). The MDD group was subject to the additional exclusion criterion of a primary psychiatric diagnosis other than MDD. The healthy control group was subject to the additional exclusion criteria of: 1) history of mood or psychotic disorders in a first- or second-degree relative (as assessed by the Family Interview for Genetics; FIGS) (Maxwell, 1992); and 2) current or lifetime DSM-IV-TR Axis I psychiatric disorder.</p>
University of Minnesota	MDD	<p>For patients: primary diagnosis of MDD and had not received any psychotropic medication treatment for the past 2 months. HC participants were eligible if they had no current or past psychiatric diagnoses.</p>	<p>Exclusion criteria for included the presence of a neurologic or other chronic medical condition, mental retardation, pervasive developmental disorder, substance use disorder, bipolar disorder, or schizophrenia</p>
University of Texas - Austin - Bipolar Seed Program	BD	<p>For patients: Meeting diagnostic criteria for bipolar disorder type I per the Structured Clinical Interview for DSM-5- Research Version, between 18-25 years of age</p>	<p>For patients: History of major medical illness or significant head trauma resulting in possible neurological or central nervous system damage, Past or current severe alcohol or other substance use disorder, Full scale IQ &lt;85, Positive pregnancy test, Having a medical condition/previous surgery preventing participation in a magnetic resonance imaging scan Not taking medications for greater than or equal to 4 weeks (i.e., participants must be stable on their medications)</p>

			For healthy controls: Meeting criteria for psychotic, affective or other anxiety disorders per the Structured Clinical Interview for DSM-5-Research Version, History of major medical illness or significant head trauma resulting in possible neurological or central nervous system damage, Past or current severe alcohol or other substance use disorder, Full scale IQ <85, Positive pregnancy test, Having a medical condition/previous surgery preventing participation in a magnetic resonance imaging scan
University of Washington/ Harvard	MDD + PTSD + GAD + Panic disorder + Social Anxiety Disorder	For all participants: 8-20 years of age, English speaking, with and without exposure to trauma. For patients: Endorsement of ideation or behaviour on the SITBI	Exclusion criteria for all participants: Psychiatric medication use (stimulants for ADHD were discontinued for the scan), MRI contraindications, active substance dependence, pervasive developmental disorder. For healthy controls: Endorsement of ideation or behaviour on the SITBI
Yale School of Medicine	BD, MDD	For individuals with MDD and BD: ages 13-25 years and meeting criteria for major depressive disorder (MDD) or bipolar disorder (BD) type 1 or type 2 according to the DSM-IV.	Exclusion criteria for all individuals: history of or current medical or neurological condition that could affect the brain (except treated hypothyroidism), and magnetic resonance imaging (MRI) contraindications.

Table S2. Scanner type and DWI acquisition parameters per site

Site	Scanner vendor and type	Voxel size and slice thickness	Number of directions & b-factor	Preprocessing software & steps
BiDirect	3T Philips Intera	0.94 x 0.94 x 3.6 mm (reconstructed)	20 directions; b-factor=1000	Eddy correction, DTIfit in FSL
Brazilian High Risk Cohort Porto Alegre	GE Signa HD 1.5-T And GE Signa HDx 1.5T	3 x 3 x 3 mm	15 directions; b-factor=800 s/mm <sup>2</sup>	Eddy current correction and DTIfit in FSL
Chiba University	3T GE Discovery MR750	1.875 x 1.875 x 2 mm	30 directions; b-factor=1000	Eddy correction, DTIfit in FSL
CHU Montpellier BICS	3T Siemens Magnetom SKYRA	0.6 x 0.6 x 2 mm	64 directions; b-factor=1000	Denosing, distortion correction (SynB0), TOPUP, Eddy, and DTIfit in FSL
CHU Montpellier IMPACT	3T Siemens Magnetom SKYRA	2.5 x 2.5 x 2.5 mm	30 directions; b-factor=1000	Denosing, distortion correction (SynB0), TOPUP, Eddy, and DTIfit in FSL
CHU Montpellier UF8555	1.5T Siemens Magnetom Avanto	2.5 x 2.5 x 2.5 mm	64 directions, b-factor=1000	Denosing, distortion correction (SynB0), TOPUP, Eddy, and DTIfit in FSL
DCHS	3T Siemens Magnetom Allegra Sygno	2 x 2 x 2 mm	45 directions; b-factor=1000	Eddy, TOPUP, DTIfit in FSL
MOODS team DEP-ARREST- CLIN	3T Philips Achieva	2 x 2 x 2 mm	60 directions b-factor=1500	Eddy correction and DTIfit in FSL
EPISCA (Leiden adolescents)	3T Philips Achieva	2.3 x 2.3 x 2.3 mm	32 directions; 1 b0 image; b- factor=1000	Eddy current and motion correction in FSL
ETPB-STB	3T GE HDx	1.875 x 1.875 x 2.5 mm	30 directions; b-factor=1100	TORTOISE
FSL_NPL- OCD sample	3T Siemens Magnetom Allegra	1.8 x 1.8 1.8 mm	30 directions; 2 b0 images, b- factor=1000	Eddy correction and DTIfit in FSL
FSL_NPL- SZ sample	3T Siemens Magnetom Allegra	1.8 x 1.8 1.8 mm	30 directions; 2 b0 images, b- factor=1000	Eddy correction and DTIfit in FSL
FOR2107 Marburg	3T Siemens Magnetom Trio	2.5 x 2.5 x 2.5 mm	30 directions; 4 b0 images; b- factor=1000	Eddy correction and DTIfit in FSL

Site	Scanner vendor and type	Voxel size and slice thickness	Number of directions & b-factor	Preprocessing software & steps
FOR2107 Muenster	3T Siemens PRISMA	2.5 x 2.5 x 2.5 mm	30 directions; 4 b0 images; b-factor=1000	Eddy correction and DTIfit in FSL
Ghent	3T Siemens Magnetom Trio Tim	2 x 2 x 2 mm	12 directions; b-factor=850	Eddy and DTIfit in FSL
GIPSI BD	3T Philips Ingenia	1.75 x 1.75 x 2 mm	16 directions; 1 b0 images, b-factor=1000	FSL
GIPSI SZ	3T Philips Ingenia	1.75 x 1.75 x 2 mm	16 directions; 1 b0 images, b-factor=1000	FSL
Grady Trauma Project Emory University	3T Siemens TIM-Trio	2 x 2 x 2 mm	138 directions; 6 b0 images; b-factor=1000	Eddy current and motion correction in FSL
Halifax	1.5T GE Signa	1.875 x 1.875 x 2 mm	31 directions; 2 b0 images; b-factor=900	Eddy current correction in FSL
Houston BD	3T Philips Ingenia	1 x 1 x 1 mm	42 directions; b-factor=1000	Eddy currents, BET, DTIfit in FSL
Jena-SB	3T Siemens Tim Trio	1.25 x 1.25 x 2.5 mm <sup>3</sup>	70 directions; $b = 1000$ s/mm <sup>2</sup> ; six $b = 0$ images	Denosing step, Eddy and DTIfit in FSL
KASP	3T GE	0.94 x 0.94 x 2.9 mm	60 directions; 10 b) images; b-factor=1000	Denosing, Gibbs ringing correction, Eddy currents, motion and DTIfit in FSL
LTS Colorado site 1	3T Siemens Tim Trio	2 x 2 x 2 mm	173 directions; b-factor=3000	FSL Topup
LTS Colorado site 2	3T Siemens Prisma	2 x 2 x 2 mm	173 directions; b-factor=3000	FSL Topup
McGill University	3T Siemens Trio	2 x 2 x 2 mm	64 directions; b-factor=1000	Eddy current correction, TOPUP
MR-IMPACT	3T Magnetom Trio Tim	2 x 2 x 2 mm	13 direction; 5 b0 images; b-factor=350/1600[62]	Eddy in FSL
Muenster Neuroimaging Cohort	3T Philips Gyroscan Intera	1.8 x 1.8 x 3.6 mm	25 directions; 5 b0 images; b-factor=1000	Eddy current & motion correction in ACID Toolbox
PAFIP	3T Philips Achieva	2 x 2 x 2 mm	64 directions; b-factor=1300	Eddy current correction in FSL

Site	Scanner vendor and type	Voxel size and slice thickness	Number of directions & b-factor	Preprocessing software & steps
San Raffaele Hospital	3T Philips Intera / 3T Philips Ingenia	2 x 2 x 2 mm	35 directions; 1 b0 image; b-factor=900/1000	Eddy correction, BET in FSL
Stanford University FAA	3T GE Discovery MR750	2 x 2 x 2 mm	96 directions; 9b0 images; b-factor=2000	Eddy current and motion correction in FSL
STRADL site Aberdeen	3T Philips Achieva-TX	2.3 x 2.3 x 2.3 mm	64 directions; b-factor=1200	Eddy in FSL
STRADL site Dundee	3T Siemens Prisma-FIT	2.3 x 2.3 x 2.3 mm	64 directions; b-factor=1200	Eddy in FSL
Sydney Bipolar Kids and Siblings	3T Philips Achieva	1 x 2.5 x 2.5 mm	32 directions; b-factor=1000	Pre-processing using FDT in FSL
Sydney Brain and Mind Centre	3T GE Discovery MR750	0.86 x 0.86 x 2 mm	69 directions; 8 b0 images; b-factor=1159	Eddy current & motion correction in FSL
TIPS-Jena	3T Siemens Prisma Fit	1 x 1 x 1 mm	54 directions; 7 b0 images; b-factor=1200	FSL Topup based pipeline
UCSF Adolescent MDD	3T GE Discovery MR750	1.875 x 1.875 x 2.5 mm	30 directions; 1 b0 image; b-factor=1500	Fieldmap correction, eddy current correction, matrix rotation in FSL & AFNI
University of Minnesota Adolescent MDD	3T Siemens Trio	2 x 2 x 2 mm	64 directions; b-factor=1000	Eddy current and distortion correction in FSL
University of Texas - Austin Bipolar Seed Program	3T Siemens Skyra	2 x 2 x 2 mm	30 directions; b-factor=1000	Eddy correction, DTIfit in FSL
University of Washington/ Harvard	3T Philips Achieva	2 x 2 x 2 mm	32 directions; b-factor=1200	Eddy correction in FSL
Yale School of Medicine	3T Siemens Trio	3 x 3 x 3 mm	32 directions; b-factor=1000	Eddy current correction in FSL

Table S3: List of white matter tracts of interest

<b>Abbreviation</b>	<b>Full tract name</b>
AverageFA	Full skeleton average FA
ACR (L+R)	Anterior corona radiata
ALIC (L+R)	Anterior limb of internal capsule
BCC	Body of corpus callosum
CC (BCC+GCC+SCC)	Corpus callosum
CGC (L+R)	Cingulum (cingulate gyrus)
CGH (L+R)	Cingulum (hippocampal portion)
CR (L+R)	Corona radiata
CST (L+R)	Corticospinal tract
EC (L+R)	External capsule
FX	Fornix

FXST (L+R)	Fornix (cres) / Stria terminalis
GCC	Genu of corpus callosum
IC (L+R)	Internal capsule
IFO (L+R)	Inferior fronto-occipital fasciculus
PCR (L+R)	Posterior corona radiata
PLIC (L+R)	Posterior limb of internal capsule
PTR (L+R)	Posterior thalamic radiation
RLIC (L+R)	Retrolemniscal part of internal capsule
SCC	Splenium of corpus callosum
SCR (L+R)	Superior corona radiata
SFO (L+R)	Superior fronto-occipital fasciculus
SLF (L+R)	Superior longitudinal fasciculus
SS (L+R)	Sagittal stratum



UNC (L+R)	Uncinate fasciculus
-----------	---------------------

Table S4. Definitions used for recent suicidal ideation and history of suicide attempt

Site	Definition recent suicidal ideation	Definition history of suicide attempt
BiDIRECT	Hamilton Depression Rating Scale suicidal ideation item scored 2 (“Wishes to be dead or thinks about possible death”) or 3 (“Ideas or gestures of suicide”)	Clinical interview: number of lifetime suicide attempts > 0
Brazilian High Risk Cohort Porto Allegre	CBCL item 91 (“Talks about killing oneself in past six months”) is scored 1 (“Somewhat or sometimes true”) or 2 (“very true or often true”)	NA
Chiba University	Beck Depression Inventory item on suicidal ideation scored 1 (“I think about ending my life but would not do it”), 2 (“I would like to end my life”), 3 (“I would kill myself if I had the opportunity”)	NA
CHU Montpellier BICS	Columbia Suicide Severity Rating Scale (C-SSRS) positive answer on one or more of the following items current non-specific active suicidal ideation, current active suicidal ideation with any method (not plan) without intent to act, current active suicidal ideation with some intent without specific plan, current active suicidal ideation with specific plan and intent	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime actual suicide attempt answered “yes”
CHU Montpellier IMPACT	Inventory of Depressive Symptomatology (Clinician-related; IDS-C) suicidal ideation item scored 2 (thinks of suicide or death multiple times per week for multiple minutes) or 3 (seriously thinks about suicide or death many times per day, or has prepared suicide or has tried to end life) OR Quick Inventory of Depression Symptomatology suicidal ideation item scored 2 (“I think about suicide or death multiple times per week for multiple minutes”) or 3 (“I think about suicide or death multiple times per day, or I have prepared suicide or I have tried to end my life”)	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime actual suicide attempt answered “yes”
CHU Montpellier UF8555	Scale for Suicidal Ideation (SSI) item: 4 (“Active Suicidal Ideation”) scored 1 (“weak”) OR 2 (“medium to strong) OR SSI item 5 (“passive suicidal thoughts) scored 2 (“would avoid doing what is necessary to stay alive; such as ending diabetes treatment”) OR Beck Depression Inventory item on suicidal ideation scored 1 (“I think about ending my life but would not do it”), 2 (“I would like to end my life”), 3 (“I would kill myself if I had the opportunity”) OR Hamilton Depression Rating Scale suicidal ideation item scored 2 (“Wishes to be dead or thinks about possible death”) or 3 (“Ideas or gestures of suicide”)	Lifetime history of suicide attempt determined using an interview
DCHS	NA	Mini International Neuropsychiatric interview (MINI) question BQ14 (“Did you ever make a suicide attempt?”) scored 4 (“yes”)
MOODS team DEP-ARREST-	Hamilton Depression Rating Scale suicidal ideation item scored 2 (“Wishes to be dead or thinks about possible death”) or 3 (“Ideas or gestures of suicide”)	Recent OR past suicide attempt determined in an interview

CLIN		
EPISCA (Leiden adolescents)	Childhood Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it") or 2 ("I want to end my life")	NA
ETPB-STB	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 ("Ideas or gestures of suicide") OR Montgomery-Asberg Depression Rating Scale (MADRS) item on suicidal ideation scored between 2 ("Weary of life, only fleeting suicidal thoughts") and 6 ("Explicit plans for suicide when there is an opportunity. Active preparations for suicide") OR Scale for Suicidal Ideation item 4 ("desire to end life") scored "weak" or "moderate to severe"	Lifetime history of suicide attempt determined using an interview
FSL_NPL-OCD sample	Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 ("Ideas or gestures of suicide")	Clinical interview: number of lifetime suicide attempts > 0
FSL_NPL - SZ sample	NA	Lifetime history of suicide attempt determined using an interview
FOR2107 Marburg	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale (HDRS) suicidal ideation item scored 2 "Wishes to be dead or thinks about possible death" or 3 ("Ideas or gestures of suicide")	NA
FOR2107 Muenster	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale (HDRS-21) suicidal ideation item scored 2 "Wishes to be dead or thinks about possible death" or 3 "Ideas or gestures of suicide"	NA
Ghent	NA	Lifetime history of suicide attempt determined using an interview
GIPSI BD	NA	Lifetime history of suicide attempt determined using an interview
GIPSI SZ	NA	Lifetime history of suicide attempt determined using an interview

Grady Trauma Project Emory University	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity")	SSH item 1 (Have you ever tried to kill yourself or commit suicide?) scored "yes"
Halifax	NA	Lifetime history of suicide attempt determined using an interview
Houston BD	NA	Clinical interview: number of lifetime suicide attempts > 0
Jena-SB	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 (Ideas or gestures of suicide)	NA
KASP	Calgary Depression Scale for Schizophrenia item 8 ("In the past two weeks, have you felt that life wasn't worth living? Did you feel like ending it all? What did you think you might do?") scored 1("Frequent thoughts or being better off dead or occasional thoughts or suicide") or higher	NA
LTS Colorado	NA	Lifetime history of suicide attempt determined using an interview
McGill University	Columbia Suicide Severity Rating Scale (C-SSRS) positive answer on one or more of the following items: current non-specific active suicidal ideation, current active suicidal ideation with any method (not plan) without intent to act, current active suicidal ideation with some intent without specific plan, current active suicidal ideation with specific plan and intent	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime actual suicide attempt answered "yes"
MR-IMPACT	Columbia Suicide Severity Rating Scale (C-SSRS) positive answer on one or more of the following items (in the past month): current non-specific active suicidal ideation, current active suicidal ideation with any method (not plan) without intent to act, current active suicidal ideation with some intent without specific plan, current active suicidal ideation with specific plan and intent	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime actual suicide attempt answered "yes"
Muenster Neuroimaging Cohort	Beck Depression Inventory-2 item on suicidal ideation scored 1 ("I have thoughts of killing myself, but I would not carry them out"), 2 ("I would like to kill myself"), 3 ("I would kill myself if I had the chance") OR Montgomery-Asberg Depression Rating Scale (MADRS) item on suicidal ideation scored between 2 ("Weary of life, only fleeting suicidal thoughts") and 6 ("Explicit plans for suicide when there is an opportunity. Active preparations for suicide")	NA

PAFIP	NA	Lifetime history of suicide attempt determined using an interview
San Raffaele Hospital	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 ("Ideas or gestures of suicide")	Clinical interview: number of lifetime suicide attempts > 0
Stanford University FAA	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 ("Ideas or gestures of suicide")	NA
STRADL	SCID item A21 ("Suicidal ideation during current major depressive episode") scored 1 ("yes") OR Quick Inventory of Depressive Symptomatology item 12 ("Thoughts of suicide or death") is scored 2 ("I think of suicide or death several times a week for several minutes") or higher	NA
Sydney Bipolar Kids and Siblings	NA	Diagnostic Interview for Genetic Studies (DIGS) question on lifetime suicide attempt answered "yes" OR Kiddie Schedule for Affective Disorders and Schizophrenia (KSADS) item on lifetime suicide attempt coded as "yes"
Sydney Brain and Mind Centre	Hamilton Depression Rating Scale suicidal ideation item scored 2 "Wishes to be dead or thinks about possible death" or 3 "Ideas or gestures of suicide"	Clinical interview: lifetime suicide attempt coded "yes"
TIPS-Jena	Beck Depression Inventory item on suicidal ideation scored 1 ("I think about ending my life but would not do it"), 2 ("I would like to end my life"), 3 ("I would kill myself if I had the opportunity") OR Hamilton Depression Rating Scale suicidal ideation item scored 2 ("Wishes to be dead or thinks about possible death") or 3 ("Ideas or gestures of suicide") OR SCID item on current suicidal ideation scored ("yes")	NA
UCSF Adolescent MDD	Columbia Suicide Severity Rating Scale (C-SSRS) positive answer on one or more of the following items (in the past week): current non-specific active suicidal ideation, current active suicidal ideation with any method (not plan) without intent to act, current active suicidal ideation with some intent without specific plan, current active suicidal ideation with specific plan and intent	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime non-interrupted attempt answered "yes"

University of Minnesota Adolescent MDD	Beck Depression Inventory-2 item on suicidal ideation scored 1 (“I have thoughts of killing myself, but I would not carry them out”), 2 (“I would like to kill myself”), 3 (“I would kill myself if I had the chance”) OR Childrens Depression Rating Scale 3 or higher	Kiddie Schedule for Affective Disorders and Schizophrenia for School Aged Children- Lifetime version (K-SADS-PL_2009) item on suicide attempt answer 1
University of Texas - Austin - Mood Addiction Interaction Neuroscience (MAIN) Program	Structured Clinical Interview DSM-5 (SCID-RV/NP for DSM-5) Suicidality module: answer “yes” to question on ideation in the past week (“Have you in the past week wished you were dead or wished you could go to sleep and not wake up?”)	Structured Clinical Interview DSM-5 (SCID-RV/NP for DSM-5) Suicidality module: answer “yes” to question “Have you ever tried to kill yourself?”
University of Washington/ Harvard	NA	Self-injurious Thoughts and Behaviors Interview (SITBI) measure on lifetime suicide attempt answered “yes”
Yale School of Medicine	Columbia Suicide Severity Rating Scale (C-SSRS) positive answer on one or more of the following items (in the past month): current non-specific active suicidal ideation, current active suicidal ideation with any method (not plan) without intent to act, current active suicidal ideation with some intent without specific plan, current active suicidal ideation with specific plan and intent	Columbia Suicide Severity Rating Scale (C-SSRS) question on lifetime non-interrupted suicide attempt answered “yes”

Table S5. Differences in regional AD between individuals with a history of suicide attempt and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	HC
ACR	-0.070	0.060	-0.188	0.048	0.245	0.557	601	511
ALIC	-0.080	0.060	-0.198	0.038	0.184	0.557	600	511
BCC	0.053	0.060	-0.065	0.172	0.377	0.674	599	507
CC	0.016	0.060	-0.102	0.134	0.788	0.946	600	508
CGC	-0.101	0.060	-0.219	0.016	0.092	0.557	604	512
CGH	0.008	0.060	-0.110	0.126	0.899	0.962	600	512
CR	-0.061	0.060	-0.179	0.057	0.311	0.649	602	511
CST	0.031	0.060	-0.087	0.149	0.606	0.891	602	510
EC	-0.055	0.060	-0.173	0.063	0.364	0.674	601	512
FX	0.033	0.060	-0.085	0.151	0.582	0.891	603	512
FXST	0.001	0.060	-0.117	0.119	0.986	0.986	604	512
GCC	-0.016	0.060	-0.134	0.102	0.795	0.946	602	509
IC	-0.077	0.060	-0.195	0.041	0.205	0.557	600	510
IFO	-0.085	0.061	-0.205	0.036	0.170	0.557	598	475
PCR	-0.075	0.060	-0.193	0.043	0.212	0.557	602	509
PLIC	-0.043	0.060	-0.161	0.074	0.471	0.785	602	511
PTR	0.006	0.060	-0.112	0.124	0.923	0.962	603	511
RLIC	-0.080	0.060	-0.198	0.038	0.184	0.557	603	512
SCC	0.024	0.060	-0.094	0.142	0.691	0.946	600	509
SCR	-0.071	0.060	-0.189	0.047	0.242	0.557	601	510
SFO	-0.011	0.060	-0.129	0.107	0.854	0.962	601	511
SLF	-0.098	0.060	-0.216	0.020	0.105	0.557	604	512
SS	-0.016	0.060	-0.134	0.102	0.795	0.946	601	511
UNC	0.154	0.063	0.030	0.277	0.015	0.375	582	447
AverageAD	-0.081	0.060	-0.199	0.037	0.181	0.557	603	511

Table S6. Differences in regional FA between individuals with a history of suicide attempt and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	HC
ACR	-0.094	0.057	-0.206	0.018	0.101	0.127	636	591
ALIC	-0.144	0.057	-0.256	-0.032	0.012	0.022	635	591
BCC	-0.182	0.057	-0.295	-0.069	0.002	0.005	630	588
CC	-0.223	0.057	-0.336	-0.111	0.000	0.001	631	591
CGC	-0.167	0.057	-0.279	-0.055	0.004	0.008	638	589
CGH	-0.040	0.057	-0.152	0.072	0.488	0.530	635	590
CR	-0.169	0.057	-0.281	-0.056	0.003	0.008	636	590
CST	-0.011	0.057	-0.123	0.101	0.849	0.849	635	588
EC	-0.168	0.057	-0.280	-0.056	0.003	0.008	637	591
FX	-0.058	0.057	-0.170	0.055	0.315	0.358	632	589
FXST	-0.201	0.057	-0.313	-0.089	0.000	0.002	636	590
GCC	-0.200	0.057	-0.313	-0.088	0.000	0.002	635	590
IC	-0.116	0.057	-0.228	-0.003	0.044	0.058	636	589
IFO	-0.130	0.058	-0.245	-0.015	0.027	0.040	633	546
PCR	-0.223	0.057	-0.335	-0.110	0.000	0.001	638	591
PLIC	-0.029	0.057	-0.141	0.083	0.609	0.634	635	590
PTR	-0.171	0.057	-0.283	-0.058	0.003	0.008	632	590
RLIC	-0.125	0.057	-0.238	-0.013	0.029	0.040	634	591
SCC	-0.202	0.057	-0.315	-0.090	0.000	0.002	632	589
SCR	-0.139	0.057	-0.251	-0.027	0.015	0.026	635	591
SFO	-0.086	0.057	-0.199	0.026	0.133	0.158	633	588
SLF	-0.226	0.057	-0.339	-0.113	0.000	0.001	633	590
SS	-0.127	0.057	-0.239	-0.015	0.027	0.040	635	591
UNC	-0.172	0.059	-0.288	-0.055	0.004	0.008	616	526
AverageFA	-0.195	0.057	-0.307	-0.083	0.001	0.003	635	589



Table S7. Differences in regional MD between individuals with a history of suicide attempt and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	HC
ACR	0.050	0.059	-0.065	0.165	0.392	0.577	614	553
ALIC	0.002	0.059	-0.114	0.117	0.979	0.997	611	550
BCC	0.141	0.059	0.025	0.256	0.017	0.086	610	549
CC	0.141	0.059	0.025	0.256	0.017	0.086	614	550
CGC	-0.013	0.059	-0.128	0.102	0.831	0.997	614	552
CGH	-0.057	0.059	-0.173	0.058	0.330	0.516	612	551
CR	0.094	0.059	-0.021	0.209	0.109	0.272	615	551
CST	0.000	0.059	-0.115	0.115	0.997	0.997	612	552
EC	0.060	0.059	-0.055	0.176	0.305	0.509	613	550
FX	0.077	0.059	-0.038	0.192	0.192	0.395	613	547
FXST	0.143	0.059	0.028	0.258	0.015	0.086	614	552
GCC	0.096	0.059	-0.018	0.211	0.101	0.272	617	553
IC	0.005	0.059	-0.110	0.120	0.934	0.997	611	550
IFO	-0.020	0.060	-0.138	0.098	0.737	0.969	608	508
PCR	0.102	0.059	-0.013	0.217	0.084	0.272	615	550
PLIC	-0.010	0.059	-0.126	0.105	0.862	0.997	609	549
PTR	0.099	0.059	-0.016	0.214	0.094	0.272	615	551
RLIC	0.034	0.059	-0.081	0.149	0.559	0.777	616	551
SCC	0.164	0.059	0.049	0.280	0.005	0.067	614	550
SCR	0.098	0.059	-0.017	0.213	0.095	0.272	616	551
SFO	0.007	0.059	-0.108	0.122	0.905	0.997	611	554
SLF	0.084	0.059	-0.031	0.200	0.152	0.346	612	551
SS	0.075	0.059	-0.040	0.190	0.205	0.395	612	553
UNC	0.220	0.061	0.099	0.340	0.000	0.009	591	487
AverageMD	0.062	0.059	-0.053	0.177	0.290	0.509	615	549

Table S8. Differences in regional RD between individuals with a history of suicide attempt and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	HC
ACR	0.096	0.059	-0.019	0.211	0.102	0.175	613	553
ALIC	0.050	0.059	-0.065	0.166	0.392	0.515	610	551
BCC	0.153	0.059	0.037	0.268	0.010	0.036	610	549
CC	0.152	0.059	0.036	0.267	0.010	0.036	612	552
CGC	0.048	0.059	-0.067	0.163	0.412	0.515	614	552
CGH	-0.027	0.059	-0.142	0.088	0.652	0.741	613	553
CR	0.148	0.059	0.033	0.263	0.012	0.037	614	552
CST	-0.011	0.059	-0.126	0.104	0.848	0.884	611	551
EC	0.086	0.059	-0.029	0.201	0.143	0.224	614	553
FX	0.104	0.059	-0.011	0.220	0.077	0.175	612	545
FXST	0.154	0.059	0.039	0.269	0.009	0.036	616	553
GCC	0.117	0.059	0.002	0.232	0.047	0.118	613	553
IC	0.059	0.059	-0.056	0.174	0.318	0.442	613	551
IFO	0.029	0.060	-0.088	0.147	0.627	0.741	612	508
PCR	0.168	0.059	0.053	0.283	0.004	0.036	617	551
PLIC	-0.008	0.059	-0.124	0.107	0.886	0.886	610	552
PTR	0.101	0.059	-0.014	0.216	0.086	0.175	614	552
RLIC	0.082	0.059	-0.033	0.197	0.162	0.238	615	553
SCC	0.166	0.059	0.050	0.281	0.005	0.036	613	550
SCR	0.137	0.059	0.021	0.252	0.020	0.057	613	552
SFO	0.017	0.059	-0.098	0.132	0.769	0.836	612	553
SLF	0.161	0.059	0.046	0.277	0.006	0.036	614	552
SS	0.095	0.059	-0.020	0.210	0.105	0.175	613	554
UNC	0.170	0.061	0.050	0.290	0.006	0.036	594	487
AverageRD	0.098	0.059	-0.017	0.213	0.096	0.175	614	552

Table S9. Differences in regional AD between individuals with a history of suicide attempt and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	CLC
ACR	0.010	0.048	-0.083	0.103	0.836	0.968	601	1680
ALIC	0.024	0.048	-0.069	0.117	0.615	0.968	600	1677
BCC	0.049	0.048	-0.044	0.142	0.304	0.968	599	1679
CC	0.038	0.048	-0.055	0.132	0.421	0.968	600	1676
CGC	0.007	0.047	-0.086	0.100	0.886	0.968	604	1683
CGH	-0.009	0.048	-0.102	0.085	0.856	0.968	600	1682
CR	0.029	0.048	-0.064	0.123	0.537	0.968	602	1676
CST	0.018	0.048	-0.075	0.111	0.709	0.968	602	1678
EC	0.011	0.048	-0.082	0.104	0.814	0.968	601	1669
FX	-0.011	0.047	-0.104	0.082	0.811	0.968	603	1685
FXST	-0.016	0.047	-0.109	0.077	0.733	0.968	604	1683
GCC	0.024	0.048	-0.069	0.118	0.608	0.968	602	1680
IC	-0.002	0.048	-0.095	0.091	0.966	0.968	600	1675
IFO	-0.042	0.048	-0.135	0.052	0.380	0.968	598	1661
PCR	0.002	0.048	-0.091	0.095	0.968	0.968	602	1675
PLIC	0.014	0.048	-0.079	0.107	0.768	0.968	602	1678
PTR	0.021	0.047	-0.072	0.114	0.665	0.968	603	1677
RLIC	-0.043	0.047	-0.136	0.050	0.363	0.968	603	1676
SCC	0.017	0.048	-0.076	0.111	0.716	0.968	600	1678
SCR	0.035	0.048	-0.058	0.128	0.464	0.968	601	1671
SFO	0.043	0.048	-0.051	0.136	0.372	0.968	601	1673
SLF	0.002	0.047	-0.091	0.095	0.960	0.968	604	1680
SS	-0.013	0.048	-0.106	0.080	0.786	0.968	601	1676
UNC	0.101	0.048	0.006	0.196	0.037	0.922	582	1615
AverageAD	-0.016	0.048	-0.110	0.077	0.729	0.968	603	1672

Table S10. Differences in regional FA between individuals with a history of suicide attempt and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	CLC
ACR	-0.102	0.046	-0.192	-0.011	0.028	0.071	636	1802
ALIC	-0.094	0.046	-0.185	-0.004	0.041	0.080	635	1794
BCC	-0.066	0.046	-0.157	0.024	0.153	0.201	630	1795
CC	-0.104	0.046	-0.195	-0.013	0.025	0.071	631	1792
CGC	-0.099	0.046	-0.189	-0.009	0.032	0.071	638	1797
CGH	-0.005	0.046	-0.096	0.085	0.912	0.956	635	1797
CR	-0.126	0.046	-0.217	-0.036	0.006	0.034	636	1795
CST	-0.002	0.046	-0.092	0.089	0.969	0.969	635	1797
EC	-0.067	0.046	-0.158	0.023	0.145	0.201	637	1792
FX	-0.032	0.046	-0.122	0.059	0.495	0.590	632	1792
FXST	-0.126	0.046	-0.217	-0.036	0.006	0.034	636	1795
GCC	-0.061	0.046	-0.151	0.030	0.190	0.237	635	1801
IC	-0.088	0.046	-0.178	0.003	0.057	0.096	636	1794
IFO	0.005	0.046	-0.086	0.096	0.918	0.956	633	1770
PCR	-0.138	0.046	-0.228	-0.047	0.003	0.034	638	1797
PLIC	-0.015	0.046	-0.106	0.075	0.744	0.846	635	1793
PTR	-0.124	0.046	-0.214	-0.033	0.008	0.034	632	1797
RLIC	-0.100	0.046	-0.190	-0.009	0.031	0.071	634	1792
SCC	-0.123	0.046	-0.214	-0.032	0.008	0.034	632	1791
SCR	-0.077	0.046	-0.167	0.014	0.098	0.153	635	1798
SFO	-0.098	0.046	-0.189	-0.007	0.034	0.071	633	1799
SLF	-0.125	0.046	-0.216	-0.034	0.007	0.034	633	1802
SS	-0.107	0.046	-0.198	-0.017	0.020	0.071	635	1794
UNC	-0.069	0.047	-0.161	0.023	0.139	0.201	616	1732
AverageFA	-0.093	0.046	-0.183	-0.002	0.045	0.080	635	1793

Table S11. Differences in regional MD between individuals with a history of suicide attempt and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	CLC
ACR	0.070	0.047	-0.021	0.162	0.133	0.278	614	1764
ALIC	0.046	0.047	-0.046	0.138	0.330	0.551	611	1759
BCC	0.081	0.047	-0.012	0.173	0.087	0.199	610	1760
CC	0.080	0.047	-0.012	0.172	0.088	0.199	614	1764
CGC	0.036	0.047	-0.056	0.128	0.442	0.631	614	1764
CGH	-0.032	0.047	-0.124	0.060	0.492	0.638	612	1764
CR	0.100	0.047	0.009	0.192	0.032	0.199	615	1762
CST	0.000	0.047	-0.092	0.092	0.993	0.993	612	1750
EC	0.030	0.047	-0.062	0.122	0.521	0.638	613	1760
FX	0.015	0.047	-0.077	0.107	0.747	0.778	613	1755
FXST	0.092	0.047	0.001	0.184	0.049	0.199	614	1759
GCC	0.035	0.047	-0.057	0.127	0.454	0.631	617	1767
IC	0.027	0.047	-0.065	0.119	0.571	0.648	611	1761
IFO	-0.067	0.047	-0.159	0.026	0.158	0.304	608	1729
PCR	0.086	0.047	-0.005	0.178	0.066	0.199	615	1757
PLIC	0.029	0.047	-0.063	0.121	0.536	0.638	609	1762
PTR	0.081	0.047	-0.011	0.173	0.085	0.199	615	1762
RLIC	0.025	0.047	-0.067	0.117	0.596	0.648	616	1764
SCC	0.088	0.047	-0.004	0.180	0.062	0.199	614	1754
SCR	0.113	0.047	0.021	0.204	0.016	0.199	616	1760
SFO	0.102	0.047	0.010	0.194	0.030	0.199	611	1755
SLF	0.084	0.047	-0.008	0.176	0.075	0.199	612	1758
SS	0.053	0.047	-0.039	0.145	0.258	0.460	612	1756
UNC	0.143	0.048	0.049	0.237	0.003	0.070	591	1692
AverageMD	0.038	0.047	-0.054	0.129	0.424	0.631	615	1763

Table S12. Differences in regional RD between individuals with a history of suicide attempt and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Attempt	CLC
ACR	0.093	0.047	0.001	0.185	0.048	0.120	613	1761
ALIC	0.046	0.047	-0.046	0.139	0.324	0.470	610	1758
BCC	0.070	0.047	-0.022	0.162	0.137	0.264	610	1763
CC	0.080	0.047	-0.012	0.172	0.088	0.183	612	1764
CGC	0.045	0.047	-0.047	0.137	0.339	0.470	614	1762
CGH	-0.039	0.047	-0.131	0.053	0.408	0.510	613	1763
CR	0.120	0.047	0.028	0.212	0.010	0.079	614	1762
CST	-0.018	0.047	-0.110	0.074	0.706	0.706	611	1762
EC	0.036	0.047	-0.056	0.128	0.443	0.527	614	1757
FX	0.019	0.047	-0.073	0.111	0.680	0.706	612	1755
FXST	0.107	0.047	0.015	0.198	0.023	0.082	616	1760
GCC	0.042	0.047	-0.050	0.134	0.374	0.492	613	1762
IC	0.024	0.047	-0.068	0.116	0.606	0.659	613	1761
IFO	-0.057	0.047	-0.149	0.036	0.229	0.382	612	1728
PCR	0.129	0.047	0.037	0.221	0.006	0.079	617	1759
PLIC	-0.026	0.047	-0.119	0.066	0.575	0.653	610	1757
PTR	0.107	0.047	0.015	0.199	0.023	0.082	614	1761
RLIC	0.058	0.047	-0.034	0.150	0.215	0.382	615	1761
SCC	0.097	0.047	0.005	0.189	0.040	0.120	613	1752
SCR	0.107	0.047	0.015	0.199	0.023	0.082	613	1763
SFO	0.090	0.047	-0.002	0.182	0.057	0.129	612	1761
SLF	0.121	0.047	0.029	0.213	0.010	0.079	614	1762
SS	0.093	0.047	0.001	0.185	0.047	0.120	613	1757
UNC	0.119	0.048	0.026	0.213	0.013	0.079	594	1692
AverageRD	0.047	0.047	-0.045	0.139	0.320	0.470	614	1762

## Recent suicidal ideation

Table S13. Differences in regional AD between individuals with recent suicidal ideation and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	HC
ACR	-0.065	0.048	-0.159	0.029	0.178	0.581	677	1217
ALIC	-0.036	0.048	-0.130	0.058	0.452	0.753	675	1206
BCC	-0.055	0.048	-0.149	0.039	0.256	0.581	675	1211
CC	-0.085	0.048	-0.179	0.009	0.075	0.520	678	1214
CGC	-0.061	0.048	-0.155	0.033	0.206	0.581	677	1214
CGH	0.056	0.048	-0.038	0.150	0.247	0.581	673	1216
CR	-0.046	0.048	-0.140	0.048	0.342	0.713	676	1216
CST	0.009	0.048	-0.085	0.104	0.845	0.845	674	1206
EC	-0.012	0.048	-0.106	0.082	0.806	0.839	669	1217
FX	0.025	0.048	-0.069	0.119	0.606	0.802	676	1214
FXST	0.041	0.048	-0.053	0.135	0.390	0.749	677	1219
GCC	-0.071	0.048	-0.165	0.023	0.138	0.576	677	1215
IC	-0.016	0.048	-0.110	0.078	0.734	0.802	674	1214
IFO	0.104	0.048	0.010	0.198	0.031	0.520	676	1213
PCR	-0.023	0.048	-0.117	0.071	0.631	0.802	675	1212
PLIC	-0.016	0.048	-0.110	0.078	0.737	0.802	673	1216
PTR	0.033	0.048	-0.061	0.127	0.495	0.773	674	1217
RLIC	0.020	0.048	-0.074	0.114	0.681	0.802	674	1220
SCC	-0.094	0.048	-0.188	0.000	0.050	0.520	679	1214
SCR	-0.029	0.048	-0.123	0.065	0.549	0.802	675	1215
SFO	-0.018	0.048	-0.113	0.076	0.702	0.802	673	1211
SLF	-0.078	0.048	-0.172	0.016	0.104	0.520	676	1217
SS	0.056	0.048	-0.038	0.150	0.242	0.581	677	1216
UNC	0.083	0.049	-0.013	0.179	0.089	0.520	661	1152
AverageAD	-0.038	0.048	-0.133	0.056	0.425	0.753	674	1214

Table S14. Differences in regional FA between individuals with recent suicidal ideation and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	HC
ACR	-0.112	0.047	-0.204	-0.020	0.018	0.064	712	1236
ALIC	-0.095	0.047	-0.187	-0.003	0.044	0.104	712	1235
BCC	-0.087	0.047	-0.179	0.006	0.066	0.128	709	1232
CC	-0.108	0.047	-0.201	-0.016	0.022	0.068	710	1233
CGC	-0.019	0.047	-0.111	0.074	0.691	0.742	711	1233
CGH	-0.074	0.047	-0.166	0.018	0.117	0.195	711	1234
CR	-0.133	0.047	-0.225	-0.041	0.005	0.031	710	1233
CST	-0.029	0.047	-0.122	0.063	0.537	0.610	710	1223
EC	-0.093	0.047	-0.186	-0.001	0.048	0.104	713	1230
FX	-0.015	0.047	-0.108	0.077	0.746	0.746	709	1230
FXST	-0.070	0.047	-0.162	0.023	0.141	0.207	711	1234
GCC	-0.148	0.047	-0.241	-0.056	0.002	0.017	711	1233
IC	-0.070	0.047	-0.162	0.023	0.140	0.207	710	1235
IFO	0.017	0.047	-0.075	0.110	0.713	0.742	710	1226
PCR	-0.118	0.047	-0.210	-0.025	0.013	0.063	710	1234
PLIC	-0.036	0.047	-0.129	0.056	0.442	0.526	706	1234
PTR	-0.079	0.047	-0.172	0.013	0.093	0.166	710	1235
RLIC	-0.052	0.047	-0.145	0.040	0.266	0.350	712	1232
SCC	-0.042	0.047	-0.135	0.050	0.371	0.464	711	1230
SCR	-0.112	0.047	-0.204	-0.019	0.018	0.064	709	1234
SFO	-0.145	0.047	-0.238	-0.053	0.002	0.017	709	1232
SLF	-0.097	0.047	-0.190	-0.005	0.039	0.104	712	1233
SS	-0.093	0.047	-0.185	0.000	0.050	0.104	710	1230
UNC	-0.061	0.048	-0.154	0.033	0.207	0.287	696	1169
AverageFA	-0.158	0.048	-0.251	-0.064	0.001	0.017	685	1231



Table S15. Differences in regional MD between individuals with recent suicidal ideation and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	HC
ACR	0.029	0.047	-0.063	0.122	0.536	0.706	703	1217
ALIC	0.053	0.048	-0.040	0.147	0.262	0.409	699	1204
BCC	0.059	0.047	-0.033	0.152	0.211	0.389	702	1213
CC	0.032	0.047	-0.061	0.125	0.499	0.693	703	1214
CGC	-0.008	0.047	-0.101	0.085	0.867	0.867	700	1214
CGH	0.089	0.047	-0.004	0.182	0.061	0.256	702	1219
CR	0.070	0.047	-0.023	0.163	0.143	0.324	701	1216
CST	-0.008	0.048	-0.101	0.085	0.863	0.867	698	1203
EC	0.082	0.048	-0.011	0.176	0.084	0.278	697	1209
FX	-0.017	0.047	-0.110	0.076	0.720	0.805	700	1209
FXST	0.103	0.047	0.010	0.196	0.030	0.189	703	1215
GCC	0.017	0.047	-0.076	0.110	0.722	0.805	704	1216
IC	0.043	0.047	-0.050	0.136	0.370	0.544	701	1212
IFO	0.078	0.047	-0.016	0.171	0.103	0.287	700	1213
PCR	0.081	0.047	-0.012	0.174	0.089	0.278	700	1217
PLIC	-0.016	0.047	-0.109	0.077	0.740	0.805	699	1215
PTR	0.094	0.047	0.001	0.187	0.048	0.241	701	1215
RLIC	0.060	0.047	-0.033	0.153	0.206	0.389	703	1214
SCC	-0.018	0.047	-0.111	0.075	0.703	0.805	701	1212
SCR	0.121	0.047	0.028	0.214	0.011	0.138	703	1214
SFO	0.059	0.048	-0.034	0.152	0.218	0.389	699	1212
SLF	0.054	0.047	-0.039	0.147	0.259	0.409	701	1216
SS	0.071	0.047	-0.022	0.164	0.136	0.324	700	1219
UNC	0.135	0.048	0.040	0.230	0.005	0.132	684	1145
AverageMD	0.109	0.048	0.015	0.203	0.023	0.189	675	1212

Table S16. Differences in regional RD between individuals with recent suicidal ideation and healthy controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	HC
ACR	0.086	0.048	-0.008	0.180	0.073	0.142	674	1214
ALIC	0.080	0.048	-0.014	0.174	0.097	0.158	674	1206
BCC	0.079	0.048	-0.015	0.173	0.101	0.158	673	1211
CC	0.068	0.048	-0.026	0.162	0.158	0.227	674	1214
CGC	-0.006	0.048	-0.100	0.088	0.900	0.938	673	1213
CGH	0.088	0.048	-0.006	0.182	0.066	0.142	675	1221
CR	0.114	0.048	0.020	0.208	0.018	0.112	674	1215
CST	0.035	0.048	-0.059	0.130	0.464	0.504	673	1202
EC	0.099	0.048	0.005	0.193	0.040	0.125	674	1213
FX	0.002	0.048	-0.092	0.097	0.963	0.963	674	1206
FXST	0.092	0.048	-0.002	0.186	0.056	0.142	676	1215
GCC	0.067	0.048	-0.027	0.161	0.163	0.227	677	1215
IC	0.084	0.048	-0.010	0.179	0.080	0.142	674	1213
IFO	0.045	0.048	-0.049	0.139	0.352	0.400	674	1211
PCR	0.101	0.048	0.007	0.195	0.036	0.125	675	1213
PLIC	0.046	0.048	-0.048	0.140	0.340	0.400	671	1215
PTR	0.105	0.048	0.010	0.199	0.030	0.125	676	1214
RLIC	0.057	0.048	-0.037	0.151	0.238	0.313	676	1215
SCC	0.050	0.048	-0.044	0.145	0.296	0.370	675	1207
SCR	0.150	0.048	0.056	0.244	0.002	0.040	677	1215
SFO	0.088	0.048	-0.006	0.182	0.068	0.142	673	1209
SLF	0.100	0.048	0.006	0.194	0.038	0.125	678	1214
SS	0.085	0.048	-0.009	0.179	0.078	0.142	676	1219
UNC	0.122	0.049	0.026	0.218	0.013	0.107	659	1143
AverageRD	0.142	0.048	0.048	0.236	0.003	0.040	677	1214

Table S17. Differences in regional AD between individuals with recent suicidal ideation and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	CLC
ACR	0.069	0.049	-0.026	0.164	0.156	0.862	677	1136
ALIC	0.007	0.049	-0.088	0.102	0.885	0.976	675	1139
BCC	-0.016	0.049	-0.111	0.079	0.746	0.938	675	1138
CC	-0.001	0.049	-0.097	0.094	0.976	0.976	678	1137
CGC	0.126	0.049	0.031	0.221	0.010	0.135	677	1137
CGH	0.006	0.049	-0.089	0.102	0.898	0.976	673	1138
CR	0.040	0.049	-0.055	0.135	0.414	0.938	676	1136
CST	-0.003	0.049	-0.098	0.093	0.959	0.976	674	1136
EC	0.063	0.049	-0.033	0.158	0.199	0.862	669	1134
FX	-0.025	0.049	-0.121	0.070	0.601	0.938	676	1130
FXST	0.031	0.049	-0.064	0.126	0.529	0.938	677	1140
GCC	0.039	0.049	-0.057	0.134	0.427	0.938	677	1136
IC	0.018	0.049	-0.077	0.113	0.709	0.938	674	1139
IFO	0.088	0.049	-0.008	0.183	0.072	0.599	676	1139
PCR	0.019	0.049	-0.076	0.114	0.699	0.938	675	1134
PLIC	0.015	0.049	-0.081	0.110	0.762	0.938	673	1139
PTR	0.054	0.049	-0.041	0.150	0.265	0.938	674	1137
RLIC	0.061	0.049	-0.034	0.157	0.207	0.862	674	1137
SCC	-0.017	0.049	-0.112	0.078	0.723	0.938	679	1137
SCR	0.013	0.049	-0.082	0.108	0.788	0.938	675	1136
SFO	-0.020	0.049	-0.115	0.075	0.683	0.938	673	1132
SLF	0.040	0.049	-0.055	0.135	0.411	0.938	676	1134
SS	0.019	0.049	-0.076	0.115	0.689	0.938	677	1134
UNC	0.027	0.049	-0.069	0.124	0.579	0.938	661	1088
AverageAD	0.124	0.049	0.029	0.220	0.011	0.135	674	1134

Table S18. Differences in regional FA between individuals with recent suicidal ideation and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	CLC
ACR	-0.048	0.047	-0.141	0.045	0.312	0.576	712	1182
ALIC	-0.026	0.047	-0.119	0.067	0.582	0.766	712	1177
BCC	0.013	0.048	-0.080	0.106	0.786	0.854	709	1174
CC	0.009	0.048	-0.084	0.102	0.847	0.883	710	1175
CGC	0.038	0.048	-0.055	0.131	0.422	0.660	711	1177
CGH	-0.065	0.048	-0.159	0.028	0.169	0.576	711	1176
CR	-0.065	0.048	-0.158	0.028	0.173	0.576	710	1179
CST	-0.005	0.048	-0.099	0.088	0.910	0.910	710	1176
EC	-0.034	0.047	-0.127	0.059	0.477	0.663	713	1175
FX	0.048	0.048	-0.045	0.141	0.316	0.576	709	1177
FXST	-0.064	0.048	-0.157	0.029	0.181	0.576	711	1176
GCC	-0.014	0.047	-0.107	0.080	0.775	0.854	711	1177
IC	-0.053	0.048	-0.146	0.040	0.268	0.576	710	1178
IFO	0.059	0.047	-0.034	0.152	0.215	0.576	710	1182
PCR	-0.058	0.048	-0.151	0.035	0.223	0.576	710	1179
PLIC	-0.015	0.048	-0.109	0.078	0.749	0.854	706	1176
PTR	-0.065	0.048	-0.158	0.028	0.172	0.576	710	1179
RLIC	-0.077	0.047	-0.170	0.016	0.104	0.576	712	1178
SCC	0.047	0.048	-0.047	0.140	0.328	0.576	711	1176
SCR	-0.054	0.048	-0.147	0.039	0.255	0.576	709	1180
SFO	-0.101	0.048	-0.195	-0.008	0.033	0.576	709	1180
SLF	-0.023	0.047	-0.116	0.070	0.632	0.790	712	1183
SS	-0.087	0.048	-0.180	0.006	0.067	0.576	710	1180
UNC	-0.035	0.048	-0.130	0.059	0.467	0.663	696	1131
AverageFA	-0.046	0.048	-0.140	0.049	0.345	0.576	685	1140

Table S19. Differences in regional MD between individuals with recent suicidal ideation and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	CLC
ACR	0.056	0.048	-0.037	0.150	0.240	0.659	703	1175
ALIC	0.047	0.048	-0.046	0.141	0.324	0.659	699	1173
BCC	-0.012	0.048	-0.105	0.082	0.805	0.839	702	1174
CC	-0.022	0.048	-0.116	0.071	0.643	0.765	703	1178
CGC	0.092	0.048	-0.002	0.186	0.054	0.659	700	1177
CGH	0.028	0.048	-0.066	0.121	0.560	0.719	702	1176
CR	0.057	0.048	-0.036	0.151	0.231	0.659	701	1176
CST	-0.015	0.048	-0.109	0.079	0.752	0.818	698	1170
EC	0.062	0.048	-0.032	0.156	0.197	0.659	697	1174
FX	-0.036	0.048	-0.129	0.058	0.459	0.719	700	1162
FXST	0.053	0.048	-0.040	0.147	0.264	0.659	703	1174
GCC	0.017	0.048	-0.077	0.110	0.725	0.818	704	1177
IC	0.048	0.048	-0.046	0.142	0.315	0.659	701	1173
IFO	0.006	0.048	-0.087	0.100	0.896	0.896	700	1175
PCR	0.033	0.048	-0.060	0.127	0.485	0.719	700	1175
PLIC	0.030	0.048	-0.064	0.124	0.530	0.719	699	1175
PTR	0.045	0.048	-0.048	0.139	0.343	0.659	701	1176
RLIC	0.069	0.048	-0.025	0.162	0.149	0.659	703	1177
SCC	-0.050	0.048	-0.143	0.044	0.300	0.659	701	1174
SCR	0.076	0.048	-0.018	0.169	0.113	0.659	703	1173
SFO	0.034	0.048	-0.059	0.128	0.473	0.719	699	1171
SLF	0.059	0.048	-0.035	0.152	0.221	0.659	701	1174
SS	0.032	0.048	-0.062	0.125	0.509	0.719	700	1171
UNC	0.027	0.048	-0.068	0.122	0.575	0.719	684	1127
AverageMD	0.090	0.049	-0.006	0.185	0.066	0.659	675	1136

Table S20. Differences in regional RD between individuals with recent suicidal ideation and clinical controls. Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	Ideation	CLC
ACR	0.080	0.049	-0.015	0.176	0.099	0.248	674	1135
ALIC	0.064	0.049	-0.031	0.160	0.187	0.334	674	1133
BCC	-0.012	0.049	-0.108	0.083	0.801	0.870	673	1137
CC	-0.006	0.049	-0.101	0.089	0.901	0.939	674	1135
CGC	0.034	0.049	-0.061	0.130	0.479	0.631	673	1136
CGH	0.047	0.049	-0.048	0.143	0.330	0.485	675	1135
CR	0.081	0.049	-0.015	0.176	0.098	0.248	674	1135
CST	-0.003	0.049	-0.099	0.092	0.946	0.946	673	1137
EC	0.074	0.049	-0.021	0.170	0.128	0.267	674	1132
FX	-0.038	0.049	-0.133	0.058	0.438	0.608	674	1122
FXST	0.089	0.049	-0.006	0.184	0.067	0.248	676	1136
GCC	0.025	0.049	-0.070	0.120	0.604	0.755	677	1134
IC	0.096	0.049	0.001	0.191	0.049	0.248	674	1136
IFO	-0.018	0.049	-0.113	0.077	0.710	0.807	674	1133
PCR	0.061	0.049	-0.034	0.156	0.210	0.338	675	1136
PLIC	0.077	0.049	-0.019	0.172	0.115	0.261	671	1135
PTR	0.066	0.049	-0.030	0.161	0.178	0.334	676	1136
RLIC	0.085	0.049	-0.010	0.180	0.081	0.248	676	1137
SCC	-0.022	0.049	-0.118	0.073	0.647	0.771	675	1133
SCR	0.095	0.049	0.000	0.190	0.051	0.248	677	1134
SFO	0.082	0.049	-0.013	0.178	0.092	0.248	673	1135
SLF	0.081	0.049	-0.014	0.176	0.095	0.248	678	1136
SS	0.099	0.049	0.004	0.194	0.042	0.248	676	1132
UNC	0.061	0.049	-0.036	0.158	0.216	0.338	659	1087
AverageRD	0.083	0.049	-0.012	0.178	0.088	0.248	677	1136

## CSSRS analyses

Table S21. Association between lifetime intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional AD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.070	-0.186	0.047	-1.174	0.241	0.367	299
ALIC	-0.080	-0.197	0.037	-1.338	0.182	0.325	299
BCC	-0.122	-0.240	-0.005	-2.050	0.041	0.206	299
CC	-0.109	-0.227	0.009	-1.825	0.069	0.288	299
CGC	0.076	-0.040	0.193	1.287	0.199	0.332	299
CGH	-0.063	-0.177	0.052	-1.076	0.283	0.393	299
CR	-0.080	-0.197	0.038	-1.338	0.182	0.325	299
CST	-0.102	-0.219	0.015	-1.723	0.086	0.298	297
EC	-0.041	-0.159	0.077	-0.690	0.491	0.548	297
FX	-0.117	-0.228	-0.006	-2.081	0.038	0.206	298
FXST	-0.058	-0.170	0.055	-1.002	0.317	0.396	299
GCC	-0.093	-0.211	0.024	-1.563	0.119	0.298	297
IC	-0.118	-0.229	-0.006	-2.077	0.039	0.206	297
IFO	-0.093	-0.209	0.024	-1.565	0.119	0.298	298
PCR	-0.040	-0.158	0.078	-0.666	0.506	0.548	299
PLIC	-0.126	-0.235	-0.017	-2.285	0.023	0.206	298
PTR	-0.052	-0.170	0.066	-0.860	0.390	0.465	299
RLIC	-0.092	-0.205	0.020	-1.612	0.108	0.298	299
SCC	-0.062	-0.179	0.056	-1.031	0.303	0.396	298
SCR	-0.082	-0.199	0.036	-1.371	0.171	0.325	299
SFO	-0.089	-0.207	0.029	-1.489	0.137	0.312	298
SLF	-0.069	-0.187	0.049	-1.154	0.250	0.367	299
SS	-0.036	-0.154	0.081	-0.608	0.544	0.548	298
UNC	-0.035	-0.151	0.081	-0.601	0.548	0.548	299
AverageAD	-0.127	-0.245	-0.010	-2.137	0.033	0.206	298

Table S22. Association between lifetime intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional FA (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.047	-0.151	0.057	-0.894	0.372	0.990	298
ALIC	-0.028	-0.144	0.088	-0.474	0.636	0.990	298
BCC	-0.010	-0.126	0.107	-0.160	0.873	0.990	298
CC	-0.038	-0.153	0.076	-0.662	0.508	0.990	298
CGC	0.002	-0.113	0.117	0.040	0.968	0.990	299
CGH	-0.004	-0.122	0.113	-0.071	0.943	0.990	299
CR	-0.005	-0.111	0.100	-0.102	0.919	0.990	298
CST	0.008	-0.110	0.126	0.137	0.891	0.990	298
EC	-0.028	-0.138	0.082	-0.496	0.620	0.990	298
FX	0.056	-0.052	0.163	1.022	0.308	0.990	296
FXST	-0.079	-0.191	0.032	-1.402	0.162	0.990	297
GCC	-0.052	-0.162	0.058	-0.931	0.353	0.990	298
IC	-0.027	-0.141	0.087	-0.470	0.638	0.990	298
IFO	-0.104	-0.219	0.011	-1.779	0.076	0.953	298
PCR	0.014	-0.098	0.126	0.249	0.803	0.990	299
PLIC	0.013	-0.106	0.131	0.209	0.835	0.990	296
PTR	-0.053	-0.158	0.052	-0.995	0.321	0.990	297
RLIC	-0.028	-0.138	0.081	-0.514	0.608	0.990	298
SCC	-0.056	-0.173	0.061	-0.946	0.345	0.990	295
SCR	0.032	-0.080	0.143	0.561	0.575	0.990	299
SFO	-0.135	-0.251	-0.020	-2.311	0.022	0.538	297
SLF	-0.009	-0.121	0.103	-0.160	0.873	0.990	299
SS	0.027	-0.081	0.135	0.490	0.625	0.990	297
UNC	-0.001	-0.115	0.114	-0.012	0.990	0.990	299
AverageFA	-0.014	-0.125	0.096	-0.258	0.797	0.990	298



Table S23. Association between lifetime intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional MD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.013	-0.126	0.101	-0.219	0.827	0.862	299
ALIC	-0.005	-0.123	0.113	-0.085	0.933	0.933	298
BCC	-0.073	-0.190	0.044	-1.234	0.218	0.683	298
CC	-0.058	-0.175	0.059	-0.982	0.327	0.683	299
CGC	0.023	-0.094	0.141	0.391	0.696	0.857	299
CGH	-0.064	-0.180	0.051	-1.096	0.274	0.683	299
CR	-0.040	-0.155	0.076	-0.678	0.498	0.779	299
CST	-0.085	-0.201	0.032	-1.430	0.154	0.683	297
EC	-0.021	-0.135	0.092	-0.367	0.714	0.857	297
FX	-0.139	-0.244	-0.035	-2.620	0.009	0.232	294
FXST	0.015	-0.104	0.133	0.246	0.806	0.862	298
GCC	-0.029	-0.144	0.087	-0.488	0.626	0.857	299
IC	-0.067	-0.185	0.050	-1.126	0.261	0.683	297
IFO	0.014	-0.103	0.131	0.231	0.818	0.862	298
PCR	-0.040	-0.158	0.077	-0.678	0.499	0.779	299
PLIC	-0.089	-0.201	0.024	-1.555	0.121	0.683	297
PTR	-0.058	-0.175	0.059	-0.980	0.328	0.683	296
RLIC	-0.080	-0.199	0.038	-1.337	0.182	0.683	298
SCC	-0.022	-0.140	0.097	-0.361	0.719	0.857	298
SCR	-0.070	-0.187	0.047	-1.177	0.240	0.683	299
SFO	0.022	-0.096	0.139	0.359	0.720	0.857	298
SLF	-0.079	-0.196	0.038	-1.337	0.182	0.683	298
SS	-0.041	-0.157	0.076	-0.686	0.493	0.779	298
UNC	-0.053	-0.169	0.064	-0.893	0.373	0.717	298
AverageMD	-0.073	-0.190	0.045	-1.217	0.225	0.683	297

Table S24. Association between lifetime intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional RD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	0.008	-0.100	0.116	0.143	0.887	0.964	298
ALIC	0.000	-0.117	0.116	-0.006	0.995	0.995	297
BCC	-0.033	-0.151	0.084	-0.554	0.580	0.926	298
CC	-0.002	-0.119	0.115	-0.038	0.970	0.995	298
CGC	-0.039	-0.155	0.076	-0.673	0.502	0.926	298
CGH	-0.042	-0.159	0.074	-0.717	0.474	0.926	299
CR	-0.016	-0.126	0.095	-0.282	0.778	0.926	298
CST	-0.071	-0.188	0.046	-1.190	0.235	0.926	297
EC	-0.024	-0.135	0.087	-0.431	0.667	0.926	297
FX	-0.152	-0.256	-0.048	-2.877	0.004	0.108	293
FXST	0.042	-0.075	0.158	0.704	0.482	0.926	299
GCC	0.024	-0.091	0.139	0.413	0.680	0.926	298
IC	-0.017	-0.134	0.100	-0.284	0.777	0.926	298
IFO	0.067	-0.049	0.182	1.136	0.257	0.926	298
PCR	-0.023	-0.138	0.093	-0.385	0.701	0.926	299
PLIC	-0.055	-0.173	0.062	-0.923	0.357	0.926	296
PTR	-0.024	-0.138	0.089	-0.426	0.671	0.926	296
RLIC	-0.018	-0.134	0.097	-0.315	0.753	0.926	298
SCC	0.014	-0.105	0.133	0.230	0.819	0.930	297
SCR	-0.040	-0.153	0.073	-0.701	0.484	0.926	298
SFO	0.061	-0.056	0.178	1.027	0.305	0.926	299
SLF	-0.034	-0.148	0.080	-0.589	0.556	0.926	299
SS	-0.046	-0.158	0.067	-0.801	0.424	0.926	297
UNC	-0.046	-0.161	0.069	-0.792	0.429	0.926	299
AverageRD	-0.052	-0.167	0.063	-0.894	0.372	0.926	298

Table S25. Association between recent intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional AD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.077	-0.188	0.034	-1.359	0.175	0.273	338
ALIC	-0.078	-0.189	0.033	-1.376	0.170	0.273	338
BCC	-0.101	-0.213	0.011	-1.780	0.076	0.190	337
CC	-0.086	-0.198	0.026	-1.507	0.133	0.255	337
CGC	-0.067	-0.179	0.045	-1.182	0.238	0.313	338
CGH	0.014	-0.097	0.124	0.241	0.810	0.843	338
CR	-0.119	-0.231	-0.008	-2.110	0.036	0.149	338
CST	-0.036	-0.149	0.076	-0.637	0.525	0.596	336
EC	-0.089	-0.201	0.023	-1.561	0.120	0.255	335
FX	-0.052	-0.161	0.056	-0.946	0.345	0.410	337
FXST	-0.082	-0.192	0.029	-1.454	0.147	0.263	337
GCC	-0.022	-0.134	0.091	-0.379	0.705	0.766	336
IC	-0.147	-0.256	-0.039	-2.665	0.008	0.083	336
IFO	-0.105	-0.216	0.006	-1.865	0.063	0.175	338
PCR	-0.087	-0.199	0.025	-1.531	0.127	0.255	338
PLIC	-0.144	-0.249	-0.038	-2.683	0.008	0.083	337
PTR	-0.068	-0.181	0.045	-1.185	0.237	0.313	337
RLIC	-0.145	-0.254	-0.035	-2.585	0.010	0.083	337
SCC	-0.074	-0.186	0.038	-1.304	0.193	0.284	336
SCR	-0.141	-0.252	-0.030	-2.488	0.013	0.083	338
SFO	-0.117	-0.229	-0.005	-2.056	0.041	0.149	337
SLF	-0.112	-0.224	-0.001	-1.981	0.048	0.151	338
SS	-0.057	-0.169	0.056	-0.986	0.325	0.406	336
UNC	0.008	-0.104	0.120	0.139	0.890	0.890	338
AverageAD	-0.116	-0.227	-0.004	-2.043	0.042	0.149	338

Table S26. Association between recent intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional FA (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	0.026	-0.078	0.130	0.490	0.625	0.855	336
ALIC	0.047	-0.063	0.157	0.842	0.401	0.855	335
BCC	-0.021	-0.133	0.091	-0.372	0.710	0.855	334
CC	-0.040	-0.149	0.070	-0.711	0.477	0.855	335
CGC	-0.017	-0.127	0.093	-0.306	0.759	0.855	338
CGH	0.013	-0.099	0.126	0.233	0.816	0.855	338
CR	0.019	-0.087	0.124	0.347	0.729	0.855	336
CST	0.052	-0.060	0.164	0.914	0.361	0.855	337
EC	-0.052	-0.160	0.057	-0.936	0.350	0.855	337
FX	-0.080	-0.189	0.029	-1.450	0.148	0.855	335
FXST	-0.013	-0.124	0.098	-0.233	0.816	0.855	334
GCC	-0.040	-0.144	0.065	-0.749	0.455	0.855	335
IC	-0.013	-0.122	0.097	-0.227	0.821	0.855	336
IFO	-0.163	-0.272	-0.054	-2.933	0.004	0.090	337
PCR	0.022	-0.088	0.132	0.399	0.690	0.855	338
PLIC	0.014	-0.098	0.126	0.242	0.809	0.855	336
PTR	-0.020	-0.127	0.087	-0.366	0.715	0.855	336
RLIC	-0.048	-0.156	0.061	-0.867	0.386	0.855	336
SCC	-0.047	-0.159	0.065	-0.825	0.410	0.855	333
SCR	-0.008	-0.117	0.101	-0.143	0.886	0.886	338
SFO	-0.050	-0.162	0.062	-0.878	0.381	0.855	336
SLF	0.036	-0.073	0.145	0.645	0.519	0.855	338
SS	0.038	-0.070	0.146	0.691	0.490	0.855	335
UNC	0.056	-0.052	0.165	1.018	0.309	0.855	338
AverageFA	0.035	-0.073	0.144	0.640	0.522	0.855	336

Table S27. Association between recent intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional MD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.068	-0.177	0.040	-1.242	0.215	0.358	337
ALIC	-0.088	-0.200	0.023	-1.562	0.119	0.271	337
BCC	-0.055	-0.166	0.057	-0.959	0.338	0.448	336
CC	-0.060	-0.171	0.051	-1.058	0.291	0.428	338
CGC	-0.097	-0.208	0.014	-1.715	0.087	0.260	337
CGH	-0.003	-0.114	0.109	-0.052	0.959	0.959	338
CR	-0.104	-0.214	0.005	-1.872	0.062	0.222	337
CST	-0.052	-0.164	0.060	-0.916	0.360	0.451	334
EC	-0.053	-0.162	0.056	-0.954	0.341	0.448	337
FX	-0.020	-0.126	0.086	-0.371	0.711	0.808	334
FXST	-0.094	-0.207	0.018	-1.650	0.100	0.260	337
GCC	-0.009	-0.117	0.100	-0.156	0.876	0.913	338
IC	-0.164	-0.275	-0.053	-2.897	0.004	0.087	336
IFO	0.017	-0.095	0.128	0.294	0.769	0.836	337
PCR	-0.093	-0.205	0.019	-1.630	0.104	0.260	337
PLIC	-0.149	-0.257	-0.041	-2.714	0.007	0.087	336
PTR	-0.074	-0.186	0.038	-1.307	0.192	0.343	334
RLIC	-0.140	-0.251	-0.029	-2.482	0.014	0.101	336
SCC	-0.063	-0.175	0.050	-1.093	0.275	0.428	336
SCR	-0.135	-0.246	-0.024	-2.395	0.017	0.101	337
SFO	-0.076	-0.188	0.037	-1.323	0.187	0.343	337
SLF	-0.132	-0.243	-0.021	-2.333	0.020	0.101	336
SS	-0.080	-0.191	0.031	-1.415	0.158	0.329	336
UNC	-0.035	-0.147	0.077	-0.612	0.541	0.644	338
AverageMD	-0.111	-0.222	0.000	-1.967	0.050	0.208	337

Table S28. Association between recent intensity of ideation (scored 0-5) in the Columbia Suicide Severity Rating Scale and regional RD (healthy controls were excluded). Beta: standardized beta, p: p-value, FDR-p: FDR corrected p-value.

Region	Beta	Lower CI	Upper CI	T-value	P	FDR-p	N
ACR	-0.060	-0.166	0.045	-1.125	0.261	0.670	334
ALIC	-0.068	-0.179	0.043	-1.201	0.231	0.670	336
BCC	-0.009	-0.121	0.103	-0.157	0.875	0.916	336
CC	0.009	-0.103	0.120	0.152	0.879	0.916	335
CGC	-0.049	-0.159	0.060	-0.885	0.377	0.685	336
CGH	-0.006	-0.118	0.107	-0.100	0.920	0.920	338
CR	-0.062	-0.169	0.045	-1.143	0.254	0.670	334
CST	-0.014	-0.126	0.098	-0.248	0.804	0.916	334
EC	-0.018	-0.125	0.090	-0.327	0.744	0.916	336
FX	-0.009	-0.113	0.096	-0.160	0.873	0.916	334
FXST	-0.080	-0.192	0.031	-1.417	0.157	0.670	338
GCC	0.030	-0.077	0.136	0.548	0.584	0.913	335
IC	-0.058	-0.169	0.053	-1.033	0.302	0.670	336
IFO	0.115	0.005	0.225	2.051	0.041	0.670	337
PCR	-0.060	-0.172	0.051	-1.062	0.289	0.670	336
PLIC	-0.056	-0.168	0.055	-0.993	0.322	0.670	335
PTR	-0.015	-0.126	0.095	-0.268	0.789	0.916	333
RLIC	-0.042	-0.153	0.069	-0.740	0.460	0.766	336
SCC	0.022	-0.091	0.134	0.376	0.707	0.916	335
SCR	-0.073	-0.182	0.036	-1.314	0.190	0.670	336
SFO	-0.025	-0.138	0.088	-0.435	0.664	0.916	338
SLF	-0.092	-0.202	0.017	-1.659	0.098	0.670	337
SS	-0.071	-0.180	0.039	-1.271	0.205	0.670	334
UNC	-0.049	-0.159	0.061	-0.873	0.383	0.685	338
AverageRD	-0.080	-0.189	0.030	-1.435	0.152	0.670	337

Table S29. Differences in regional AD between people with a lifetime history of an actual suicide attempt and those without a history of any suicide attempt determined using the Columbia Suicide Severity Rating Scale (healthy controls were excluded).

D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval, HC: healthy controls, CC: clinical controls.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N actual attempt	N no attempt
ACR	-0.100	0.108	0.357	0.733	-0.311	0.111	134	242
ALIC	-0.102	0.108	0.347	0.733	-0.314	0.109	134	241
BCC	-0.244	0.108	0.026	0.177	-0.456	-0.032	134	240
CC	-0.256	0.108	0.019	0.177	-0.468	-0.044	134	239
CGC	0.025	0.108	0.821	0.933	-0.186	0.235	134	243
CGH	0.056	0.108	0.605	0.756	-0.155	0.267	134	244
CR	-0.081	0.108	0.459	0.733	-0.292	0.131	134	240
CST	0.007	0.108	0.951	0.988	-0.205	0.218	133	242
EC	-0.117	0.108	0.285	0.713	-0.329	0.095	133	240
FX	0.032	0.108	0.769	0.915	-0.179	0.243	134	244
FXST	-0.068	0.108	0.529	0.735	-0.280	0.143	134	242
GCC	-0.236	0.108	0.031	0.177	-0.448	-0.024	133	244
IC	-0.137	0.108	0.211	0.658	-0.349	0.075	133	239
IFO	-0.085	0.108	0.433	0.733	-0.296	0.126	134	242
PCR	-0.008	0.108	0.942	0.988	-0.219	0.203	134	242
PLIC	-0.078	0.108	0.474	0.733	-0.290	0.133	134	240
PTR	-0.161	0.108	0.141	0.504	-0.373	0.051	134	239
RLIC	-0.230	0.108	0.035	0.177	-0.443	-0.018	134	238
SCC	-0.120	0.108	0.271	0.713	-0.332	0.091	134	239
SCR	-0.074	0.108	0.499	0.733	-0.286	0.138	134	238
SFO	-0.002	0.108	0.988	0.988	-0.213	0.209	134	242
SLF	-0.260	0.108	0.017	0.177	-0.473	-0.048	134	240
SS	-0.060	0.108	0.582	0.756	-0.271	0.151	134	242
UNC	0.078	0.108	0.472	0.733	-0.133	0.289	134	242
AverageAD	-0.210	0.108	0.055	0.228	-0.422	0.002	134	239

Table S30. Differences in regional FA between people with a lifetime history of an actual suicide attempt and those without a history of any suicide attempt determined using the Columbia Suicide Severity Rating Scale (healthy controls were excluded).

D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval, HC: healthy controls, CC: clinical controls.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N actual attempt	N no attempt
ACR	0.045	0.108	0.678	0.983	-0.167	0.257	132	243
ALIC	-0.022	0.108	0.838	0.983	-0.234	0.189	133	241
BCC	-0.002	0.108	0.983	0.983	-0.215	0.210	131	244
CC	-0.034	0.108	0.757	0.983	-0.246	0.178	132	242
CGC	-0.129	0.108	0.235	0.983	-0.340	0.082	134	244
CGH	-0.065	0.108	0.549	0.983	-0.276	0.146	134	242
CR	0.045	0.108	0.679	0.983	-0.167	0.257	132	242
CST	0.078	0.108	0.471	0.983	-0.133	0.290	134	242
EC	0.005	0.108	0.964	0.983	-0.207	0.216	133	243
FX	-0.009	0.108	0.933	0.983	-0.221	0.203	132	242
FXST	-0.132	0.108	0.227	0.983	-0.345	0.080	132	241
GCC	0.039	0.108	0.719	0.983	-0.173	0.251	132	243
IC	-0.006	0.108	0.957	0.983	-0.218	0.206	133	241
IFO	0.012	0.108	0.911	0.983	-0.199	0.223	134	243
PCR	-0.049	0.108	0.653	0.983	-0.260	0.162	134	244
PLIC	0.103	0.108	0.344	0.983	-0.108	0.315	133	242
PTR	-0.127	0.108	0.246	0.983	-0.339	0.085	132	242
RLIC	-0.062	0.108	0.571	0.983	-0.274	0.150	133	241
SCC	-0.093	0.109	0.397	0.983	-0.307	0.121	130	237
SCR	0.065	0.108	0.547	0.983	-0.145	0.276	134	244
SFO	-0.113	0.108	0.301	0.983	-0.325	0.099	133	242
SLF	-0.122	0.108	0.262	0.983	-0.333	0.089	134	244
SS	0.051	0.108	0.643	0.983	-0.162	0.263	132	239
UNC	-0.096	0.108	0.376	0.983	-0.307	0.115	134	243
AverageFA	-0.022	0.108	0.837	0.983	-0.235	0.190	132	241



Table S31. Differences in regional MD between people with a lifetime history of an actual suicide attempt and those without a history of any suicide attempt determined using the Columbia Suicide Severity Rating Scale (healthy controls were excluded).

D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval, HC: healthy controls, CC: clinical controls.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N actual attempt	N no attempt
ACR	-0.033	0.108	0.760	0.966	-0.244	0.178	134	242
ALIC	0.015	0.108	0.888	0.966	-0.196	0.227	133	244
BCC	-0.063	0.108	0.563	0.966	-0.275	0.149	132	244
CC	-0.092	0.108	0.399	0.966	-0.302	0.119	134	244
CGC	-0.015	0.108	0.889	0.966	-0.226	0.196	134	243
CGH	0.024	0.108	0.827	0.966	-0.187	0.235	134	243
CR	0.006	0.108	0.957	0.997	-0.205	0.217	134	243
CST	-0.040	0.108	0.712	0.966	-0.252	0.172	134	237
EC	-0.090	0.108	0.409	0.966	-0.302	0.121	133	243
FX	-0.027	0.108	0.806	0.966	-0.239	0.185	132	243
FXST	0.051	0.108	0.643	0.966	-0.161	0.262	133	244
GCC	-0.203	0.108	0.063	0.966	-0.414	0.009	134	244
IC	-0.070	0.108	0.522	0.966	-0.281	0.142	133	243
IFO	-0.109	0.108	0.316	0.966	-0.321	0.102	133	244
PCR	0.038	0.108	0.725	0.966	-0.173	0.249	134	243
PLIC	-0.065	0.108	0.548	0.966	-0.277	0.146	133	243
PTR	-0.085	0.108	0.437	0.966	-0.296	0.127	133	242
RLIC	-0.150	0.108	0.171	0.966	-0.361	0.062	133	242
SCC	-0.020	0.108	0.857	0.966	-0.231	0.192	133	242
SCR	0.000	0.108	0.997	0.997	-0.212	0.211	134	242
SFO	0.090	0.108	0.408	0.966	-0.121	0.301	134	243
SLF	-0.077	0.108	0.480	0.966	-0.288	0.134	134	242
SS	-0.141	0.108	0.195	0.966	-0.353	0.070	133	243
UNC	0.126	0.108	0.250	0.966	-0.086	0.338	132	244
AverageMD	-0.086	0.108	0.432	0.966	-0.297	0.126	133	243

Table S32. Differences in regional RD between people with a lifetime history of an actual suicide attempt and those without a history of any suicide attempt determined using the Columbia Suicide Severity Rating Scale (healthy controls were excluded).

D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval, HC: healthy controls, CC: clinical controls.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N actual attempt	N no attempt
ACR	-0.066	0.108	0.548	0.833	-0.278	0.147	132	241
ALIC	-0.084	0.108	0.443	0.833	-0.296	0.128	132	244
BCC	-0.028	0.108	0.794	0.870	-0.240	0.183	132	244
CC	-0.047	0.108	0.665	0.833	-0.259	0.165	132	244
CGC	-0.078	0.108	0.474	0.833	-0.289	0.133	133	244
CGH	-0.001	0.108	0.991	0.991	-0.212	0.210	134	243
CR	-0.011	0.108	0.919	0.957	-0.223	0.201	133	241
CST	-0.058	0.108	0.593	0.833	-0.270	0.153	133	240
EC	-0.127	0.108	0.244	0.833	-0.339	0.084	133	243
FX	-0.063	0.108	0.563	0.833	-0.276	0.149	131	243
FXST	0.059	0.108	0.588	0.833	-0.152	0.270	134	244
GCC	-0.104	0.108	0.340	0.833	-0.316	0.108	132	243
IC	-0.102	0.108	0.348	0.833	-0.314	0.109	133	243
IFO	-0.109	0.108	0.316	0.833	-0.320	0.102	134	244
PCR	0.047	0.108	0.667	0.833	-0.164	0.258	134	242
PLIC	-0.125	0.108	0.252	0.833	-0.337	0.087	133	241
PTR	-0.049	0.108	0.651	0.833	-0.262	0.163	132	241
RLIC	-0.071	0.108	0.513	0.833	-0.283	0.140	133	242
SCC	0.028	0.108	0.798	0.870	-0.184	0.240	132	242
SCR	-0.058	0.108	0.597	0.833	-0.269	0.154	133	243
SFO	0.089	0.108	0.412	0.833	-0.122	0.300	134	244
SLF	0.028	0.108	0.800	0.870	-0.183	0.238	134	243
SS	-0.186	0.108	0.090	0.833	-0.398	0.027	132	242
UNC	0.074	0.108	0.494	0.833	-0.136	0.285	134	244
AverageRD	-0.099	0.108	0.362	0.833	-0.311	0.112	133	244

Table S33. Differences in regional AD between people with a lifetime history of an actual suicide and those without a history of an actual suicide attempt but with lifetime suicidal ideation determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N ideation	N attempt
ACR	-0.254	0.126	0.047	0.343	-0.500	-0.007	122	134
ALIC	-0.098	0.125	0.439	0.610	-0.344	0.147	122	134
BCC	-0.171	0.125	0.179	0.448	-0.417	0.075	122	134
CC	-0.178	0.125	0.161	0.448	-0.424	0.067	122	134
CGC	-0.100	0.125	0.430	0.610	-0.346	0.145	122	134
CGH	0.034	0.125	0.792	0.825	-0.212	0.279	122	134
CR	-0.175	0.125	0.168	0.448	-0.421	0.070	122	134
CST	0.120	0.126	0.345	0.576	-0.126	0.367	121	133
EC	-0.126	0.126	0.324	0.576	-0.372	0.121	121	133
FX	0.175	0.126	0.169	0.448	-0.071	0.422	121	134
FXST	-0.173	0.125	0.175	0.448	-0.418	0.073	122	134
GCC	-0.273	0.126	0.033	0.343	-0.521	-0.026	121	133
IC	-0.037	0.126	0.773	0.825	-0.283	0.209	121	133
IFO	-0.123	0.126	0.334	0.576	-0.369	0.123	121	134
PCR	-0.152	0.125	0.232	0.511	-0.398	0.094	122	134
PLIC	-0.005	0.125	0.971	0.971	-0.250	0.241	121	134
PTR	-0.232	0.126	0.069	0.343	-0.478	0.014	122	134
RLIC	-0.106	0.125	0.404	0.610	-0.352	0.139	122	134
SCC	-0.041	0.125	0.747	0.825	-0.287	0.205	121	134
SCR	-0.045	0.125	0.725	0.825	-0.290	0.201	122	134
SFO	-0.053	0.125	0.678	0.825	-0.299	0.193	121	134
SLF	-0.346	0.126	0.007	0.173	-0.593	-0.098	122	134
SS	-0.148	0.126	0.245	0.511	-0.394	0.098	121	134
UNC	0.081	0.125	0.522	0.686	-0.164	0.327	122	134
AverageAD	-0.244	0.126	0.056	0.343	-0.491	0.002	121	134

Table S34. Differences in regional FA between people with a lifetime history of an actual suicide and those without a history of an actual suicide attempt but with lifetime suicidal ideation determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N ideation	N attempt
ACR	0.113	0.126	0.378	0.865	-0.134	0.359	122	132
ALIC	0.075	0.125	0.558	0.865	-0.171	0.320	122	133
BCC	0.045	0.126	0.727	0.865	-0.202	0.291	122	131
CC	0.037	0.126	0.772	0.877	-0.209	0.283	122	132
CGC	-0.098	0.125	0.440	0.865	-0.344	0.147	122	134
CGH	-0.095	0.125	0.453	0.865	-0.341	0.150	122	134
CR	0.110	0.126	0.389	0.865	-0.136	0.356	122	132
CST	0.117	0.126	0.359	0.865	-0.129	0.363	121	134
EC	0.046	0.125	0.715	0.865	-0.199	0.292	122	133
FX	-0.078	0.126	0.541	0.865	-0.324	0.168	122	132
FXST	-0.123	0.126	0.334	0.865	-0.370	0.123	122	132
GCC	0.051	0.126	0.689	0.865	-0.195	0.297	122	132
IC	0.099	0.125	0.436	0.865	-0.147	0.345	122	133
IFO	-0.070	0.125	0.580	0.865	-0.316	0.175	121	134
PCR	-0.022	0.125	0.865	0.901	-0.267	0.224	122	134
PLIC	0.137	0.126	0.285	0.865	-0.110	0.383	121	133
PTR	-0.182	0.126	0.155	0.865	-0.428	0.065	122	132
RLIC	0.008	0.125	0.947	0.947	-0.237	0.254	122	133
SCC	0.045	0.126	0.725	0.865	-0.202	0.292	122	130
SCR	0.102	0.125	0.423	0.865	-0.144	0.347	122	134
SFO	0.022	0.126	0.862	0.901	-0.224	0.268	121	133
SLF	-0.125	0.125	0.326	0.865	-0.370	0.121	122	134
SS	0.139	0.126	0.275	0.865	-0.107	0.386	122	132
UNC	-0.068	0.125	0.594	0.865	-0.313	0.178	122	134
AverageFA	0.062	0.126	0.628	0.865	-0.184	0.308	122	132

Table S35. Differences in regional MD between people with a lifetime history of an actual suicide and those without a history of an actual suicide attempt but with lifetime suicidal ideation determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N ideation	N attempt
ACR	-0.218	0.126	0.088	0.476	-0.464	0.028	122	134
ALIC	-0.081	0.125	0.526	0.722	-0.327	0.165	122	133
BCC	-0.097	0.126	0.447	0.684	-0.343	0.149	122	132
CC	-0.118	0.125	0.353	0.684	-0.364	0.127	122	134
CGC	-0.168	0.125	0.186	0.476	-0.414	0.077	122	134
CGH	0.066	0.125	0.603	0.722	-0.179	0.311	122	134
CR	-0.169	0.125	0.185	0.476	-0.414	0.077	122	134
CST	-0.011	0.126	0.931	0.948	-0.257	0.235	120	134
EC	-0.198	0.126	0.122	0.476	-0.444	0.049	121	133
FX	0.154	0.126	0.231	0.524	-0.094	0.401	120	132
FXST	-0.039	0.125	0.758	0.824	-0.285	0.207	122	133
GCC	-0.183	0.125	0.151	0.476	-0.428	0.063	122	134
IC	-0.102	0.126	0.426	0.684	-0.348	0.145	121	133
IFO	-0.074	0.125	0.562	0.722	-0.320	0.172	122	133
PCR	-0.110	0.125	0.387	0.684	-0.355	0.136	122	134
PLIC	-0.065	0.126	0.609	0.722	-0.312	0.181	121	133
PTR	-0.174	0.126	0.174	0.476	-0.420	0.073	121	133
RLIC	-0.181	0.126	0.155	0.476	-0.427	0.065	122	133
SCC	-0.060	0.125	0.636	0.722	-0.306	0.185	122	133
SCR	-0.097	0.125	0.444	0.684	-0.343	0.148	122	134
SFO	-0.008	0.125	0.948	0.948	-0.253	0.237	122	134
SLF	-0.175	0.125	0.170	0.476	-0.421	0.071	122	134
SS	-0.218	0.126	0.088	0.476	-0.464	0.029	122	133
UNC	0.093	0.126	0.465	0.684	-0.153	0.339	122	132
AverageMD	-0.167	0.126	0.191	0.476	-0.414	0.079	121	133

Table S36. Differences in regional RD between people with a lifetime history of an actual suicide and those without a history of an actual suicide attempt but with lifetime suicidal ideation determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	Lower CI	Upper CI	N ideation	N attempt
ACR	-0.204	0.126	0.111	0.684	-0.450	0.043	122	132
ALIC	-0.137	0.126	0.284	0.684	-0.383	0.110	122	132
BCC	-0.067	0.126	0.600	0.770	-0.313	0.179	122	132
CC	-0.089	0.126	0.487	0.770	-0.335	0.158	122	132
CGC	-0.127	0.125	0.319	0.684	-0.373	0.119	122	133
CGH	0.052	0.125	0.684	0.770	-0.194	0.297	122	134
CR	-0.145	0.126	0.255	0.684	-0.391	0.101	122	133
CST	-0.072	0.126	0.574	0.770	-0.318	0.175	121	133
EC	-0.163	0.126	0.200	0.684	-0.409	0.083	122	133
FX	0.125	0.126	0.331	0.684	-0.123	0.373	120	131
FXST	0.044	0.125	0.730	0.770	-0.202	0.289	122	134
GCC	-0.121	0.126	0.341	0.684	-0.368	0.125	122	132
IC	-0.166	0.126	0.192	0.684	-0.413	0.080	122	133
IFO	-0.049	0.125	0.703	0.770	-0.294	0.197	121	134
PCR	-0.052	0.125	0.682	0.770	-0.297	0.193	122	134
PLIC	-0.158	0.126	0.217	0.684	-0.404	0.089	121	133
PTR	-0.079	0.126	0.536	0.770	-0.326	0.168	121	132
RLIC	-0.118	0.125	0.356	0.684	-0.364	0.128	122	133
SCC	-0.050	0.126	0.694	0.770	-0.296	0.196	122	132
SCR	-0.138	0.126	0.278	0.684	-0.384	0.108	122	133
SFO	0.042	0.125	0.739	0.770	-0.203	0.288	122	134
SLF	-0.032	0.125	0.804	0.804	-0.277	0.214	122	134
SS	-0.248	0.126	0.052	0.684	-0.495	-0.001	122	132
UNC	0.083	0.125	0.511	0.770	-0.162	0.329	122	134
AverageRD	-0.133	0.125	0.298	0.684	-0.379	0.113	122	133

Table S37. Differences in regional AD between people with a lifetime history an interrupted, aborted or actual attempt and those without a history of an interrupted, aborted or actual suicide attempt determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	N attempt	N no attempt
ACR	-0.074	0.099	0.457	0.722	177	242
ALIC	-0.132	0.099	0.187	0.451	177	241
BCC	-0.245	0.099	0.015	0.147	177	240
CC	-0.235	0.100	0.019	0.147	177	239
CGC	0.007	0.099	0.942	0.942	177	243
CGH	0.056	0.099	0.578	0.753	177	244
CR	-0.088	0.099	0.377	0.674	177	240
CST	-0.064	0.099	0.520	0.722	176	242
EC	-0.144	0.100	0.151	0.420	175	240
FX	-0.044	0.099	0.663	0.753	176	244
FXST	-0.047	0.099	0.640	0.753	177	242
GCC	-0.198	0.099	0.048	0.200	175	244
IC	-0.176	0.100	0.080	0.278	175	239
IFO	-0.069	0.099	0.488	0.722	177	242
PCR	-0.012	0.099	0.905	0.942	177	242
PLIC	-0.109	0.099	0.275	0.528	176	240
PTR	-0.170	0.099	0.089	0.278	177	239
RLIC	-0.219	0.100	0.029	0.147	177	238
SCC	-0.124	0.099	0.216	0.451	176	239
SCR	-0.128	0.099	0.201	0.451	177	238
SFO	-0.065	0.099	0.513	0.722	177	242
SLF	-0.266	0.100	0.008	0.147	177	240
SS	-0.051	0.099	0.612	0.753	176	242
UNC	0.015	0.099	0.882	0.942	177	242
AverageAD	-0.219	0.099	0.029	0.147	177	239

Table S38. Differences in regional FA between people with a lifetime history an interrupted, aborted or actual attempt and those without a history of an interrupted, aborted or actual suicide attempt determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	N attempt	N no attempt
ACR	0.018	0.099	0.858	0.968	175	243
ALIC	-0.020	0.099	0.838	0.968	176	241
BCC	-0.009	0.099	0.930	0.968	174	244
CC	-0.051	0.099	0.608	0.968	175	242
CGC	-0.123	0.099	0.218	0.968	177	244
CGH	-0.056	0.099	0.573	0.968	177	242
CR	0.030	0.099	0.768	0.968	175	242
CST	-0.012	0.099	0.901	0.968	177	242
EC	-0.010	0.099	0.923	0.968	176	243
FX	0.000	0.099	0.999	0.999	175	242
FXST	-0.149	0.099	0.138	0.968	175	241
GCC	0.012	0.099	0.902	0.968	175	243
IC	-0.013	0.099	0.898	0.968	176	241
IFO	0.042	0.099	0.674	0.968	176	243
PCR	-0.020	0.099	0.843	0.968	177	244
PLIC	0.077	0.099	0.441	0.968	176	242
PTR	-0.063	0.099	0.528	0.968	175	242
RLIC	-0.052	0.099	0.606	0.968	176	241
SCC	-0.129	0.100	0.203	0.968	173	237
SCR	0.059	0.099	0.551	0.968	177	244
SFO	-0.143	0.099	0.155	0.968	175	242
SLF	-0.084	0.099	0.398	0.968	177	244
SS	0.049	0.100	0.629	0.968	175	239
UNC	-0.121	0.099	0.225	0.968	177	243
AverageFA	-0.032	0.099	0.746	0.968	175	241



Table S39. Differences in regional MD between people with a lifetime history an interrupted, aborted or actual attempt and those without a history of an interrupted, aborted or actual suicide attempt determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	N attempt	N no attempt
ACR	-0.009	0.099	0.930	0.999	177	242
ALIC	-0.030	0.099	0.767	0.965	176	244
BCC	-0.085	0.099	0.396	0.761	175	244
CC	-0.088	0.099	0.377	0.761	177	244
CGC	-0.017	0.099	0.868	0.999	177	243
CGH	0.031	0.099	0.758	0.965	177	243
CR	-0.005	0.099	0.962	0.999	177	243
CST	-0.048	0.099	0.634	0.950	177	237
EC	-0.092	0.099	0.358	0.761	176	243
FX	-0.095	0.099	0.346	0.761	174	243
FXST	0.061	0.099	0.541	0.902	176	244
GCC	-0.173	0.099	0.082	0.761	177	244
IC	-0.105	0.099	0.292	0.761	175	243
IFO	-0.129	0.099	0.196	0.761	176	244
PCR	0.012	0.099	0.906	0.999	177	243
PLIC	-0.093	0.099	0.352	0.761	175	243
PTR	-0.079	0.099	0.428	0.765	175	242
RLIC	-0.126	0.099	0.207	0.761	176	242
SCC	0.000	0.099	0.999	0.999	176	242
SCR	-0.046	0.099	0.646	0.950	177	242
SFO	0.029	0.099	0.772	0.965	177	243
SLF	-0.109	0.099	0.275	0.761	177	242
SS	-0.114	0.099	0.253	0.761	176	243
UNC	0.089	0.099	0.374	0.761	175	244
AverageMD	-0.095	0.099	0.342	0.761	176	243

Table S40. Differences in regional RD between people with a lifetime history an interrupted, aborted or actual attempt and those without a history of an interrupted, aborted or actual suicide attempt determined with the Columbia Suicide Severity Rating Scale (healthy controls were excluded). D: Cohen's d effect size, SE: standard error; p: p-value, FDR-p: FDR corrected p-value, CI: confidence interval.

Region	D	SE	P	FDR-p	N attempt	N no attempt
ACR	-0.029	0.099	0.772	0.938	175	241
ALIC	-0.093	0.099	0.350	0.919	175	244
BCC	-0.033	0.099	0.740	0.938	175	244
CC	-0.029	0.099	0.771	0.938	175	244
CGC	-0.078	0.099	0.437	0.919	175	244
CGH	0.009	0.099	0.930	0.938	177	243
CR	-0.008	0.099	0.938	0.938	176	241
CST	0.011	0.099	0.912	0.938	176	240
EC	-0.118	0.099	0.236	0.919	176	243
FX	-0.121	0.100	0.228	0.919	173	243
FXST	0.070	0.099	0.484	0.919	177	244
GCC	-0.073	0.099	0.464	0.919	175	243
IC	-0.095	0.099	0.339	0.919	176	243
IFO	-0.117	0.099	0.243	0.919	176	244
PCR	0.013	0.099	0.897	0.938	177	242
PLIC	-0.103	0.099	0.303	0.919	176	241
PTR	-0.047	0.099	0.638	0.938	174	241
RLIC	-0.054	0.099	0.588	0.919	176	242
SCC	0.059	0.099	0.553	0.919	175	242
SCR	-0.071	0.099	0.477	0.919	176	243
SFO	0.056	0.099	0.573	0.919	177	244
SLF	-0.012	0.099	0.906	0.938	177	243
SS	-0.151	0.099	0.133	0.919	175	242
UNC	0.083	0.099	0.403	0.919	177	244
AverageRD	-0.094	0.099	0.344	0.919	176	244

### Supplemental Note 1.

While the main analyses of this manuscript focused on comparing healthy controls and clinical controls to participants with recent suicidal ideation or a lifetime suicide attempt, we also performed supplementary analyses in which we compare WM microstructure between healthy controls and clinical controls. These analyses may not reveal new information related to the neural correlates of suicidality, but may help identify alterations related to having a psychiatric diagnosis and contribute to the interpretation of our main findings. These supplementary analyses were performed similar to the main analyses (with age, sex, age-by-sex, age<sup>2</sup> and age<sup>2</sup>-by-sex included as covariates). In order to compare to our main findings, we compared healthy controls and clinical controls in samples included in the analyses on lifetime suicide attempt ( $N_{CLC} = 1871$ ,  $N_{HC} = 642$ ) and in samples included in the analyses on recent suicidal ideation ( $N_{CLC} = 1184$ ,  $N_{HC} = 1240$ ). Results are presented in supplemental tables S41-48. Results showed no significant differences between in global or regional AD, FA, MD or RD between healthy controls and clinical controls in samples included in the analyses on suicide attempt (supplemental tables S41-44). In samples included in the ideation analyses, we observed lower global AD and regional AD in the ACR, CGC, GCC and SLF in the CLC group compared to HC (Cohen's  $d$  range: 0.112-0.184; supplemental table S45). In addition, we observed lower global FA and regional FA in the GCC in the CLC group compared to HC (Cohen's  $D=0.123$  and  $0.127$  respectively; supplemental table S46).

Table S41 Differences in regional AD between clinical controls and healthy controls in samples included in the analysis on suicide attempt  
Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	-0.056	0.051	-0.155	0.043	0.269	0.770	511	1680
ALIC	-0.085	0.051	-0.184	0.014	0.095	0.770	511	1677
BCC	0.037	0.051	-0.062	0.137	0.461	0.770	507	1679
CC	0.009	0.051	-0.091	0.108	0.863	0.926	508	1676
CGC	-0.084	0.050	-0.183	0.015	0.096	0.770	512	1683
CGH	0.036	0.050	-0.063	0.135	0.478	0.770	512	1682
CR	-0.062	0.051	-0.161	0.037	0.223	0.770	511	1676
CST	0.032	0.051	-0.067	0.131	0.524	0.770	510	1678
EC	-0.057	0.051	-0.156	0.042	0.260	0.770	512	1669
FX	0.050	0.050	-0.049	0.149	0.323	0.770	512	1685
FXST	0.023	0.050	-0.076	0.122	0.652	0.841	512	1683
GCC	-0.005	0.051	-0.104	0.094	0.926	0.926	509	1680
IC	-0.044	0.051	-0.143	0.055	0.387	0.770	510	1675
IFO	-0.028	0.052	-0.130	0.074	0.595	0.827	475	1661
PCR	-0.055	0.051	-0.154	0.045	0.281	0.770	509	1675
PLIC	-0.034	0.051	-0.133	0.065	0.498	0.770	511	1678
PTR	-0.006	0.051	-0.105	0.093	0.911	0.926	511	1677
RLIC	-0.005	0.050	-0.104	0.094	0.924	0.926	512	1676
SCC	0.021	0.051	-0.078	0.120	0.674	0.841	509	1678
SCR	-0.077	0.051	-0.177	0.022	0.127	0.770	510	1671
SFO	-0.039	0.051	-0.138	0.060	0.441	0.770	511	1673
SLF	-0.067	0.050	-0.166	0.032	0.184	0.770	512	1680
SS	0.019	0.051	-0.080	0.118	0.707	0.841	511	1676
UNC	0.059	0.053	-0.046	0.164	0.269	0.770	447	1615
AverageAD	-0.037	0.051	-0.136	0.062	0.461	0.770	511	1672

Table S42 Differences in regional FA between clinical controls and healthy controls in samples included in the analysis on suicide attempt  
 Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	0.007	0.047	-0.086	0.100	0.877	0.913	591	1802
ALIC	-0.033	0.047	-0.126	0.060	0.482	0.753	591	1794
BCC	-0.120	0.048	-0.213	-0.026	0.012	0.075	588	1795
CC	-0.126	0.047	-0.219	-0.033	0.008	0.068	591	1792
CGC	-0.070	0.047	-0.163	0.024	0.144	0.299	589	1797
CGH	-0.024	0.047	-0.117	0.069	0.607	0.779	590	1797
CR	-0.044	0.047	-0.137	0.049	0.357	0.595	590	1795
CST	-0.020	0.048	-0.113	0.073	0.670	0.779	588	1797
EC	-0.091	0.047	-0.184	0.002	0.055	0.165	591	1792
FX	-0.031	0.047	-0.124	0.062	0.512	0.753	589	1792
FXST	-0.070	0.047	-0.163	0.023	0.138	0.299	590	1795
GCC	-0.131	0.047	-0.224	-0.038	0.006	0.068	590	1801
IC	-0.015	0.047	-0.108	0.078	0.750	0.816	589	1794
IFO	-0.118	0.049	-0.214	-0.022	0.016	0.082	546	1770
PCR	-0.094	0.047	-0.187	-0.001	0.048	0.165	591	1797
PLIC	-0.005	0.047	-0.098	0.088	0.924	0.924	590	1793
PTR	-0.060	0.047	-0.153	0.033	0.204	0.364	590	1797
RLIC	-0.020	0.047	-0.113	0.073	0.676	0.779	591	1792
SCC	-0.075	0.048	-0.168	0.018	0.113	0.283	589	1791
SCR	-0.064	0.047	-0.157	0.029	0.180	0.347	591	1798
SFO	0.019	0.048	-0.074	0.112	0.685	0.779	588	1799
SLF	-0.090	0.047	-0.183	0.003	0.059	0.165	590	1802
SS	-0.024	0.047	-0.117	0.069	0.617	0.779	591	1794
UNC	-0.144	0.050	-0.241	-0.046	0.004	0.068	526	1732
AverageFA	-0.107	0.048	-0.200	-0.014	0.025	0.102	589	1793

Table S43 Differences in regional MD between clinical controls and healthy controls in samples included in the analysis on suicide attempt  
Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	-0.003	0.049	-0.098	0.093	0.954	0.954	553	1764
ALIC	-0.045	0.049	-0.141	0.051	0.360	0.853	550	1759
BCC	0.083	0.049	-0.012	0.179	0.088	0.500	549	1760
CC	0.078	0.049	-0.018	0.174	0.111	0.500	550	1764
CGC	-0.027	0.049	-0.123	0.068	0.577	0.853	552	1764
CGH	-0.011	0.049	-0.107	0.085	0.823	0.935	551	1764
CR	0.012	0.049	-0.083	0.108	0.800	0.935	551	1762
CST	0.022	0.049	-0.074	0.117	0.660	0.868	552	1750
EC	0.029	0.049	-0.067	0.125	0.554	0.853	550	1760
FX	0.073	0.049	-0.023	0.169	0.137	0.500	547	1755
FXST	0.051	0.049	-0.045	0.146	0.301	0.853	552	1759
GCC	0.081	0.049	-0.014	0.177	0.095	0.500	553	1767
IC	-0.004	0.049	-0.100	0.092	0.937	0.954	550	1761
IFO	0.037	0.050	-0.062	0.136	0.461	0.853	508	1729
PCR	0.028	0.049	-0.068	0.124	0.565	0.853	550	1757
PLIC	-0.015	0.049	-0.111	0.080	0.753	0.935	549	1762
PTR	0.035	0.049	-0.060	0.131	0.472	0.853	551	1762
RLIC	0.025	0.049	-0.071	0.120	0.614	0.853	551	1764
SCC	0.078	0.049	-0.018	0.174	0.111	0.500	550	1754
SCR	0.008	0.049	-0.088	0.104	0.872	0.948	551	1760
SFO	-0.072	0.049	-0.168	0.023	0.140	0.500	554	1755
SLF	0.025	0.049	-0.071	0.121	0.612	0.853	551	1758
SS	0.040	0.049	-0.055	0.136	0.408	0.853	553	1756
UNC	0.092	0.051	-0.009	0.193	0.075	0.500	487	1692
AverageMD	0.044	0.049	-0.052	0.140	0.368	0.853	549	1763

Table S44 Differences in regional RD between clinical controls and healthy controls in samples included in the analysis on suicide attempt  
Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	0.016	0.049	-0.080	0.111	0.750	0.812	553	1761
ALIC	-0.005	0.049	-0.100	0.091	0.922	0.922	551	1758
BCC	0.089	0.049	-0.007	0.185	0.070	0.630	549	1763
CC	0.074	0.049	-0.022	0.170	0.130	0.630	552	1764
CGC	0.014	0.049	-0.082	0.109	0.779	0.812	552	1762
CGH	0.019	0.049	-0.077	0.114	0.702	0.812	553	1763
CR	0.040	0.049	-0.055	0.136	0.409	0.723	552	1762
CST	0.017	0.049	-0.078	0.113	0.723	0.812	551	1762
EC	0.052	0.049	-0.043	0.148	0.286	0.630	553	1757
FX	0.091	0.049	-0.005	0.187	0.064	0.630	545	1755
FXST	0.049	0.049	-0.046	0.145	0.312	0.630	553	1760
GCC	0.074	0.049	-0.021	0.170	0.129	0.630	553	1762
IC	0.036	0.049	-0.060	0.132	0.462	0.723	551	1761
IFO	0.071	0.050	-0.028	0.170	0.162	0.630	508	1728
PCR	0.048	0.049	-0.048	0.144	0.328	0.630	551	1759
PLIC	0.019	0.049	-0.077	0.115	0.699	0.812	552	1757
PTR	0.020	0.049	-0.076	0.116	0.683	0.812	552	1761
RLIC	0.027	0.049	-0.069	0.122	0.581	0.812	553	1761
SCC	0.058	0.049	-0.038	0.153	0.239	0.630	550	1752
SCR	0.038	0.049	-0.058	0.133	0.439	0.723	552	1763
SFO	-0.058	0.049	-0.153	0.038	0.236	0.630	553	1761
SLF	0.048	0.049	-0.048	0.144	0.326	0.630	552	1762
SS	0.019	0.049	-0.077	0.114	0.699	0.812	554	1757
UNC	0.080	0.051	-0.021	0.181	0.121	0.630	487	1692
AverageRD	0.066	0.049	-0.030	0.161	0.179	0.630	552	1762

Table S45 Differences in regional AD between clinical controls and healthy controls in samples included in the analysis on suicidal ideation  
 Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	-0.132	0.041	-0.213	-0.051	<b>0.001</b>	<b>0.012</b>	1217	1136
ALIC	-0.050	0.041	-0.131	0.031	0.229	0.446	1206	1139
BCC	-0.032	0.041	-0.113	0.049	0.442	0.614	1211	1138
CC	-0.080	0.041	-0.161	0.001	0.054	0.209	1214	1137
CGC	-0.184	0.041	-0.265	-0.103	<b>0.000</b>	<b>0.000</b>	1214	1137
CGH	0.049	0.041	-0.031	0.130	0.232	0.446	1216	1138
CR	-0.078	0.041	-0.159	0.003	0.059	0.209	1216	1136
CST	0.018	0.041	-0.063	0.099	0.666	0.724	1206	1136
EC	-0.072	0.041	-0.153	0.009	0.081	0.228	1217	1134
FX	0.051	0.041	-0.030	0.132	0.219	0.446	1214	1130
FXST	0.008	0.041	-0.073	0.089	0.842	0.877	1219	1140
GCC	-0.113	0.041	-0.194	-0.032	<b>0.006</b>	<b>0.033</b>	1215	1136
IC	-0.032	0.041	-0.113	0.049	0.442	0.614	1214	1139
IFO	0.022	0.041	-0.059	0.103	0.597	0.679	1213	1139
PCR	-0.039	0.041	-0.120	0.042	0.342	0.562	1212	1134
PLIC	-0.023	0.041	-0.104	0.058	0.579	0.679	1216	1139
PTR	-0.023	0.041	-0.104	0.057	0.572	0.679	1217	1137
RLIC	-0.038	0.041	-0.119	0.043	0.360	0.562	1220	1137
SCC	-0.072	0.041	-0.153	0.009	0.082	0.228	1214	1137
SCR	-0.029	0.041	-0.110	0.052	0.478	0.629	1215	1136
SFO	0.006	0.041	-0.075	0.087	0.888	0.888	1211	1132
SLF	-0.112	0.041	-0.193	-0.031	<b>0.007</b>	<b>0.033</b>	1217	1134
SS	0.041	0.041	-0.040	0.122	0.322	0.562	1216	1134
UNC	0.063	0.042	-0.020	0.146	0.139	0.347	1152	1088
AverageAD	-0.157	0.041	-0.239	-0.076	<b>0.000</b>	<b>0.002</b>	1214	1134



Table S46 Differences in regional FA between clinical controls and healthy controls in samples included in the analysis on suicidal ideation  
 Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	-0.061	0.041	-0.141	0.019	0.135	0.303	1236	1182
ALIC	-0.071	0.041	-0.151	0.009	0.083	0.226	1235	1177
BCC	-0.093	0.041	-0.173	-0.013	0.024	0.147	1232	1174
CC	-0.111	0.041	-0.191	-0.032	0.006	0.053	1233	1175
CGC	-0.059	0.041	-0.139	0.020	0.145	0.303	1233	1177
CGH	-0.018	0.041	-0.097	0.062	0.666	0.742	1234	1176
CR	-0.069	0.041	-0.149	0.010	0.089	0.226	1233	1179
CST	-0.022	0.041	-0.102	0.059	0.599	0.742	1223	1176
EC	-0.069	0.041	-0.149	0.011	0.090	0.226	1230	1175
FX	-0.057	0.041	-0.137	0.023	0.162	0.311	1230	1177
FXST	-0.014	0.041	-0.094	0.066	0.736	0.742	1234	1176
GCC	-0.127	0.041	-0.207	-0.047	0.002	0.036	1233	1177
IC	-0.019	0.041	-0.099	0.061	0.640	0.742	1235	1178
IFO	-0.039	0.041	-0.119	0.041	0.342	0.534	1226	1182
PCR	-0.071	0.041	-0.151	0.009	0.083	0.226	1234	1179
PLIC	-0.016	0.041	-0.096	0.063	0.688	0.742	1234	1176
PTR	-0.026	0.041	-0.106	0.053	0.518	0.720	1235	1179
RLIC	0.021	0.041	-0.059	0.101	0.608	0.742	1232	1178
SCC	-0.083	0.041	-0.163	-0.003	0.043	0.217	1230	1176
SCR	-0.052	0.041	-0.132	0.028	0.203	0.363	1234	1180
SFO	-0.040	0.041	-0.120	0.040	0.325	0.534	1232	1180
SLF	-0.075	0.041	-0.154	0.005	0.068	0.226	1233	1183
SS	-0.013	0.041	-0.093	0.066	0.742	0.742	1230	1180
UNC	-0.035	0.042	-0.117	0.047	0.402	0.592	1169	1131
AverageFA	-0.123	0.041	-0.203	-0.042	0.003	0.036	1231	1140

Table S47 Differences in regional MD between clinical controls and healthy controls in samples included in the analysis on suicidal ideation  
 Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	-0.032	0.041	-0.112	0.048	0.435	0.777	1217	1175
ALIC	0.003	0.041	-0.077	0.084	0.940	0.970	1204	1173
BCC	0.070	0.041	-0.010	0.150	0.089	0.507	1213	1174
CC	0.051	0.041	-0.029	0.132	0.210	0.507	1214	1178
CGC	-0.094	0.041	-0.174	-0.014	0.022	0.276	1214	1177
CGH	0.061	0.041	-0.019	0.141	0.137	0.507	1219	1176
CR	0.012	0.041	-0.068	0.092	0.771	0.948	1216	1176
CST	0.011	0.041	-0.070	0.091	0.797	0.948	1203	1170
EC	0.024	0.041	-0.056	0.104	0.562	0.826	1209	1174
FX	0.021	0.041	-0.060	0.101	0.617	0.857	1209	1162
FXST	0.052	0.041	-0.029	0.132	0.208	0.507	1215	1174
GCC	-0.007	0.041	-0.087	0.073	0.863	0.948	1216	1177
IC	-0.007	0.041	-0.087	0.074	0.871	0.948	1212	1173
IFO	0.074	0.041	-0.006	0.154	0.070	0.507	1213	1175
PCR	0.052	0.041	-0.028	0.132	0.202	0.507	1217	1175
PLIC	-0.052	0.041	-0.133	0.028	0.201	0.507	1215	1175
PTR	0.052	0.041	-0.028	0.133	0.200	0.507	1215	1176
RLIC	-0.007	0.041	-0.087	0.074	0.872	0.948	1214	1177
SCC	0.035	0.041	-0.046	0.115	0.399	0.766	1212	1174
SCR	0.050	0.041	-0.030	0.130	0.223	0.507	1214	1173
SFO	0.025	0.041	-0.055	0.106	0.539	0.826	1212	1171
SLF	-0.002	0.041	-0.082	0.079	0.970	0.970	1216	1174
SS	0.045	0.041	-0.035	0.125	0.270	0.564	1219	1171
UNC	0.115	0.042	0.033	0.197	0.006	0.155	1145	1127
AverageMD	0.028	0.041	-0.053	0.109	0.505	0.826	1212	1136

Table S48 Differences in regional RD between clinical controls and healthy controls in samples included in the analysis on suicidal ideation  
Age, sex, age-by-sex, age<sup>2</sup>, age<sup>2</sup>-by-sex included as covariates.

Region	Cohen's d	SE	CI LB	CI UB	P-value	FDR P-value	HC	CLC
ACR	0.006	0.041	-0.075	0.087	0.883	0.920	1214	1135
ALIC	0.022	0.041	-0.059	0.103	0.593	0.743	1206	1133
BCC	0.091	0.041	0.010	0.172	0.028	0.421	1211	1137
CC	0.072	0.041	-0.009	0.153	0.081	0.421	1214	1135
CGC	-0.033	0.041	-0.114	0.048	0.425	0.626	1213	1136
CGH	0.048	0.041	-0.033	0.129	0.244	0.611	1221	1135
CR	0.039	0.041	-0.042	0.120	0.350	0.622	1215	1135
CST	0.038	0.041	-0.043	0.119	0.357	0.622	1202	1137
EC	0.037	0.041	-0.044	0.118	0.374	0.622	1213	1132
FX	0.042	0.041	-0.040	0.123	0.315	0.622	1206	1122
FXST	0.015	0.041	-0.066	0.096	0.712	0.847	1215	1136
GCC	0.040	0.041	-0.041	0.121	0.329	0.622	1215	1134
IC	-0.008	0.041	-0.089	0.073	0.846	0.920	1213	1136
IFO	0.068	0.041	-0.013	0.149	0.101	0.421	1211	1133
PCR	0.048	0.041	-0.033	0.129	0.243	0.611	1213	1136
PLIC	-0.035	0.041	-0.116	0.046	0.398	0.622	1215	1135
PTR	0.053	0.041	-0.028	0.134	0.200	0.611	1214	1136
RLIC	-0.022	0.041	-0.103	0.059	0.594	0.743	1215	1137
SCC	0.073	0.041	-0.008	0.154	0.078	0.421	1207	1133
SCR	0.058	0.041	-0.023	0.139	0.158	0.563	1215	1134
SFO	0.007	0.041	-0.074	0.088	0.870	0.920	1209	1135
SLF	0.025	0.041	-0.056	0.106	0.547	0.743	1214	1136
SS	-0.002	0.041	-0.083	0.078	0.953	0.953	1219	1132
UNC	0.075	0.042	-0.009	0.158	0.079	0.421	1143	1087
AverageRD	0.071	0.041	-0.010	0.152	0.084	0.421	1214	1136

## Supplemental note 2.

The ENIGMA-Suicidal Thoughts and Behaviours consortium is transdiagnostic, and includes participants with various types of psychiatric illness, including major depressive disorder, bipolar disorder, anxiety disorders, obsessive-compulsive disorder, posttraumatic stress disorder and psychotic disorders. We were interested to examine if the association between suicidality and WM microstructure differed according to diagnosis type. In supplemental analyses, we aimed to examine interaction effects between group (clinical controls and recent suicidal ideation or clinical controls and lifetime suicide attempt) and type of lifetime psychiatric disorder. Healthy controls were not included in this analysis as they did not have a psychiatric disorder. As information on lifetime type of psychiatric disorder was only available for a subsample of participants (some sites only assessed current disorders), these analyses were performed as supplemental analyses.

Lifetime diagnosis type was categorized into six main lifetime diagnosis types: MDD, BD, anxiety disorders, PTSD, OCD and psychotic disorders. In the suicide attempt-by-diagnosis interaction analyses, we were only able to include MDD (N=1139, of whom 355 attempted suicide), BD (N=553; of whom 141 attempted suicide) and psychotic disorders (N=198; of whom 44 attempted suicide) given the small sample sizes per lifetime diagnosis type. Similarly, in the recent suicidal ideation-by-diagnosis interaction analyses, we were only able to include MDD (N=1128 of whom 424 had recent suicidal ideation) and BD (N=225; of whom 114 had recent suicidal ideation).

Similar to our main analyses, we used multiple linear regression models in R, in which we included the group-by-diagnosis type interaction term, and main effects of group and diagnosis type. Results of these analyses are presented in supplemental tables S49-56. Results show that there are no significant interaction effects between suicide attempt group (clinical control or lifetime suicide attempt) and diagnosis type (MDD, BD or psychotic disorder) on WM microstructure (supplemental tables S49-52). In contrast, we observed many significant interaction effects between suicidal ideation group (clinical control or recent suicidal ideation) and diagnosis type (MDD or BD) on WM microstructure, specifically on regional FA, MD or RD (supplemental tables 53-56). These findings indicate that the association between ideation and these WM microstructure metrics differs in MDD patients and BD patients. To follow-up on this, we performed stratified analyses, looking at the effect of recent suicidal ideation (clinical controls vs. recent ideation) in MDD patients and BD patients separately (with age, sex, age-by-sex, age<sup>2</sup> and age<sup>2</sup>-by-sex included as covariates) for regions where we found significant interaction effects. Results of these stratified analyses are presented in supplemental table S57. For most of these associations, the strength of the association appeared to differ between MDD and BD patients, but the associations were not significant in the MDD only sample or BD only sample. Plotting the results showed that the difference between clinical controls and ideators appeared larger in the BD group, than in the MDD group.

Table S49. Interaction of attempt group-by-type of psychiatric diagnosis on regional AD measures

Region	P-value	FDR P-value	Attempt	CLC
ACR	0.151	0.497	503	1240
ALIC	0.407	0.848	502	1238
BCC	0.490	0.942	501	1238
CC	0.293	0.740	502	1238
CGC	0.759	0.959	505	1244
CGH	0.122	0.497	502	1244
CR	0.853	0.959	503	1236
CST	0.148	0.497	503	1239
EC	0.730	0.959	503	1233
FX	0.159	0.497	504	1245
FXST	0.022	0.281	505	1242
GCC	0.021	0.281	503	1242
IC	0.882	0.959	502	1236
IFO	0.075	0.497	503	1243
PCR	0.328	0.745	503	1236
PLIC	0.840	0.959	503	1239
PTR	0.155	0.497	504	1236
RLIC	0.296	0.740	504	1236
SCC	0.775	0.959	502	1241
SCR	0.973	0.973	503	1231
SFO	0.622	0.959	503	1238
SLF	0.878	0.959	505	1240
SS	0.577	0.959	504	1239
UNC	0.850	0.959	495	1220
AverageAD	0.949	0.973	505	1232

Table S50. Interaction of attempt group-by-type of psychiatric diagnosis on regional FA measures

Region	P-value	FDR P-value	Attempt	CLC
ACR	0.377	0.884	525	1286
ALIC	0.442	0.884	523	1280
BCC	0.441	0.884	521	1283
CC	0.201	0.839	521	1278
CGC	0.652	0.884	526	1286
CGH	0.773	0.884	524	1283
CR	0.621	0.884	525	1280
CST	0.720	0.884	523	1283
EC	0.506	0.884	525	1279
FX	0.285	0.884	520	1279
FXST	0.720	0.884	524	1282
GCC	0.085	0.597	524	1285
IC	0.486	0.884	525	1279
IFO	0.038	0.475	526	1283
PCR	0.778	0.884	526	1282
PLIC	0.441	0.884	524	1280
PTR	0.767	0.884	521	1284
RLIC	0.967	0.981	523	1280
SCC	0.027	0.475	522	1278
SCR	0.981	0.981	524	1283
SFO	0.920	0.981	521	1283
SLF	0.294	0.884	522	1286
SS	0.172	0.839	523	1278
UNC	0.096	0.597	516	1263
AverageFA	0.748	0.884	525	1278

Table S51. Interaction of attempt group-by-type of psychiatric diagnosis on regional MD measures

Region	P-value	FDR P-value	Attempt	CLC
ACR	0.024	0.218	511	1264
ALIC	0.799	0.868	508	1262
BCC	0.942	0.942	509	1263
CC	0.385	0.561	511	1267
CGC	0.215	0.518	510	1267
CGH	0.375	0.561	509	1268
CR	0.107	0.447	512	1264
CST	0.315	0.561	508	1258
EC	0.142	0.506	510	1263
FX	0.067	0.336	509	1256
FXST	0.028	0.218	511	1265
GCC	0.005	0.117	513	1269
IC	0.375	0.561	508	1262
IFO	0.849	0.884	509	1261
PCR	0.182	0.506	511	1260
PLIC	0.592	0.672	506	1263
PTR	0.174	0.506	511	1264
RLIC	0.365	0.561	512	1266
SCC	0.559	0.672	511	1261
SCR	0.588	0.672	513	1262
SFO	0.427	0.561	509	1260
SLF	0.410	0.561	510	1260
SS	0.035	0.218	510	1262
UNC	0.271	0.561	499	1241
AverageMD	0.228	0.518	512	1264

Table S52. Interaction of attempt group-by-type of psychiatric diagnosis on regional RD measures

Region	P-value	FDR P-value	Attempt	CLC
ACR	0.028	0.231	511	1262
ALIC	0.608	0.799	507	1263
BCC	0.341	0.609	509	1265
CC	0.119	0.430	510	1267
CGC	0.210	0.478	510	1267
CGH	0.985	0.985	510	1268
CR	0.169	0.430	511	1263
CST	0.474	0.698	507	1266
EC	0.156	0.430	511	1263
FX	0.064	0.401	508	1256
FXST	0.794	0.863	513	1267
GCC	0.002	0.059	510	1264
IC	0.922	0.960	510	1264
IFO	0.373	0.622	513	1259
PCR	0.294	0.565	513	1263
PLIC	0.752	0.855	508	1262
PTR	0.403	0.630	510	1263
RLIC	0.721	0.855	511	1265
SCC	0.128	0.430	510	1260
SCR	0.673	0.841	511	1265
SFO	0.536	0.745	509	1264
SLF	0.274	0.565	511	1264
SS	0.026	0.231	510	1264
UNC	0.090	0.430	502	1242
AverageRD	0.172	0.430	511	1264



Table S53. Interaction of ideation group-by-type of psychiatric diagnosis on regional AD measures

Region	P	FDR-P	Ideation	CLC
ACR	0.336	0.621	502	769
ALIC	0.398	0.621	499	773
BCC	0.027	0.167	499	769
CC	0.056	0.215	502	770
CGC	0.060	0.215	502	772
CGH	0.365	0.621	498	771
CR	0.777	0.809	501	769
CST	0.303	0.621	498	768
EC	0.033	0.167	496	769
FX	0.000	0.000	501	766
FXST	0.019	0.161	501	772
GCC	0.381	0.621	501	769
IC	0.723	0.786	499	772
IFO	0.848	0.848	502	772
PCR	0.693	0.786	500	768
PLIC	0.077	0.242	498	772
PTR	0.434	0.638	499	770
RLIC	0.712	0.786	498	771
SCC	0.354	0.621	503	771
SCR	0.542	0.713	500	769
SFO	0.158	0.438	499	768
SLF	0.506	0.703	503	768
SS	0.681	0.786	501	770
UNC	0.011	0.140	497	756
AverageAD	0.392	0.621	499	767

Table S54. Interaction of ideation group-by-type of psychiatric diagnosis on regional FA measures

Region	P	FDR-P	Ideation	CLC
ACR	0.000	0.000	537	813
ALIC	0.375	0.395	537	810
BCC	0.001	0.004	534	810
CC	0.000	0.001	535	808
CGC	0.290	0.362	537	813
CGH	0.379	0.395	536	810
CR	0.000	0.001	536	810
CST	0.378	0.395	535	809
EC	0.031	0.052	537	809
FX	0.001	0.004	535	812
FXST	0.001	0.003	535	811
GCC	0.000	0.000	536	809
IC	0.061	0.090	534	810
IFO	0.905	0.905	534	814
PCR	0.007	0.015	536	811
PLIC	0.152	0.200	531	809
PTR	0.000	0.001	535	811
RLIC	0.044	0.069	536	811
SCC	0.019	0.037	536	809
SCR	0.013	0.027	535	812
SFO	0.368	0.395	534	814
SLF	0.021	0.037	537	814
SS	0.001	0.002	534	811
UNC	0.073	0.102	532	798
AverageFA	0.004	0.011	509	771

Table S55. Interaction of ideation group-by-type of psychiatric diagnosis on regional MD measures

Region	P	FDR-P	Ideation	CLC
ACR	0.003	0.009	528	808
ALIC	0.342	0.389	524	807
BCC	0.001	0.003	527	807
CC	0.002	0.009	527	810
CGC	0.008	0.019	526	810
CGH	0.888	0.888	527	810
CR	0.003	0.009	527	809
CST	0.051	0.086	523	805
EC	0.000	0.001	523	809
FX	0.000	0.000	528	799
FXST	0.364	0.395	528	809
GCC	0.000	0.003	529	809
IC	0.264	0.330	525	806
IFO	0.295	0.352	526	807
PCR	0.017	0.032	526	807
PLIC	0.583	0.608	523	808
PTR	0.013	0.027	525	808
RLIC	0.072	0.106	528	809
SCC	0.141	0.186	526	807
SCR	0.004	0.012	528	807
SFO	0.059	0.092	524	805
SLF	0.008	0.019	526	807
SS	0.027	0.047	525	807
UNC	0.000	0.003	521	794
AverageMD	0.098	0.136	500	769

Table S56. Interaction of ideation group-by-type of psychiatric diagnosis on regional RD measures

Region	P	FDR-P	Ideation	CLC
ACR	0.000	0.000	501	767
ALIC	0.258	0.281	499	768
BCC	0.002	0.006	499	769
CC	0.000	0.002	500	768
CGC	0.068	0.090	499	770
CGH	0.618	0.644	500	769
CR	0.000	0.000	500	767
CST	0.121	0.144	498	770
EC	0.000	0.001	500	767
FX	0.000	0.000	502	759
FXST	0.014	0.026	501	770
GCC	0.001	0.002	503	766
IC	0.021	0.035	498	769
IFO	0.657	0.657	500	765
PCR	0.007	0.014	500	769
PLIC	0.186	0.212	497	768
PTR	0.004	0.009	500	768
RLIC	0.057	0.079	501	768
SCC	0.040	0.059	501	766
SCR	0.001	0.002	501	767
SFO	0.116	0.144	498	770
SLF	0.001	0.002	502	768
SS	0.003	0.008	500	768
UNC	0.024	0.037	496	755
AverageRD	0.003	0.008	502	768

Table S57. Follow-up stratified analyses: association between group (clinical controls or recent suicidal ideation) and brain metrics that showed significant interaction effects in MDD patients only or in BD patients only.

Measure	Region	MDD only			BD only		
		Cohen's D	SE	P-value	Cohen's D	SE	P-value
AD	FX	-0.119	0.064	0.065	-0.127	0.134	0.351
FA	ACR	-0.043	0.062	0.483	0.042	0.133	0.757
FA	BCC	0.042	0.062	0.499	0.039	0.134	0.774
FA	CC	0.009	0.062	0.885	0.000	0.134	1.000
FA	CR	-0.089	0.062	0.149	0.041	0.133	0.762
FA	FX	0.060	0.062	0.328	0.253	0.134	0.063
FA	FXST	-0.079	0.062	0.201	-0.098	0.134	0.469
FA	GCC	-0.008	0.062	0.903	0.007	0.134	0.956
FA	PCR	-0.090	0.062	0.146	-0.068	0.134	0.616
FA	PTR	-0.125	0.062	0.043	-0.107	0.134	0.433
FA	SCC	0.029	0.062	0.636	-0.103	0.134	0.451
FA	SCR	-0.064	0.062	0.299	0.080	0.133	0.556
FA	SLF	-0.051	0.062	0.411	-0.001	0.133	0.992
FA	SS	-0.078	0.062	0.206	-0.056	0.134	0.683
FA	AverageFA	-0.076	0.064	0.236	-0.001	0.133	0.997
MD	ACR	0.016	0.137	0.798	0.121	0.134	0.373
MD	BCC	-0.090	0.032	0.149	0.186	0.135	0.176
MD	CC	-0.086	0.036	0.168	0.101	0.134	0.458
MD	CGC	0.023	0.145	0.707	0.254	0.134	0.063
MD	CR	0.012	0.133	0.847	0.153	0.134	0.263
MD	EC	-0.021	0.101	0.737	0.070	0.134	0.608
MD	FX	-0.133	-0.011	0.033	-0.098	0.134	0.472
MD	GCC	-0.035	0.086	0.573	0.132	0.134	0.331
MD	PCR	-0.001	0.120	0.984	0.183	0.134	0.179
MD	PTR	0.022	0.144	0.720	0.218	0.134	0.109
MD	SCR	0.023	0.144	0.716	0.169	0.134	0.214
MD	SLF	-0.009	0.113	0.884	0.169	0.134	0.216
MD	SS	-0.023	0.099	0.711	0.143	0.134	0.293
MD	UNC	-0.042	0.080	0.505	0.334	0.139	0.018
RD	ACR	0.050	0.064	0.440	0.095	0.134	0.486
RD	BCC	-0.071	0.064	0.271	0.056	0.134	0.684
RD	CC	-0.055	0.064	0.395	0.097	0.135	0.479
RD	CR	0.062	0.064	0.334	0.181	0.134	0.184

RD	EC	0.025	0.064	0.699	0.028	0.135	0.838
RD	FX	-0.119	0.064	0.065	-0.144	0.134	0.289
RD	FXST	0.110	0.064	0.087	0.194	0.134	0.154
RD	GCC	0.005	0.064	0.937	0.042	0.134	0.758
RD	IC	0.035	0.064	0.591	0.122	0.134	0.370
RD	PCR	0.063	0.064	0.329	0.149	0.134	0.275
RD	PTR	0.087	0.064	0.178	0.142	0.134	0.298
RD	SCR	0.079	0.064	0.217	0.091	0.134	0.505
RD	SLF	0.033	0.064	0.610	0.182	0.134	0.182
RD	SS	0.069	0.064	0.284	0.135	0.134	0.321
RD	UNC	0.018	0.064	0.786	0.223	0.138	0.112
RD	AverageRD	0.033	0.064	0.610	0.143	0.134	0.294