

Supplement Table 1: Definitions of covariates and diseases variables in UK Biobank

	HES statistics ICD-10 and ICD9 codes	HES statistics OPCS4 codes	Self-reported Data-Field 20002	Self-reported operations Data-Field 20004	Other data fields
Type-2 diabetes	E10, E11, E13, E14		1223		Field 2443 = 1
Constipation	K590		1599		Field 6154 = 6
Depression	F33		1286		
Bladder dysfunction	N31		1202		
Sleep duration					Field 1160
Sleeplessness					Field 1200 = 3
Erectile dysfunction			1518		
Parkinson's disease	G20		1262		
<i>Prevalent neurodegenerative disease</i>					
Extrapyramidal and movement disorders	G20-G26		1262		
Other degenerative diseases of the nervous system	G30-G32				
Multi-system degeneration of the autonomic nervous system	G903				
<i>Prevalent Cardiovascular disease</i>					
Ischemic Heart Disease	411.1, 410.11, 410.41, 410.81, 410.91, 410.71, 410.01, 410.21, 410.31, 410.51, 410.61, 411.89, 429.2, 414.0, 414.10, 414.19, 414.8, 414.9, I20-I25	K40, K41, K42, K44, K45, K49, K50, K75		1070, 1095, 1523	
Cardiomyopathies	425.4, 425.11, 425.18, 425, 425.3, 425.5, 425.9, 425.2, 425.8, 425.7, I42-I43				
Heart Failure	428.0, 428.1, 428.9, I13.0, I13.2, I50	K59.6, K61.7, K60.7, K576,			
Arrhythmias	427.3, 427.31, 427.32, 427.1, 427.9, 427.5, I48, I47.2, I47.0, I49.0, I46.0, I46.1, I49.0, I44.1, I44.2, I49.9, G45	K641, X504, X504			
Stroke	434.91, 434.01, 434.11, 436, I63, I61, I64				
Other cardiovascular diseases	I70, I73, I74, I71, I11, I34, I35, I36, I37, Q20, Q21, Q22, Q23, Q24, B33.2, I40, I41, I51.4	K59, K72			
<i>Active cancer diagnosis</i>					
Malignant neoplasms	C00-C97				

Supplemental Table 2
Characteristics of UK Biobank participants with and without consent for the exercise test

	Consent for exercise test			P
	Yes (n=96 524; 19.2%)	No (n=405 840; 80.8%)	N	
Sex, male, %	46.1%	45.5%	502 364	<0.001
Age, years, mean (SD)	56.9 (8.0)	56.4 (8.1)	502 364	<0.001
Ethnicity, white, %	92.1%	95.2%	499 587	<0.001
BMI, mean (SD)	27.3 (4.8)	27.5 (4.8)	499 259	<0.001
Townsend deprivation index ^a , mean (SD)	-1.2 (3.0)	-1.3 (3.1)	501 740	<0.001
Type 2 Diabetes Mellitus, yes, %	5.5%	5.4%	502 364	0.13
Current Smoking, yes, %	9.1%	11.0%	499 415	<0.001
Constipation, yes, %	3.7%	3.8%	502 364	0.38
Depression, yes, %	5.9%	6.1%	502 364	0.017
Bladder dysfunction, %	0.9 %	0.5 %	502 364	<0.001
Erectile dysfunction, %	0.4 %	0.1 %	229 067	<0.001
Sleep duration, hours, mean (SD)	7.1 (1.2)	7.1 (1.3)	501 476	0.025
Sleeplessness, %	27.0%	28.6%	499 984	<0.001

There are statistically significant but clinically irrelevant differences in baseline characteristics between the 96,524 individuals who provided consent for the exercise test, and the remaining part of the UK Biobank cohort. *BMI = Body Mass Index, SD = standard deviation.*

^a *Townsend deprivation index is a measure of material deprivation.*

Supplemental Table 3**Characteristics of participants consenting for the exercise test in relation to eligibility for present study**

	Included in study			P
	Yes (n=69 288; 71.8%)	No (n=27 236; 28.2%)	N	
Sex, male, %	47.6%	42.4%	96 524	<0.001
Age, years, mean (SD)	56.0 (8.0)	59.1 (7.5)	96 524	<0.001
Ethnicity, white, %	93.1%	89.5%	95 896	<0.001
BMI, mean (SD)	27.0 (4.4)	28.2 (5.5)	96 178	<0.001
Townsend deprivation index ^a , mean (SD)	-1.4 (2.9)	-0.8 (3.2)	96 420	<0.001
Type 2 Diabetes Mellitus, yes, %	4.0%	9.5%	96 524	<0.001
Current Smoking, yes, %	8.5%	10.5%	95 938	<0.001
Constipation, yes, %	2.9%	5.8%	96 524	<0.001
Depression, yes, %	5.2%	7.8%	96 524	<0.001
Bladder dysfunction, %	0.8%	1.4%	96 524	<0.001
Erectile dysfunction, %	0.4%	0.5%	44 498	0.032
Sleep duration, hours, mean (SD)	7.1 (1.1)	7.1 (1.5)	96 308	0.52
Sleeplessness, %	24.8%	32.8%	96 053	<0.001

BMI = Body Mass Index, SD = standard deviation. ^a Townsend deprivation index is a measure of material deprivation.

Supplemental Table 4**Hazard ratios of characteristics at baseline – univariate associations with incident PD**

	Hazard Ratio (95% CI)	p
Sex, male	2.5 (2.0-3.2)	<0.001
Age, per year increase	1.1 (1.1-1.2)	<0.001
Ethnicity, white	1.6 (1.0-2.7)	0.070
Townsend deprivation index ^a , per unit increase	1.0 (0.9-1.0)	0.039
BMI, per unit increase	1.0 (1.0-1.1)	0.016
Type 2 Diabetes Mellitus, yes	3.6 (2.6-4.9)	<0.001
Current Smoking, yes	0.4 (0.2-0.7)	0.001
Constipation, yes	2.1 (1.3-3.3)	0.002
Depression, yes	1.7 (1.2-2.5)	0.006
Bladder dysfunction, yes	1.8 (0.8-4.1)	0.15
Sleep duration, per hour increase	1.2 (1.1-1.3)	0.004
Sleeplessness, yes	0.9 (0.7-1.2)	0.40
HRI-exc, per 10 beats less increase	1.3 (1.2-1.4)	<0.001
HRD-rec, per 10 beats less recovery	1.6 (1.4-1.7)	<0.001

^a *Townsend deprivation index is a measure of material deprivation.*

BMI = Body Mass Index

HRI-exc = Heart rate increase during exercise

HRD-rec = Heart rate decrease during recovery

Supplemental Table 5
Associations of clinical and autonomic exercise parameters with incident PD:
multivariate analysis

	Hazard Ratio (95% CI)	P
Sex, male	2.3 (1.8-2.9)	<0.001
Age, per year increase	1.1 (1.1-1.2)	<0.001
Ethnicity, white	1.0 (0.6-1.8)	0.88
BMI, per unit increase	1.0 (1.0-1.0)	0.86
Townsend deprivation index ^a , per unit increase	1.0 (1.0-1.0)	0.98
Type 2 Diabetes Mellitus, yes	2.2 (1.6-3.1)	<0.001
Current Smoking, yes	0.4 (0.2-0.8)	0.005
Constipation, yes	2.1 (1.3-3.3)	0.003
Depression, yes	2.0 (1.4-3.0)	0.001
Bladder dysfunction, yes	1.2 (0.5-2.7)	0.65
Sleep duration, per hour	1.0 (0.9-1.2)	0.38
Sleeplessness, yes	0.8 (0.6-1.1)	0.17
HRD-rec, per 10 beats less recovery	1.3 (1.1-1.4)	<0.001

^a *Townsend deprivation index is a measure of material deprivation.*

BMI = Body Mass Index,

HRD-rec = Heart rate decrease during recovery phase after exercise

Supplemental Table 6**Adjusted hazard ratios of incident PD according to heart rate decrease during recovery in prespecified sensitivity analyses**

Pre-specified sensitivity analyses			
Marker	N	HR per 10 bpm (95% CI)	P Value
Original analysis			
HRD-rec	69,288	1.3 (1.1 - 1.4)	<0.001
Exclusion of individuals using heart rate modulating medication			
HRD-rec	65,802	1.3 (1.1 - 1.5)	<0.001
Exclusion of individuals using dopaminergic and/or (other) psychoactive medication			
HRD-rec	64,578	1.3 (1.2 - 1.5)	<0.001
Exclusion of individuals with missing data			
HRD-rec	66,282	1.3 (1.1 - 1.4)	<0.001

Medication data (UK Biobank Data Field: 20003) were mapped to the Anatomical Therapeutic Chemical (ATC) classification system (<https://github.com/PhilAppleby/ukbb-srmed>). Heart rate modulating medication: beta and calcium blockers, ATC codes: C07, C08; psychoactive medication: ATC codes: N04, N05, N06, N07. Analyses were adjusted for both clinical variables (age, sex, ethnicity, body mass index (BMI), Townsend deprivation index, type-2 diabetes, and smoking) and prodromal autonomic variables (constipation, sleep duration, sleeplessness, depression, and bladder dysfunction).

HRD-rec = heart rate decrease during recovery. Hazard ratios (HR) and confidence intervals (CI) are expressed per 10 beat/min impairment

Supplemental Table 7**Adjusted hazard ratios of incident PD according to heart rate decrease during recovery in post-hoc sensitivity analyses**

Post-hoc sensitivity analyses			
Analysis stratified by workload^a			
HRD-rec only considering subjects exercising at 50% of maximum predicted workload	58,728	1.3 (1.1 – 1.5)	0.001
HRD-rec only considering subjects exercising at 35% of maximum predicted workload	8381	1.2 (0.8 – 1.6)	0.829
Imposing longer lag time to diagnosis			
HRD-rec only considering cases after median follow-up time (9.3 yrs)	67,252	1.3 (1.1 – 1.5)	0.007
HRD-rec only considering cases after 75th % of follow-up time (11.0 yrs)	54,417	1.2 (0.9 – 1.5)	0.163
Incident PD defined as new out-of-hospital and/or in-hospital diagnosis of PD^b			
HRD-rec	37,458	1.2 (1.0 – 1.5)	0.013

^a The interaction term between HRD-rec and workload (35% or 50% of maximum predicted workload) was $p = 0.548$.

^a Incident PD defined by self-reported data (see Supplemental Table 1), primary care data (diagnostic Read code F12..00 for Parkinson's disease), and/or hospitalization (ICD 10 G20); there were 171 cases (0.5%). Median time to out-of-hospital diagnosis was 4.07 years (IQR: 2.25 – 5.09).