Risk factor	TB Confirmed (n)	RR-TB (%)	OR (95% CI)	P value	AOR (95% CI)	P value
Age						
18-29	382	40 (10.5)				
30-44	486	63 (13.0)	1.31 (0.86, 2.04)	.215	1.18 (0.76, 1.85)	.462
45-60	169	16 (9.5)	1.07 (0.56, 1.99)	.830	0.97 (0.50, 1.82)	.928
≥ 60	52	5 (9.6)	1.44 (0.46, 3.72)	.482	1.32 (0.42, 3.45)	.597
Sex						
Male	1081	123 (11.4)	0.20 (0.01, 5.14)	.260	0.15 (0.01, 3.99)	.192
Female	8	1 (12.5)				
TB History						
Yes	397	65 (16.4)	2.00 (1.36, 2.96)	<.001	1.96 (1.32, 2.91)	.001
No	692	59 (8.5)				
TB Contact	556	56 (10.1)	0.61 (0.32, 1.11)	.119	0.59 (0.31, 1.08)	.103
Peru Origin	1054	122 (11.6)	2.72 (0.80, 17.02)	.176	2.25 (0.65, 14.21)	.279

Supplemental Table. Multivariable logistic regression analysis of demographic characteristics and risk factors for rifampicin-resistant TB among PDL with confirmed TB.

Data are presented as no. (%) unless otherwise indicated. CI, confidence interval; PDL, persons deprived of liberty; TB, tuberculosis; RR-TB, rifampicin-resistant tuberculosis; OR, odds ratio; AOR, adjusted odds ratio; PDL, persons deprived of liberty.