Supporting Information

Supporting materials

Supplementary figures



Figure S1. Significant interaction between LTL and *APOE*- ϵ 4 status (P*int* = 0.047) on the association with CSF YKL-40 levels at baseline (N = 346). Model was adjusted for age and sex.



FigureS2. Significant interaction between LTL and *APOE*- ϵ 4 status (P*int* = 0.036) on the association with the change in CSF sTREM2 levels over time (N = 237). Model was adjusted for age and sex.



FigureS3. Significant associations between LTL and log-transformed CSF biomarkers levels at baseline in APOE- ϵ 4 carriers (N = 184). Models were adjusted for age and sex.



FigureS4. Interaction between LTL and AT status on CSF A β 42/40 levels at baseline (A-T+ vs. A-T-: Pint = 0.002; A+T- vs. A-T-: P = 0.09) (N = 346). Model was adjusted for age, sex and *APOE*- ϵ 4.



FigureS5. Significant associations between LTL and change in (A) CSF α -synuclein and (B) CSF IL-6 levels in the A-T- group. Models were adjusted for age, sex, *APOE*- ϵ 4 status and time difference between the two visits.



FigureS6. Significant associations between LTL and baseline levels of (A) CSF NfL, (B) CSF S100B, and (C) CSF YKL-50 at baseline in the A+T- group. Models were adjusted for age, sex, *APOE*-ε4 status.



FigureS7. Significant associations between LTL and baseline levels of (A) CSF t-tau, (B) CSF α -synuclein, (C) CSF sTREM and change over time of (D) CSF sTREM2 in the A+T+ group. Models were adjusted for age, sex and *APOE*- ϵ 4 status. Model was further adjusted for time difference between visits for the change in CSF biomarkers levels over time.



FigureS8. Mediator role of CSF glial and inflammatory biomarkers on the association between leukocyte telomere length and aging and AD cortical thickness signatures. Models were adjusted for age, sex, *APOE*- ϵ 4 status and firmware MRI version. All β are standardized coefficients.