

Supplementary Table 1. Patient scenarios used in the study

Scenario No.	Mobility issues	Cognitive ability and personality (Does not call nurses)	Urinary issues	Presence of tubes or drains	Medication use	Age
1	Need assistance in transfer	No cognitive impairment, but reserved and does not call nurses when necessary	Nocturia	No tube or drain inserted	Sleeping pill	Less than 65 years old
2	Independent	No cognitive impairment and calls nurses when necessary	Nocturia	One tube or drain inserted	Sleeping pill	80 years old and over
3	Walks alone but unstable	No cognitive impairment, but reserved and does not call nurses when necessary	Nocturia	One tube or drain inserted	Anti-hypertensive	65-79 years old
4	Walks alone but unstable	Impaired cognition and does not call nurses when necessary	No urinary problems	One tube or drain inserted	Sleeping pill	Less than 65 years old
5	Walks alone but unstable	No cognitive impairment and calls nurses when necessary	No urinary problems	Multiple tubes and/or drains inserted	Anti-hypertensive	80 years old and over
6	Needs assistance in transfer	No cognitive impairment and calls nurses when necessary	Frequency	One tube or drain inserted	Anti-hypertensive	Less than 65 years old
7	Walk alone but unstable	Impaired cognition and does not call nurses when necessary	Nocturia	No tube or drain inserted	Antibiotic	80 years old and over
8	Needs assistance in transfer	Impaired cognition and does not call nurses when necessary	Frequency	Multiple tubes and/or drains inserted	Sleeping pill	80 years old and over
9	Independent	No cognitive impairment, but reserved and does not call nurses	Frequency	No tube or drain inserted	Anti-hypertensive	80 years old and over

		when necessary				
10	Walks alone but unstable	No cognitive impairment, but reserved and does not call nurses when necessary	Frequency	Multiple tubes and/or drains inserted	Antibiotic	Less than 65 years old
11	Walks alone but unstable	No cognitive impairment and calls nurses when necessary	Frequency	No tube or drain inserted	Sleeping pill	65-79 years old
12	Needs assistance in transfer	No cognitive impairment and calls nurses when necessary	Nocturia	Multiple tubes and/or drains inserted	Antibiotic	65-79 years old
13	Independent	No cognitive impairment, but reserved and does not call nurses when necessary	No urinary problems	Multiple tubes and/or drains inserted	Sleeping pill	65-79 years old
14	Independent	Impaired cognition and does not call nurses when necessary	Frequency	One tube or drain inserted	Antibiotic	65-79 years old
15	Needs assistance in transfer	No cognitive impairment, but reserved and does not call nurses when necessary	No urinary problems	One tube or drain inserted	Antibiotic	80 years old and over
16	Needs assistance in transfer	Impaired cognition and does not call nurses when necessary	No urinary problems	No tube or drain inserted	Anti-hypertensive	65-79 years old
17	Needs assistance in transfer	No cognitive impairment and calls nurses when necessary	No urinary problems	No tube or drain inserted	Antibiotic	Less than 65 years old
18	Needs assistance in transfer	Impaired cognition and does not call nurses when necessary	Nocturia	Multiple tubes and/or drains inserted	Anti-hypertensive	Less than 65 years old

Supplementary Table 2. Factor analysis of the cognitive scales and reliability coefficients

Variables	Factor 1	Factor 2	Factor 3
BN item1	0.665		
BN item2	0.740		
BN item3	0.593		
AB item1		0.670	
AB item2		0.659	
AB item3		0.622	
BB item1			0.651
BB item2			0.697
BB item3			0.436
Eigenvalue	2.169	2.030	1.587
Variance explained (Cumulative variance)	0.241	0.226 (0.467)	0.176 (0.643)
KR 20 coefficients	0.323	0.562	0.434

Note: BN: Base-rate neglect, AB: Availability bias, BB: Belief bias

Bartlett's test of sphericity: Chi-squared (36) = 804.64, $p < .001$, The Kaiser, Meyer, Olkin (KMO) = 0.59.

Supplementary Table 3. Frequency of the responses in cognitive bias

Total score	Base-rate neglect	Belief bias	Availability bias
0	4 (1.19%)	28 (8.36%)	98 (29.25%)
1	16 (4.78%)	68 (20.30%)	93 (27.76%)
2	83 (24.78%)	113 (33.73%)	88 (26.27%)
3	230 (68.66%)	119 (35.52%)	54 (16.12%)
Missing	2 (0.60%)	7 (2.09%)	2 (0.60%)
Mean (SD)	2.62 (0.64)	1.99 (0.95)	1.29 (1.06)

Note: Means and the standard deviations were calculated after replacing missing values with the median values.

Supplementary Table 4. Spearman correlation coefficients between variables

	Experience	Age	Education	Sex	Base-rate neglect	Belief bias	Availability bias
Age	0.921*						
Education	-0.291*	-0.271*					
Sex	-0.004	0.028	0.073				
Base-rate neglect	-0.042	-0.074	-0.142*	-0.090			
Belief bias	0.100	0.061	-0.060	0.025	0.056		
Availability bias	0.098	0.113*	0.025	0.105	-0.050	0.072	
Ns Judgement (Mean)	-0.052	-0.079	-0.138*	-0.150*	0.099	-0.029	-0.166*

Note: * $p < 0.05$. Missing values were replaced with the median values in the continuous/ordinal variables, and the most representative category in the nominal variables.

The dummy variables are as follows: Education (0: Diploma, 1: Degree), Sex (0: Women, 1: Men). The scores of the bias scales ranged from 0 to 3.