

Supplementary Data for “Peritraumatic C-reactive protein levels predict pain outcomes following traumatic stress exposure in a sex-dependent manner”

Supplementary Table 1. Full model statistics for the relationship between peritraumatic (i.e. ED) CRP levels and longitudinal pain outcomes (week 8, month 3, month 6) in men and women

	Men (n=99)				Women (n=220)			
	β	SEM	t-value	<i>p</i>	β	SEM	t-value	<i>p</i>
Time	-0.36	0.08	-4.71	<0.001	-0.16	0.05	-3.39	<0.001
ED CRP	-0.24	0.11	-2.13	0.037	0.05	0.07	0.72	0.470
Age	0.03	0.02	1.50	0.137	0.02	0.01	1.44	0.150
ED pain	0.38	0.10	3.70	<0.001	0.38	0.08	4.88	<0.001
Highest grade: high school	-0.56	0.91	-0.62	0.539	-0.19	0.73	-0.26	0.799
Highest grade: some college	0.15	0.83	0.18	0.855	0.21	0.66	0.31	0.756
Highest grade: \geq bachelor's degree	0.17	1.04	0.16	0.872	-0.09	0.74	-0.12	0.904
Race	0.39	0.59	0.65	0.516	0.11	0.44	0.24	0.812
Plate 2	-0.55	1.10	-0.50	0.620	1.46	0.79	1.85	0.066
Plate 3	0.79	1.18	0.67	0.502	1.68	0.79	2.13	0.034
Plate 4	0.19	1.50	0.13	0.900	1.58	0.85	1.86	0.064
Plate 5	-0.57	1.71	-0.33	0.739	0.89	0.94	0.95	0.344
Plate 6	-0.09	1.83	-0.05	0.959	1.10	0.98	1.13	0.262
Plate 7	-0.81	1.40	-0.58	0.565	1.62	0.91	1.78	0.076
Plate 8	1.52	1.61	0.94	0.349	2.37	0.92	2.58	0.011
Plate 9	0.44	1.16	0.38	0.703	2.09	0.74	2.82	0.005
Plate 10	-2.49	1.46	-1.71	0.092	1.73	0.85	2.05	0.042
Plate 11	3.15	2.06	1.53	0.130	0.81	1.21	0.67	0.503
Plate 12	0.72	1.38	0.52	0.607	1.31	0.86	1.53	0.129
Plate 13	-2.28	1.79	-1.27	0.208	2.24	1.12	2.00	0.047

ED: emergency department, CRP: C-reactive protein, SEM: standard error of the mean
CRP was log₂-adjusted

Supplementary Table 2. Full model statistics for the relationship between peritraumatic (i.e. ED) CRP levels and longitudinal pain outcomes (week 8, month 3, month 6) adjusted for BMI, in men and women

	Men (n=78)				Women (n=197)			
	β	SEM	t-value	<i>p</i>	β	SEM	t-value	<i>p</i>
Time	-0.35	0.08	-4.32	<0.001	-0.14	0.05	-2.77	0.006
CRP	-0.23	0.14	-1.63	0.110	0.04	0.09	0.46	0.643
Age	0.01	0.02	0.36	0.720	0.02	0.02	1.29	0.199
ED pain	0.24	0.12	1.93	0.059	0.36	0.08	4.38	<0.001
Highest grade: high school	-0.54	1.12	-0.48	0.630	0.14	0.76	0.19	0.850
Highest grade: some college	0.13	1.00	0.13	0.899	0.17	0.69	0.24	0.808
Highest grade: \geq bachelor's degree	-0.42	1.27	-0.33	0.743	0.06	0.76	0.08	0.933
Race	0.29	0.70	0.42	0.679	0.29	0.47	0.60	0.547
BMI	0.00	0.05	-0.02	0.987	0.03	0.03	0.86	0.392
Plate 2	0.74	1.50	0.49	0.623	1.61	0.87	1.85	0.066
Plate 3	0.54	1.75	0.31	0.759	2.19	0.87	2.52	0.013
Plate 4	2.87	1.98	1.45	0.152	1.76	0.94	1.86	0.064
Plate 5	0.07	1.95	0.04	0.970	0.90	0.99	0.91	0.365
Plate 6	0.91	2.45	0.37	0.713	1.30	1.09	1.19	0.236
Plate 7	-0.61	1.87	-0.33	0.745	1.78	1.01	1.76	0.080
Plate 8	2.45	1.89	1.30	0.201	2.44	0.96	2.54	0.012
Plate 9	1.80	1.54	1.17	0.248	1.93	0.78	2.48	0.014
Plate 10	-1.21	1.78	-0.68	0.499	1.62	0.88	1.83	0.068
Plate 11	4.27	2.31	1.85	0.070	0.68	1.23	0.55	0.584
Plate 12	2.05	1.71	1.20	0.235	1.28	0.90	1.43	0.156
Plate 13	-1.27	2.02	-0.63	0.533	2.15	1.15	1.86	0.065

ED:emergency department, CRP: C-reactive protein, BMI: body mass index, SEM: standard error of the mean
CRP was log₂-adjusted

Supplementary Table 3. Full model statistics for the relationship between peritraumatic (i.e. ED) CRP levels and longitudinal pain outcomes (week 8, month 3, month 6) in men and women, the subcohort of participants with available BMI data (BMI not included as a covariate)

	Men (n=78)				Women (n=197)			
	β	SEM	t-value	p	β	SEM	t-value	p
Time	-0.35	0.08	-4.32	<0.001	-0.14	0.05	-2.77	0.006
CRP	-0.23	0.14	-1.69	0.096	0.07	0.08	0.94	0.346
Age	0.01	0.02	0.37	0.716	0.02	0.01	1.42	0.159
ED pain	0.24	0.12	1.95	0.056	0.37	0.08	4.58	<0.001
Highest grade: high school	-0.54	1.11	-0.49	0.627	0.15	0.76	0.20	0.842
Highest grade: some college	0.13	0.99	0.13	0.898	0.23	0.69	0.33	0.743
Highest grade: \geq bachelor's degree	-0.42	1.23	-0.34	0.732	0.06	0.76	0.08	0.935
Race	0.29	0.68	0.42	0.674	0.30	0.47	0.63	0.526
Plate 2	0.75	1.46	0.51	0.611	1.52	0.86	1.76	0.080
Plate 3	0.55	1.66	0.33	0.742	2.17	0.87	2.50	0.014
Plate 4	2.87	1.96	1.47	0.148	1.70	0.94	1.81	0.072
Plate 5	0.08	1.88	0.04	0.966	0.83	0.99	0.83	0.405
Plate 6	0.91	2.41	0.38	0.706	1.23	1.09	1.14	0.258
Plate 7	-0.61	1.86	-0.33	0.742	1.71	1.01	1.70	0.091
Plate 8	2.46	1.87	1.32	0.194	2.31	0.95	2.44	0.016
Plate 9	1.81	1.50	1.20	0.234	1.95	0.78	2.50	0.013
Plate 10	-1.21	1.74	-0.70	0.490	1.64	0.88	1.86	0.065
Plate 11	4.27	2.29	1.87	0.067	0.68	1.23	0.55	0.581
Plate 12	2.06	1.67	1.23	0.224	1.25	0.90	1.40	0.165
Plate 13	-1.27	2.00	-0.63	0.529	2.18	1.15	1.89	0.061

ED: emergency department, CRP: C-reactive protein, BMI: body mass index, SEM: standard error of the mean
CRP was log₂-adjusted

Supplementary Table 4. Counts of participants with increasing versus decreasing CRP and Pain trends.

		CRP Trend from ED to 6M			
		Decreasing CRP		Increasing CRP	
		Men	Women	Men	Women
Pain Trend from ED to 6M	Decreasing Pain	n= 39	n=72	n=27	n=62
	Increasing Pain	n=6	n=28	n=1	n=26
	Unchanged	n=2	n=10	n=4	n=14

Men: χ^2 -test $p=0.115$

Women: χ^2 -test $p=0.552$

CRP: C-reactive protein, ED: emergency department, 6M: six months