

Supplementary to: Substituting red and processed meat, poultry, and fish with legumes and risk of non-alcoholic fatty liver disease in a large cohort

Fie Langmann¹, Daniel B. Ibsen^{1,2}, Luke W. Johnston², Aurora Perez-Cornago³, Christina C. Dahm¹

1. Aarhus University, Department of Public Health, Bartholins Allé 2, 8000 Aarhus C, Denmark.

2. Steno Diabetes Center Aarhus, Aarhus University Hospital, Palle Juul-Jensens Boulevard 11, 8200 Aarhus C, Denmark

3. European Commission, Joint Research Centre (JRC), Ispra, Italy

Contents

Supplementary Table 1.....	3
Supplementary Table 2.....	5
Supplementary Figure 1	8

Supplementary Table 1. Description of dietary components included in the definition of food groups for the study.

Food group	Dietary components
Legumes	Legumes and pulses, soy desserts and yoghurt, soy milk, meat substitutes made from soy.
Red and processed meat	Beef, pork, lamb, offal, sausages, bacon, ham, and liver pâté.
Poultry	Poultry with or without skin and fried poultry with batter or breadcrumbs.
Fish	Oily fish, white fish, tinned tuna, fried fish with batter or breadcrumbs, and shellfish.
Eggs and egg dishes	Eggs and egg-based dishes like scrambled eggs, omelets, and quiche.
Refined cereal	White pasta and rice, white bread, biscuit cereal, biscuits, savory crackers, mixed bread (50/50 brown and seeded), other bread, and other cereal with high sugar content.
Whole grain cereal	Whole meal bread, whole meal pasta, brown rice, bran cereal, oat cereal (with and without added sugar), muesli, and other whole grains.
Dairy	Milk (whole, semi-skimmed, skimmed), cream, cheese (low, medium, and high fat), yoghurt (low or full fat), milk-based and powdered drinks, and milk-dairy desserts.
Fruits	Berries, citrus fruits, apples, pears, dried fruits, stewed fruits, and other fruits.
Vegetables	Green leafy vegetables, cabbages, onion, garlic, root vegetables, raw salad, peas, sweetcorn, vegetable dips (guacamole and hummus), tomatoes, mushrooms, mixed vegetables, and vegetable side dishes.
Potatoes	Fried/roasted potatoes, mashed potatoes, and baked or boiled potatoes and sweet potatoes.
Nuts	Salted and unsalted nuts and seeds.
Fats	Olive oil, animal-based fat spreads, and plant-based fat spreads.

Mixed dishes	Pizza, grain dishes with added fat, samosa, pakora, soups, sushi, and vegetarian non-soy meat substitutes like quorn.
Sauce and condiments	Sauces and condiments with low or high fat content.
Snacks and sweets	Nut-based spreads, savoury snacks, chocolate confectionery, added sugars and preserves, cakes, pastry, other desserts, and other sweets.
Non-alcoholic beverages	Coffee, tea, rice milk, oat milk, water (still or sparkling), sugar-sweetened soft drinks, artificially sweetened soft drinks, and fruit juices.
Alcoholic beverages	Beer, cider, spirits, and fortified and non-fortified wines.

Supplementary Table 2. Baseline characteristics of participants in the UK Biobank cohort across incident NAFLD (N = 123,586)

Characteristic	All participants N = 124,194	With NAFLD^a N = 1201
Age	57.0 (44.0, 66.0)	57.0 (45.6, 66.0)
Sex		
Female	69,273 (56%)	618 (51%)
Male	54,921 (44%)	583 (49%)
Yearly income, £		
>100,000	8,904 (7.2%)	54 (4.5%)
52,000-100,000	28,831 (23%)	209 (17%)
31,000-51,999	32,819 (26%)	289 (24%)
18,000-30,999	26,854 (22%)	292 (24%)
<18,000	15,602 (13%)	227 (19%)
Unknown	11,184 (9.0%)	130 (11%)
Educational level^b		
High	58,435 (47%)	423 (35%)
Intermediate	41,127 (33%)	442 (37%)
Low	24,632 (20%)	336 (28%)
Deprivation score^c	-2.4 (-4.7, 2.6)	-1.7 (-4.5, 3.6)
Cohabitation		
Alone	22,134 (18%)	241 (20%)
With spouse/partner	92,515 (74%)	833 (69%)
Other non-partner	9,366 (7.5%)	121 (10%)
Unknown	179 (0.1%)	6 (0.5%)
Ethnicity		
White	119,968 (97%)	1,147 (96%)
Other	4,226 (3.4%)	54 (4.5%)
Physical activity^d		
High	21,596 (17%)	176 (15%)
Moderate	55,037 (44%)	469 (39%)
Low	30,167 (24%)	370 (31%)
Unknown	17,394 (14%)	186 (15%)

Smoking status

Current, >15 cigarettes/day	1,797 (1.4%)	33 (2.7%)
Current, <15 cigarettes/day	3,379 (2.7%)	45 (3.7%)
Former	44,379 (36%)	509 (42%)
Never	70,978 (57%)	570 (47%)
Unknown	3,661 (2.9%)	44 (3.7%)

Alcohol consumption, g/day 11.4 (0.0, 44.0) 9.0 (0.0, 51.4)

UK-region of recruitment

East Midlands	9,931 (8.0%)	101 (8.4%)
London	25,432 (20%)	282 (23%)
North East	11,794 (9.5%)	156 (13%)
North West	15,291 (12%)	205 (17%)
Scotland	6,577 (5.3%)	15 (1.2%)
South East	11,656 (9.4%)	72 (6.0%)
South West	12,917 (10%)	122 (10%)
Wales	3,876 (3.1%)	11 (0.9%)
West Midlands	7,542 (6.1%)	66 (5.5%)
Yorkshire and Humber	19,178 (15%)	171 (14%)

BMI \geq 30 kg/m² 24,556 (20%) 592 (49%)

History of NAFLD-related diseases^e 45,747 (37%) 715 (60%)

Family history of related diseases^f 96,936 (78%) 982 (82%)


Food group intakes, g/week^g

Legumes	0.0 (0.0, 472.5)	0.0 (0.0, 472.5)
Red and processed meat	371.0 (0.0, 843.5)	420.0 (0.0, 1,022.0)
Poultry	175.0 (0.0, 606.7)	227.5 (0.0, 606.7)
Fish	175.0 (0.0, 595.0)	161.0 (0.0, 581.0)
Eggs and egg dishes	0.0 (0.0, 420.0)	0.0 (0.0, 513.3)
Refined cereal	850.5 (257.3, 1,674.6)	855.8 (269.5, 1,659.0)
Whole grain cereal	474.8 (0.0, 1,424.5)	350.0 (0.0, 1,361.5)
Dairy	1,942.5 (717.5, 3,456.3)	1,837.5 (600.8, 3,517.5)
Fruits	1,341.2 (280.0, 2,905.0)	1,190.0 (70.0, 2,968.0)

Vegetables	1,190.0 (330.8, 2,570.6)	1,114.8 (236.3, 2,637.3)
Potatoes	625.6 (0.0, 1,260.0)	630.0 (0.0, 1,365.0)
Nuts	7.0 (0.0, 150.5)	0.0 (0.0, 140.0)
Fats	79.6 (7.6, 180.9)	86.9 (8.4, 198.6)
Mixed dishes	210.0 (0.0, 1,166.7)	210.0 (0.0, 1,166.7)
Sauce and condiments	116.7 (0.0, 385.0)	105.0 (0.0, 385.0)
Snacks and sweets	518.0 (112.0, 1,194.7)	556.5 (116.7, 1,354.5)
Non-alcoholic beverages	10,733.3 (7,157.5, 15,295.0)	10,990.0 (7,093.3, 15,890.0)
Alcoholic beverages	938.0 (0.0, 4,676.0)	721.9 (0.0, 5,496.8)
Total weight of food and beverages	22,023.0 (16,611.0, 28,878.6)	22,325.6 (16,421.8, 29,921.8)

Continuous variables are presented as median (10%, 90%) and categorical values are presented as numbers (%).

^aNAFLD, non-alcoholic fatty liver disease. ^bEducational level was defined as low (Certificate of Secondary Education (CSE), National Vocational Qualifications, Higher National Diploma, Higher National Certificates, other professional qualifications, or equivalent), intermediate (A levels, O levels, General Certificate of Secondary Education, or equivalent), and high (College or University degree). ^cDeprivation was assessed with the Townsend Deprivation Index based on four indicators of material deprivation: non-home ownership, non-car ownership, unemployment, and overcrowding. Positive values indicate that individuals live in areas with high material deprivation and negative values indicate relative affluence.⁽⁸⁴⁾ ^dPhysical activity was based on total metabolic equivalent task (MET) minutes per week for all activity including walking, moderate, and vigorous activity, and defined as low (0-9.9 METs/week), moderate (10-49.9 METs/week), high (≥ 50 METs/week), and unknown. ^eHistory of NAFLD-related diseases was defined as participant's previous diagnosis of diabetes, alcoholic liver disease, angina pectoris, hypertension, myocardial infarction, stroke, elevated cholesterol, or gallbladder disease), ^fFamily history of related diseases was defined as diagnosis of diabetes, heart disease, stroke, hypertension, or elevated cholesterol among the mother, father, and/or biological sibling(s) of the participant. ^gThe definition and inclusion of foods in each food group can be found in Supplementary Table 1.



Supplementary Figure 1. Directed Acyclic Graph representing the association between replacing meat, poultry, or fish with legumes and risk of non-alcoholic fatty liver disease (NAFLD) and the assumed relationship with covariates. Socio-economic status comprises factors like ethnicity, Townsend Deprivation Index, geographical living area, cohabitation status, educational level, and yearly income. Related diseases include cancer, hypertension, stroke, myocardial infarction, elevated cholesterol, gallbladder diseases, alcoholic liver disease, diabetes or family history of hypertension, diabetes, stroke, or heart disease. ● exposure, ● outcome, ● ancestor of outcome, ● ancestor of exposure and outcome, — causal path, — biasing path