



(B)

P-Value Matrix with Statistical Significance Highlighted (p < 0.05)													
CCL5		2.7e-64	8.4e-27	2.9e-39	0.032	1.4e-18	3.1e-67	3.9e-41	2.7e-77	1.8e-62	3.4e-46	3.9e-07	2e-12
T_cells	2.7e-64		1.5e-25	1.8e-45		3.3e-18	3.4e-45	5.3e-47	3.3e-53	2.8e-38	2.5e-27	1e-05	1e-09
B_cells	- 8.4e-27	1.5e-25		5.2e-21	0.0074	5.7e-07	5.8e-35	2e-24	3.9e-30	1.1e-21	1.8e-15	2.6e-08	0.0003
NK_cells	- 2.9e-39	1.8e-45	5.2e-21			3.3e-13	1.2e-35	1.9e-27	7.4e-40	1.5e-35	3.7e-27	2.1e-06	1.1e-06
Granulocyte	0.032		0.0074			0.00068		2.5e-06	0.0074				0.0067
Macrophage	1.4e-18	3.3e-18	5.7e-07	3.3e-13	0.00068		6.1e-15	4.7e-18	9e-18	9.8e-19	1e-14	0.0014	4.6e-12
Th1_cells	- 3.1e-67	3.4e-45	5.8e-35	1.2e-35		6.1e-15		4.2e-33	1.8e-56	5.5e-65	4.5e-45	6.8e-09	2.5e-07
Th2_cells	3.9e-41	5.3e-47	2e-24	1.9e-27	2.5e-06	4.7e-18	4.2e-33		1.3e-38	1.4e-28	2.4e-17	2.8e-07	1.6e-12
CD8_T_cells	- 2.7e-77	3.3e-53	3.9e-30	7.4e-40	0.0074	9e-18	1.8e-56	1.3e-38		1.7e-57	7.8e-40	3.3e-08	5.3e-08
PRF1	- 1.8e-62	2.8e-38	1.1e-21	1.5e-35		9.8e-19	5.5e-65	1.4e-28	1.7e-57		1.4e-60	9.4e-07	7.5e-07
GZMB	3.4e-46	2.5e-27	1.8e-15	3.7e-27		le-14	4.5e-45	2.4e-17	7.8e-40	1.4e-60		9.3e-05	1.2e-05
Tregs	- 3.9e-07	1e-05	2.6e-08	2.1e-06		0.0014	6.8e-09	2.8e-07	3.3e-08	9.4e-07	9.3e-05		0.0074
MDSCs	- 2e-12	1e-09	0.0003	1.1e-06	0.0067	4.6e-12	2.5e-07	1.6e-12	5.3e-08	7.5e-07	1.2e-05	0.0074	0
	CCL5	T_cells	B_cells -	NK_cells	Granulocyte -	Macrophage -	Th1_cells -	Th2_cells -	CD8_T_cells -	PRF1 -	- GZMB	Tregs -	MDSCs -

Figure S2. The non-parametric correlation analysis between CCL5 expression and various immune cells.

(A) Non-parametric Spearman rho between CCL5 and the selected immune genes are calculated and analyzed to determine the strength and direction of their linear relationships. (B) The correlation of the *p*-value matrix with incorporated in the figure.