

Severe *GBA1* variants drive the GBA-PD clinical phenotype: implications for counselling and clinical trials

Elisa Menozzi, MD, PhD^{1,2}, Sara Lucas Del Pozo, MD^{1,2}, Jane Macnaughtan, MD, PhD^{2,3}, Roxana Meزابrovski^{1,2}, Sofia Koletsi^{1,2}, Pierfrancesco Mitrotti, MD^{4,5}, Luca Gallo, MD^{4,5}, Rosaria Calabrese, PsyD^{4,5}, Marco Toffoli, MD, PhD^{1,2}, Nadine Loefflad^{1,2}, Franco Valzania, MD⁶, Francesco Cavallieri, MD, PhD⁶, Valentina Fioravanti, MD, PhD⁶, Selen Yalkic^{1,2}, Naomi Limbachiya^{1,2}, Fabio Blandini, MD, PhD^{4,8§}, Micol Avenali, MD, PhD^{4,5§}, Anthony HV Schapira, MD, DSc, FRCP, FMedSci^{1,2§}

¹Department of Clinical and Movement Neurosciences, Queen Square Institute of Neurology, University College London (UCL), London, United Kingdom

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815

³Liver Failure Group, Institute for Liver and Digestive Health, University College London, London, UK

⁴Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy

⁵IRCCS Mondino Foundation, Pavia, Italy

⁶Neurology Unit, Neuromotor and Rehabilitation Department, Azienda USL-IRCCS di Reggio Emilia, Reggio Emilia, Italy

⁸Ca' Granda IRCCS Foundation, Ospedale Maggiore Policlinico, Milan, Italy

§senior authors

Corresponding author:

Anthony HV Schapira

Address: Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, London NW3 2PF, UK

Email: a.schapira@ucl.ac.uk

Telephone: +44 2078302012

Word count: 1,700 words

Running title: Severe *GBA1* variants driving GBA-PD phenotype

Abstract (147 words)

Background

Variants in the *GBA1* gene are the commonest genetic risk factor for Parkinson disease (PD). Genotype-phenotype correlations exist but with conflicting data, particularly in the cognitive domain.

Objectives

Comparing clinical phenotypes in a multicentre, international cohort incorporating GBA-PD and idiopathic PD (iPD) patients.

Methods

Patients underwent a comprehensive assessment of motor and non-motor functions. Two-group (GBA-PD vs iPD) and multiple-group comparisons (iPD, risk, mild, and severe variant GBA-PD) were performed.

Results

Three hundred fifteen PD patients were recruited: 186 iPD, 39 severe GBA-PD, 24 mild GBA-PD, 56 risk GBA-PD, and 10 patients carrying variants of unknown significance. Groups were matched for sex, disease duration and medications. Mild and severe GBA-PD were younger and developed PD earlier. Severe GBA-PD had worse depression, cognitive impairment and hyposmia, and a trend for higher rates of motor complications.

Conclusions

Only severe variant GBA-PD have a distinctive, more severe clinical profile.

Main text (1,700 words)

Introduction

Variants in the *GBA1* gene are highly prevalent in the Parkinson disease (PD) population (~10-15%).¹ In recent years, the increased recourse to *GBA1* genotyping in PD,² the different response of patients carrying *GBA1* variants (GBA-PD) to advanced treatments for PD (i.e., deep brain stimulation-DBS),³ and the growing number of clinical trials testing *GBA1*-targeted therapies,⁴ have created the need for standardized guidelines for genetic counselling, management and selection of patients for clinical trials.¹

Major challenges in manifest GBA-PD include the differences in clinical severity and progression according to *GBA1* variant type.⁴ Several case-control studies have reported that PD patients carrying severe *GBA1* variants (e.g., p.L483P) compared to carriers of mild (e.g., p.N409S) or risk (e.g., p.E365K) variants, present with a more severe clinical phenotype characterised by faster progression, and worse psychiatric and cognitive dysfunction, worsened by interventions such as DBS.^{3,5-8} However, recent studies have contradicted these results, reporting similar cognitive and motor deterioration in *GBA1* variant carriers, regardless of variant type.⁹⁻¹² Phase II trials are now underway evaluating changes in cognitive function at 12 months as primary outcomes, recruiting either mild or severe variant GBA-PD.¹³ An understanding of the natural clinical history of PD associated with different *GBA1* variants is crucial to appropriate trial design.

Here, we present our experience from a large, multi-centre, international case-control study incorporating PD patients negative (idiopathic PD-iPD) or positive for *GBA1* variants (GBA-PD), the latter group being representative of all variant types (risk, mild, and severe). We highlight how the severe *GBA1* variant carriers represent the only group with a remarkably different and more severe clinical phenotype compared to iPD. Our findings caution the PD community to avoid generalisation and consider GBA-PD as a clinically heterogeneous condition.

Methods

Study design and clinical data

Participants were recruited via the RAPSODI study (University College London) in the United Kingdom,¹⁴ and two neurology tertiary centres in Italy (IRCCS Mondino Foundation, Pavia, and Azienda USL-IRCCS, Reggio Emilia). *GBA1* variants were classified into four classes (severe, mild, risk, and unknown significance) as previously reported (a list of *GBA1* variants included in the study is presented in Supplementary Table 1).¹⁵ Information about family history, disease onset, and medications was collected, and levodopa equivalent daily dose (LEDD) calculated for each participant.^{16, 17} Motor and non-motor functions were evaluated through an extensive clinical examination (see Supplementary Methods).

This study was approved by the local Ethics Committees (London – Queen Square REC: 15/LO/1155; EC of Pavia: code P-20210009687; EC of Area Vasta Emilia Nord: code 2021/0092531). All participants signed informed consent upon enrolment.

Statistical analysis

A first set of statistical analyses were completed comparing iPD to GBA-PD. In a second set, patients were stratified according to their *GBA1* variant severity into risk, mild, and severe (carriers of variants of unknown significance were excluded from this comparison because of the low numbers of individuals in this group). Study data were collected and managed using REDCap electronic data capture tools.¹⁸ Statistical analyses were performed using R version 4.2.3.¹⁹ Full statistical analysis methods can be found in Supplementary Methods.

Results

Demographic and clinical features of iPD and GBA-PD groups are summarised in Table 1. A total of 315 PD patients (186 iPD, 129 GBA-PD) were recruited. Within the GBA-PD, 43% (N=56) carried *GBA1* risk variants, 30% (N=39) severe variants, 19% (N=24) mild variants, and 8% (N=10) carried variants of unknown significance.

Two-group comparison: iPD vs GBA-PD

Compared with iPD, GBA-PD patients showed a significantly younger age (61 vs 65 years) and more frequent positive family history for PD (36.4% vs 23% of participants). No differences in sex distribution were found between groups. In terms of disease-associated features, GBA-PD patients developed the disease 5 years earlier and had undergone DBS more frequently than iPD patients. No differences in disease duration or LEDD were identified.

GBA-PD patients presented with more severe mood disorders (HADS depression: OR, 2.1; 95% CI: 1.2-3.4; $P=0.004$, and BDI: OR, 2.4; 95% CI: 1.6-3.8; $p<0.0001$), anxiety (HADS anxiety: OR, 1.8; 95% CI: 1.1-2.9; $p=0.02$), olfactory dysfunction (UPSIT: $\beta=-2.2$, $p=0.002$), subjective motor disability (MDS-UPDRS part II: $p=0.02$) and motor complications (MDS-UPDRS part IV: $p=0.002$). Despite similar global cognitive function as measured by the total MOCA score, the GBA-PD group performed significantly worse in tasks evaluating visuospatial and executive functions (all p values <0.05 , see Table 2; individual p values, ORs and CIs are reported in Supplementary Table 2).

Clinical features in stratified *GBA1*-variant type groups

When patients were stratified according to *GBA1* variant type, severe and mild GBA-PD were significantly younger than both risk GBA-PD ($p=0.03$, and $p=0.004$, respectively) and iPD (both $p<0.001$), with no difference in age between severe and mild GBA-PD. Severe and mild GBA-PD also developed PD earlier than risk GBA-PD ($p=0.005$ and $p<0.001$, respectively) and iPD (both $p<0.001$). Severe GBA-PD patients had a more frequent positive family history for PD (46%) when compared to iPD patients and were more frequently subjected to DBS procedures (Table 1). Groups were homogenous in terms of disease duration and LEDD.

When clinical features were compared, severe GBA-PD patients presented with a distinct clinical profile when compared to iPD patients, but not to the risk or mild GBA-PD patients (Table 1). This profile was characterised by more severe depression (HADS depression: OR, 2.7; 95% CI: 1.3-5.5; $P=0.005$, and BDI: OR, 3.8; 95% CI: 2-7.5; $p<0.0001$), olfactory dysfunction (UPSIT: $\beta=-3.8$, $p=0.0007$) and cognitive impairment (MOCA: $\beta=-1.9$, $p=0.001$), and a trend for more severe motor complications and constipation (see Table 1). Cognitive performances were worse in severe GBA-PD, also compared with risk and mild GBA-PD patients (all p values <0.01); these results remained statistically significant even after removing subjects with DBS, only with a trend observed for the comparison between mild and severe carriers ($p=0.057$), which might be due to low numbers of individuals (20 and 26, respectively) and thus loss of statistical power. Severe GBA-PD were mostly affected in visuospatial and executive function and attention abilities (Table 2 and Supplementary Table 2). Interestingly, we detected a trend for higher scores in the SCOPA-pupillomotor subdomain in severe GBA-PD when compared to both mild and risk GBA-PD (see Table 1). No other statistically significant differences emerged among the four groups in the severity of motor and non-motor symptoms when results were adjusted for multiple comparisons.

Discussion

In this large, multi-centre, international case-control study, we demonstrated the importance of considering the type of *GBA1* genetic variant in research studies and clinical trials conducted on GBA-PD. Reporting findings on GBA-PD without stratifying according to *GBA1* variant type, can be misleading: differences can be detected related to iPD, however these are mainly driven by severe variant carriers.

Our results align with previous findings showing a more severe phenotype in severe variant carriers.^{6, 8, 20} In a previous study comparing a large cohort of GBA-PD (139 mild, 48 severe) to 152 iPD patients, more severe depression, hallucinations, worse hyposmia, and higher frequencies of REM sleep behaviour disorder (RBD) were detected in severe GBA-PD compared to the other two groups.²⁰ However, this study did not include risk variant carriers, who numerically represent the predominant group of GBA-PD individuals, especially in those of Caucasian background.¹

In contrast to previous reports, we did not detect any cognitive impairment in carriers of risk or mild variants. One study found that both pathogenic (mix of mild and severe) variant carriers (N=60) and p.E365K carriers (N=65) had higher incidence of dementia and a greater impairment in working memory/executive function and visuospatial abilities compared to iPD, however the disease duration in these groups was mostly >7.5years, so additional factors beyond *GBA1* status might have contributed to these results.¹¹ Cognitive decline at 7 years from diagnosis was faster in pathogenic and risk variant carriers than iPD in another study, however, the association with progression rate to dementia was much smaller in the risk variant carriers group, and no distinction was made between severe and mild variant carriers.²¹

Our results showed a trend toward more severe cardiovascular and thermoregulatory dysfunction in GBA-PD vs iPD, but we did not see any differences across variant types. Previous reports also found more impaired cardiovascular autonomic control characterised by a lower parasympathetic modulation at rest and a lower parasympathetic modulation in response to active standing, in GBA-PD compared to iPD.²² GBA-PD also presented a lower heart-to-mediastinum uptake ratio in ¹²³I-meta-iodobenzylguanidine scintigraphy suggestive of reduced cardiac sympathetic denervation,²³ and greater cardiac sympatho-vagal demodulation.²⁴ The latter study detected greater impairment in cardiovagal function in severe GBA-PD compared to risk and mild GBA-PD combined together, but numbers were small.²⁴ Nonetheless, these findings make this area worth investigating in the future. In our study, we showed a trend for more severe impairment in pupillomotor function in severe variant carriers compared to mild or risk, which might be related to a pupillary parasympathetic dysfunction which might be more extreme in severe GBA-PD. Interestingly, pupillomotor abnormalities suggestive of dysfunctional parasympathetic innervation were also detected in individuals with neuronopathic Gaucher disease, caused by biallelic pathogenic variants in *GBA1*,²⁵ or with Fabry disease, another lysosomal storage disorder.²⁶

Several disease-modifying therapies targeting the *GBA1* pathway are in Phase II/III of clinical trial stage, aiming to halt/slow GBA-PD progression.^{4, 27} Recently, a Phase II, randomized-controlled trial (MOVES-PD), testing the effect of the glucosylceramide synthase inhibitor Venglustat in GBA-PD patients, did not meet its primary endpoint, and showed paradoxical worsening of motor symptoms in mild GBA-PD treated with Venglustat.²⁸ This highlights the need to refine patients' selection and stratify individuals for variant severity in trials targeting the GBA-PD population in order to optimise chances of success.²⁸

Our study has some limitations. First, the relatively small number of individuals with severe and mild GBA-PD might have undermined the results. Second, this is a case-control study, however a 2-year follow-up assessment of this cohort is in progress.

Despite these limitations, we believe that this study provides clinicians with useful information to approach their GBA-PD patients in the clinic, tailoring their counselling on prognosis and disease trajectories, and integrating our recent consensus guidance on PD risk.¹ Moreover, we hope that these findings will inform researchers involved in clinical trials to improve the design of such studies and guide the choice of realistic outcomes in the correct patient groups. Finally, this study provides insight into the pathophysiological mechanisms underlying disease severity in GBA-PD.

Conflict of Interest Statement:

EM, SLDP, JM, SK, PM, LG, RC, MT, NL, FV, FC, VF, NL, FB and MA have no conflicts to disclose. RM and SY are supported by a Royal Free Charity fellowship.

AHVS has provided paid consultancy to Capsida, Neurocrine and Auxilius, is the Chief Investigator of the ambroxol phase III study and a Principal Investigator of the MOVES-PD study.

Acknowledgments:

This research was funded in part by Aligning Science Across Parkinson's (Grant number: ASAP-000420) through the Michael J. Fox Foundation for Parkinson's Research (MJFF) and by the EU Joint Programme—Neurodegenerative Research (JPND: GBA-PaCTS) through the MRC grant code MR/T046007/1. For the purpose of open access, the author has applied a CC BY 4.0 public copyright license to all Author Accepted Manuscripts arising from this submission.

Authors' Roles

1. Research project: A. Conception, B. Organization, C. Execution.
2. Statistical Analysis: A. Design, B. Execution, C. Review and Critique.
3. Manuscript Preparation: A. Writing of the first draft, B. Review and Critique.

EM: 1A, 1B, 1C, 2A, 2B, 2C, 3A.

SLDP: 1A, 1B, 1C, 3B.

JM: 1A, 1B, 1C, 3B.

RM: 1B, 1C, 3B.

SK: 1B, 3B.

PM: 1B, 1C, 3B.

LG: 1B, 1C, 3B.

RC: 1B, 1C, 3B.

MT: 1B, 1C, 3B.

NL: 1B, 1C, 3B.

FV: 1C, 3B.

FC: 1C, 3B.

VF: 1C, 3B.

SY: 1B, 1C, 3B.

NL: 1B, 1C, 3B.

FB: 1A, 1B, 2C, 3B.

MA: 1A, 1B, 1C, 2C, 3B.

AHVS: 1A, 1B, 2C, 3B.

Table 1. Demographics and clinical features of study cohort.

	Negative - iPD (N=186)	GBA-PD (N=129)	p value	Risk (N=56)	Mild (N=24)	Severe (N=39)	P1 (N vs R)	P2 (N vs M)	P3 (N vs S)	P4 (R vs M)	P5 (R vs S)	P6 (M vs S)
GBR (N)	93	63		35	13	12						
ITA (N)	93	66		21	11	27						
Age	65 ± 9	61 ± 9	***	63 ± 8	56 ± 10	59 ± 8	ns	***	***	**	*	ns
Sex (% F, N)	42% (78)	47% (61)	ns	37.5% (21)	54% (13)	54% (21)	ns	ns	ns	ns	ns	ns
PD family history (% N)	23% (43)	36.4% (47)	*	34% (19)	25% (6)	46% (18)	ns	ns	*	ns	ns	ns
Education (yrs)	13.9 ± 4.4	14.2 ± 4.1	*	14.3 ± 4.6	14.2 ± 3.8	14.2 ± 4.1	ns	ns	ns	ns	ns	ns
BMI (kg/m ²)	25.8 ± 5.1	25.5 ± 4.5	ns	25.9 ± 4.9	23.9 ± 3.1	25.6 ± 4.5	ns	ns	ns	ns	ns	ns
PD Age at onset	59 ± 9.3	54 ± 10	***	58 ± 8.9	49 ± 11.2	52 ± 9.7	ns	***	***	***	**	ns
PD Duration	6 ± 5.1	6.8 ± 5	ns	6 ± 3.7	7.4 ± 5.8	7.3 ± 5.6	ns	ns	ns	ns	ns	ns
H&Y stage (median)	2	2	ns	2	2	2	ns	ns	ns	ns	ns	ns
LEDD	549.2 ± 432.6	645.9 ± 477.6	ns	615 ± 528.6	654.6 ± 474.6	724.7 ± 434.7	ns	ns	ns	ns	ns	ns
DBS (N, %)	15 (8%)	25 (19.4%)	**	6 (11%)	4 (17%)	13 (33%)	ns	ns	*	ns	ns	ns
Apomorphine pump infusion (N, %)	-	1 (0.8%)	-	-	1 (4%)	-	-	-	-	-	-	-
LCIG (N, %)	-	1 (0.8%)	-	-	1 (4%)	-	-	-	-	-	-	-
MDS-UPDRS part I	9.9 ± 6.9	10.9 ± 6.2	ns	10.3 ± 6.4	10.3 ± 5.5	11.8 ± 6.2	ns	ns	ns	ns	ns	ns
MDS-UPDRS part II	10.4 ± 7	11.5 ± 6.1	*	11.1 ± 6.1	11.2 ± 6.7	12 ± 6	ns	ns	ns	ns	ns	ns
MDS-UPDRS part III	27.1 ± 12.9	27.9 ± 13.1	ns	28 ± 12.4	28.4 ± 12.1	28.2 ± 14.6	ns	ns	ns	ns	ns	ns
MDS-UPDRS part IV	2.6 ± 3.4	3.8 ± 3.6	**	3.4 ± 3.3	4.2 ± 4.2	4 ± 3.7	ns	ns	ns (un,p=0.01; adj,p=0.07)	ns	ns	ns
MDS-UPDRS total	49.6 ± 23	53.8 ± 20.7	*	52.8 ± 19.6	54.2 ± 20	55.3 ± 21.2	ns	ns	ns	ns	ns	ns
SCOPA-AUT total	14.3 ± 7.9	14.6 ± 7.7	ns	13.8 ± 7.2	14.2 ± 8	15.9 ± 7.3	ns	ns	ns	ns	ns	ns
SCOPA-AUT Gastrointestinal	3.8 ± 3	4 ± 2.8	ns	3.8 ± 2.6	3.4 ± 2.8	4.6 ± 3.1	ns	ns	ns	ns	ns	ns
SCOPA-AUT Urinary	5.1 ± 3.1	4.7 ± 3	ns	5 ± 3.2	3.9 ± 2.7	4.8 ± 2.6	ns	ns	ns	ns	ns	ns
SCOPA-AUT Cardiovascular	0.7 ± 1	0.9 ± 1.1	ns (p=0.058)	0.8 ± 1	0.8 ± 1.2	1 ± 1.2	ns	ns	ns	ns	ns	ns
SCOPA-AUT Thermoregulatory	2 ± 1.9	2.6 ± 2.4	ns (p=0.084)	2.3 ± 2.1	2.8 ± 3.1	2.5 ± 2.3	ns	ns	ns	ns	ns	ns
SCOPA-AUT Pupilomotor	0.5 ± 0.7	0.5 ± 0.9	ns	0.4 ± 0.8	0.4 ± 0.9	0.8 ± 1	ns	ns	ns	ns	ns (un,p=0.01; adj,p=0.09)	ns (un,p=0.04; adj,p=0.13)
SCOPA-AUT Sexual	1.6 ± 1.8	1.5 ± 1.8	ns	1.2 ± 1.7	1.9 ± 1.9	1.6 ± 1.6	ns	ns	ns	ns	ns	ns
WCSS	5.7 ± 4.4	6.1 ± 4.6	ns	5.2 ± 3.8	5.7 ± 4.2	7.2 ± 5.1	ns	ns	ns (un,p=0.09; adj,p=0.28)	ns	ns (un,p=0.07; adj,p=0.28)	ns
RBDSQ	5 ± 3.6	5.7 ± 3.6	ns	5.6 ± 3.7	5.4 ± 2.7	5.7 ± 3.9	ns	ns	ns	ns	ns	ns
UPSIT	18.8 ± 6.7	17.5 ± 5.5	**	17.5 ± 5.1	18.8 ± 5.2	16.4 ± 6.3	ns	ns	***	ns	ns	ns
HADS anxiety	5.3 ± 3.9	6.4 ± 3.6	*	6.1 ± 3.6	6.6 ± 4.6	6.5 ± 3	ns	ns	ns	ns	ns	ns
HADS depression	4.9 ± 3.5	6.2 ± 3.9	**	5.3 ± 3.7	6.7 ± 3.4	6.8 ± 4.5	ns	ns	**	ns	ns	ns
BDI	8.6 ± 6.5	11.5 ± 6.5	***	10.2 ± 5.9	12.9 ± 8.8	12 ± 5.2	ns	ns	***	ns	*	ns
MOCA	25.7 ± 3.5	25.7 ± 4.1	ns	26.2 ± 3.5	27.2 ± 2.5	24.4 ± 5.1	ns	ns	**	ns	**	**

Legend: BMI, body mass index; BDI, Beck Depression Inventory; DBS, deep brain stimulation; DQS, Dietary Quality Score; HADS, Hospital Anxiety and Depression Scale; H&Y, Hoehn & Yahr; LCIG, levodopa-carbidopa intestinal gel; LEDD, levodopa equivalent daily dose; GBR, Great Britain; ITA, Italy; MDS-UPDRS, Movement Disorder Society (MDS) Unified Parkinson's Disease Rating Scale; MOCA, Montreal Cognitive Assessment; PD, Parkinson disease; SCOPA-AUT, Scales for Outcomes in Parkinson's disease; RBDSQ, REM Sleep Behavior Disorder Questionnaire; UPSIT, University of Pennsylvania Smell Identification Test; WCSS, Wexner Constipation Scoring System. N, negative; M, mild; R, risk; S, severe. p value: comparison between iPD and GBA-PD. P1: negative vs risk; P2: negative vs mild; P3: negative vs severe; P4: risk vs mild; P5: risk vs severe; P6: mild vs severe. P values: ns, p ≥ 0.05; *, p < 0.05; **, p < 0.01; ***, p < 0.001; adj.p, adjusted p value; un.p, unadjusted p value.

Table 2. Performances in specific cognitive abilities (MOCA sub-scores).

MOCA sub-scores	Negative – iPD (N=186)	GBA-PD (N=129)	<i>p</i> value	Risk (N=56)	Mild (N=24)	Severe (N=39)	<i>P1</i> (<i>N</i> vs <i>R</i>)	<i>P2</i> (<i>N</i> vs <i>M</i>)	<i>P3</i> (<i>N</i> vs <i>S</i>)	<i>P4</i> (<i>R</i> vs <i>M</i>)	<i>P5</i> (<i>R</i> vs <i>S</i>)	<i>P6</i> (<i>M</i> vs <i>S</i>)
Diagram	0.9 ± 0.3	0.8 ± 0.4	*	0.9 ± 0.4	0.8 ± 0.4	0.8 ± 0.4	ns	*	*	ns	ns	ns
Cube	0.8 ± 0.4	0.7 ± 0.5	*	0.8 ± 0.4	0.8 ± 0.4	0.6 ± 0.5	ns	ns	***	ns	**	ns
Clock	2.6 ± 0.7	2.5 ± 0.8	*	2.5 ± 0.8	2.6 ± 0.6	2.3 ± 0.9	ns	ns	**	ns	ns	ns
Visuospatial/executive functions	4.2 ± 1.2	4 ± 1.4	*	4.1 ± 1.3	4.1 ± 1.2	3.7 ± 1.5	ns	ns	***	ns	**	ns
Naming	3 ± 0.2	3 ± 0.5	ns	3 ± 0.4	3 ± 0.3	3 ± 0.6	ns	ns	ns	ns	ns	ns
Digits	1.8 ± 0.5	1.8 ± 0.4	ns	1.8 ± 0.4	1.9 ± 0.3	1.8 ± 0.5	ns	ns	ns	ns	ns	ns
Letters	0.9 ± 0.3	0.9 ± 0.3	ns	1 ± 0.2	1 ± 0	0.8 ± 0.4	ns	ns	ns	ns	*	ns
Subtraction	2.7 ± 0.6	2.7 ± 0.7	ns	2.8 ± 0.5	2.9 ± 0.4	2.4 ± 1	ns	ns	*	ns	**	*
Attention	5.4 ± 0.9	5.4 ± 1	ns	5.6 ± 0.7	5.8 ± 0.4	4.9 ± 1.5	ns	ns	ns	ns	**	*
Repeat	1.7 ± 0.5	1.8 ± 0.5	ns	1.8 ± 0.5	1.8 ± 0.4	1.8 ± 0.4	ns	ns	ns	ns	ns	ns
Fluency	0.8 ± 0.4	0.8 ± 0.4	ns	0.7 ± 0.4	0.9 ± 0.4	0.7 ± 0.4	ns	ns	ns	ns	ns	ns
Language	2.5 ± 0.7	2.5 ± 0.7	ns	2.5 ± 0.7	2.7 ± 0.6	2.5 ± 0.8	ns	ns	ns	ns	ns	ns
Abstraction	1.7 ± 0.5	1.7 ± 0.5	ns	1.8 ± 0.5	1.8 ± 0.4	1.6 ± 0.7	ns	ns	ns	ns	ns	ns
Delayed recall	2.8 ± 1.7	3.2 ± 1.5	ns	3.2 ± 1.5	3.8 ± 1.2	2.9 ± 1.6	ns	ns	ns	ns	ns	ns
Orientation	5.9 ± 0.4	5.8 ± 0.6	ns	5.9 ± 0.5	5.9 ± 0.3	5.6 ± 0.8	ns	ns	*	ns	ns	ns

Legend: PD, Parkinson disease. N, negative; M, mild; R, risk; S, severe. *p* value: comparison between iPD and GBA-PD. *P1*: negative vs risk; *P2*: negative vs mild; *P3*: negative vs severe; *P4*: risk vs mild; *P5*: risk vs severe; *P6*: mild vs severe. *P* values: ns: $p \geq 0.05$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

References

1. Vieira SRL, Meزابrovski R, Toffoli M, et al. Consensus Guidance for Genetic Counseling in GBA1 Variants: A Focus on Parkinson's Disease. *Mov Disord* 2024.
2. Skrahina V, Gaber H, Vollstedt EJ, et al. The Rostock International Parkinson's Disease (ROPAD) Study: Protocol and Initial Findings. *Mov Disord* 2021;36(4):1005-1010.
3. Pal G, Mangone G, Hill EJ, et al. Parkinson Disease and Subthalamic Nucleus Deep Brain Stimulation: Cognitive Effects in GBA Mutation Carriers. *Ann Neurol* 2022;91(3):424-435.
4. Menozzi E, Toffoli M, Schapira AHV. Targeting the GBA1 pathway to slow Parkinson disease: Insights into clinical aspects, pathogenic mechanisms and new therapeutic avenues. *Pharmacol Ther* 2023;246:108419.
5. Menozzi E, Schapira AHV. Exploring the Genotype-Phenotype Correlation in GBA-Parkinson Disease: Clinical Aspects, Biomarkers, and Potential Modifiers. *Front Neurol* 2021;12:694764.
6. Petrucci S, Ginevrino M, Trezzi I, et al. GBA-Related Parkinson's Disease: Dissection of Genotype-Phenotype Correlates in a Large Italian Cohort. *Mov Disord* 2020;35(11):2106-2111.
7. Gan-Or Z, Amshalom I, Kilarski LL, et al. Differential effects of severe vs mild GBA mutations on Parkinson disease. *Neurology* 2015;84(9):880-887.
8. Liu G, Boot B, Locascio JJ, et al. Specifically neuropathic Gaucher's mutations accelerate cognitive decline in Parkinson's. *Ann Neurol* 2016;80(5):674-685.
9. Stoker TB, Camacho M, Winder-Rhodes S, et al. Impact of GBA1 variants on long-term clinical progression and mortality in incident Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2020;91(7):695-702.
10. Straniero L, Asselta R, Bonvegna S, et al. The SPID-GBA study: Sex distribution, Penetrance, Incidence, and Dementia in GBA-PD. *Neurol Genet* 2020;6(6):e523.
11. Mata IF, Leverenz JB, Weintraub D, et al. GBA Variants are associated with a distinct pattern of cognitive deficits in Parkinson's disease. *Mov Disord* 2016;31(1):95-102.
12. Davis MY, Johnson CO, Leverenz JB, et al. Association of GBA Mutations and the E326K Polymorphism With Motor and Cognitive Progression in Parkinson Disease. *JAMA Neurol* 2016;73(10):1217-1224.
13. Colucci F, Avenali M, De Micco R, et al. Ambroxol as a disease-modifying treatment to reduce the risk of cognitive impairment in GBA-associated Parkinson's disease: a multicentre, randomised, double-blind, placebo-controlled, phase II trial. The AMBITIOUS study protocol. *BMJ Neurol Open* 2023;5(2):e000535.
14. Higgins AL, Toffoli M, Mullin S, et al. The remote assessment of parkinsonism supporting the ongoing development of interventions in Gaucher disease. *Neurodegener Dis Manag* 2021;11(6):451-458.
15. Parlar SC, Grenn FP, Kim JJ, Baluwendrat C, Gan-Or Z. Classification of GBA1 Variants in Parkinson's Disease: The GBA1-PD Browser. *Mov Disord* 2023;38(3):489-495.
16. Tomlinson CL, Stowe R, Patel S, Rick C, Gray R, Clarke CE. Systematic review of levodopa dose equivalency reporting in Parkinson's disease. *Mov Disord* 2010;25(15):2649-2653.
17. Nyholm D, Jost WH. An updated calculator for determining levodopa-equivalent dose. *Neurol Res Pract* 2021;3(1):58.
18. Harris PA, Taylor R, Minor BL, et al. The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019;95:103208.
19. Team R. A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. In: Core, editor. Available from: <https://www.R-project.org/R-projectViennaAustria.2019>.
20. Thaler A, Bregman N, Gurevich T, et al. Parkinson's disease phenotype is influenced by the severity of the mutations in the GBA gene. *Parkinsonism Relat Disord* 2018;55:45-49.
21. Lunde KA, Chung J, Dalen I, et al. Association of glucocerebrosidase polymorphisms and mutations with dementia in incident Parkinson's disease. *Alzheimers Dement* 2018;14(10):1293-1301.
22. Carandina A, Lazzeri G, Rodrigues GD, et al. Dysautonomia in Parkinson's Disease: Impact of Glucocerebrosidase Gene Mutations on Cardiovascular Autonomic Control. *Front Neurosci* 2022;16:842498.

23. Kim MS, Park DG, An YS, Yoon JH. Dual-phase (18) F-FP-CIT positron emission tomography and cardiac (123) I-MIBG scintigraphy of Parkinson's disease patients with GBA mutations: evidence of the body-first type? *Eur J Neurol* 2023;30(2):344-352.
24. Devigili G, Straccia G, Cereda E, et al. Unraveling Autonomic Dysfunction in GBA-Related Parkinson's Disease. *Mov Disord Clin Pract* 2023;10(11):1620-1638.
25. Narita A, Shirai K, Kubota N, et al. Abnormal pupillary light reflex with chromatic pupillometry in Gaucher disease. *Ann Clin Transl Neurol* 2014;1(2):135-140.
26. Bitirgen G, Turkmen K, Zengin N, Malik RA. Altered pupillary light responses are associated with the severity of autonomic symptoms in patients with Fabry disease. *Sci Rep* 2021;11(1):8146.
27. Menozzi E, Schapira AHV. Prospects for Disease Slowing in Parkinson Disease. *Annu Rev Pharmacol Toxicol* 2024.
28. Giladi N, Alcalay RN, Cutter G, et al. Safety and efficacy of venglustat in GBA1-associated Parkinson's disease: an international, multicentre, double-blind, randomised, placebo-controlled, phase 2 trial. *Lancet Neurol* 2023;22(8):661-671.