

1 **Care seeking for diarrheal illness: a systematic review and meta-analysis**

2

3 Kirsten E. Wiens¹, Marissa H. Miller¹, Daniel J. Costello^{1,4}, Ashlynn P. Solomon¹, Skye M.

4 Hilbert¹, Andrea G. Shipper², Elizabeth C. Lee³, Andrew S. Azman^{3,5}

5

6 ¹ Department of Epidemiology and Biostatistics, College of Public Health, Temple University,

7 Philadelphia, USA

8 ² Charles Library, Temple University, Philadelphia, USA

9 ³ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Johns

10 Hopkins University, Baltimore, USA

11 ⁴ Department of Epidemiology, Milken Institute School of Public Health, The George Washington

12 University, Washington, DC, USA

13 ⁵ Geneva Centre for Emerging Viral Diseases and Division of Tropical and Humanitarian

14 Medicine, Geneva University Hospitals, Geneva, Switzerland

15

16 **Corresponding author**

17 Kirsten E. Wiens, PhD

18 Assistant Professor

19 Department of Epidemiology and Biostatistics

20 Temple University College of Public Health

21 1301 Cecil B. Moore Avenue, Ritter Annex 911

22 Philadelphia, PA 19122

23 kirsten.wiens@temple.edu

24

25 Abstract

26 **Background**

27 Monitoring and treating diarrheal illness often rely on individuals seeking care at hospitals or
28 clinics. Cases that seek care through pharmacies and community health workers (CHW) are
29 frequently excluded from disease burden estimates, which are used to allocate mitigation
30 resources. Studies on care seeking behavior can help identify these gaps but typically focus on
31 children under five, even though diarrheal diseases like cholera and Enterotoxigenic E. coli
32 affect all age groups. This study aimed to estimate the proportion of individuals seeking care for
33 themselves or their children with diarrhea, considering different age groups, case definitions,
34 and study settings.

35

36 **Methods and Findings**

37 We conducted a systematic review of population-based primary research studies published
38 during 2000–2024 that examined care-seeking behavior for diarrhea. We included 166 studies
39 from 62 countries. Hospitals and clinics were the most common source of care sought outside
40 the home, with CHW and health posts rarely reported. Using a random-effects meta-analysis,
41 we found substantial heterogeneity in care seeking between studies from low- and middle-
42 income countries ($I^2 = 99.3$) and estimated that the proportion of diarrhea cases seeking care at
43 a hospital or clinic was 32.8% on average (95% Confidence Interval (CI) 28.1% to 37.9%;
44 prediction interval 3.3% to 87.5%). Although there were trends toward higher care-seeking for
45 children compared to adults, substantial variation existed between studies, and the differences
46 were not significant. We estimated that the adjusted odds of seeking care at a hospital or clinic
47 were significantly higher for severe diarrhea and cholera compared to general diarrhea (Odds
48 Ratio 3.43; 95% CI 1.71 to 6.88).

49

50 **Conclusions**

51 Our findings confirm that passive surveillance through hospitals and clinics may substantially
52 undercount the number of people with diarrhea, particularly those with milder symptoms,
53 although this proportion varied widely. Additionally, our findings underscore the importance of
54 including care seeking questions across all age groups in future studies, as we cannot assume
55 lower care seeking for adults across all settings. Our study was limited by fewer data on care-
56 seeking from health posts, traditional healers, and CHW compared to hospitals and clinics,
57 highlighting a need for further research on these sources of care.

58

59 Introduction

60 Diarrheal diseases are a leading cause of illness and death worldwide, particularly among
61 children under five [1,2]. Diarrhea burden estimates rely partly on data from population-based
62 surveys focused on children, which use caregiver recall of recent symptoms [1,2]. In adults and
63 for specific pathogens, estimates are often based on passive surveillance data, such as hospital
64 records and vital statistics [1,2]. These data are especially valuable for tracking seasonal
65 outbreaks of pathogens like cholera that may not appear in surveys but depend on individuals
66 seeking care for often severe symptoms [3,4]. A key challenge in accurately assessing diarrheal
67 disease burden by age and etiology is determining how many cases go unreported due to
68 barriers in care-seeking and identifying which populations or illnesses are most likely to be
69 undercounted.

70

71 Estimates of diarrhea burden strongly influence investments in disease mitigation resources,
72 including broad investments in improved water and sanitation and more pathogen-specific tools
73 like vaccines. However, diarrheal pathogens vary in the symptoms they cause as well as the
74 age groups they primarily affect. For example, rotavirus, *Escherichia coli*, and Salmonella can
75 cause gastroenteritis with diarrhea and/or vomiting, while cholera is marked by acute watery
76 diarrhea and/or vomiting. Rotavirus primarily affects children, whereas cholera is a leading

77 cause of diarrhea illness and deaths across all age groups [1]. Strategies for controlling
78 rotavirus differ from those used for cholera, as its primary mode of transmission is person-to-
79 person via the fecal-oral route, rather than through consuming contaminated food and water.
80 Enteric pathogens also cause a range of disease severity, from mild infection to severe
81 dehydration requiring rehydration to prevent severe outcomes including death. Given these
82 variations, relying solely on population-level estimates of general diarrhea in children and
83 passive surveillance data may not optimize burden reduction efforts.

84

85 Programs such as the Demographic and Healthcare Surveys (<https://dhsprogram.com/>) that
86 assess care seeking behavior for diarrhea around the world provide an opportunity to examine
87 these biases in disease burden estimates [5–9]. Previous analyses of these data have
88 estimated that the proportion of caregivers that seek care at health facilities for a child with
89 diarrhea was on average 45% in sub-Saharan Africa [10] and 49-51% in West and Central
90 Africa [11], with substantial variation by country [12] and type of care sought [13]. Importantly,
91 these studies all focused on care seeking for children, and did not incorporate data from studies
92 that have reported care seeking across age groups including adults [14,15]. In addition, while
93 one of these reviews found that more people sought care for general childhood illnesses when
94 illness was more severe [13], none of these studies examined care seeking behavior by severity
95 of diarrheal illness or etiology, limiting the applicability of their results.

96

97 In this study, we sought to address these knowledge gaps through a systematic review and
98 meta-analysis of surveys conducted during 2000-2024 on diarrhea care seeking behavior
99 across all age groups and diarrhea case definitions. Our primary research question was: what
100 proportion of individuals of any age seek care at hospitals, clinics, or other sources when they
101 have diarrhea? In addition, we asked: how does seeking care at hospitals or clinics vary by age
102 group, severity of diarrheal illness, and other factors?

103

104 Methods

105

106 **Systematic review**

107 We developed the methods for this study based on the Preferred Reporting Items for
108 Systematic Reviews and Meta-Analyses (PRISMA) guidelines [16] and submitted the protocol to
109 the International Prospective Register of Systematic Reviews (PROSPERO) on February 24th,
110 2023 (review ID: CRD42023402435) [17].

111

112 We searched PubMed, Embase, Web of Science, and Global Index Medicus on January 27,
113 2023 using search terms developed by a medical librarian (AGS); searches were updated on
114 September 3, 2024 (Supplementary Methods). We included studies that 1) included sampling
115 on or after January 1, 2000, 2) were primary research studies, 3) were conducted in human
116 populations, 4) included questions about diarrhea care seeking, perceptions knowledge, and/or
117 management, 5) surveyed individuals of any age about their own care seeking for diarrheal
118 illness, 6) reported the number of individuals surveyed about care seeking for diarrheal illness
119 either for themselves or a child, 7) reported the number or proportion who said that they would
120 or did seek care, and 8) were written in English, French or Spanish. We excluded studies that 1)
121 were conducted in special populations, e.g., HIV- or malaria-infected individuals, healthcare
122 professionals, deployed troops, travelers, or long-term care facility residents, 2) focused
123 exclusively on non-infectious diarrhea, 3) did not separate diarrhea-related care seeking from
124 other illnesses, 4) did not report the sampling method and study population, 5) were case
125 studies, case-control studies where cases selected based on care seeking behavior, or
126 qualitative studies, 6) used convenience sampling, 7) were reviews or secondary data analyses,
127 8) were conference abstracts, or 9) did not have full texts available.

128

129 Studies were screened by title and abstract by two blinded and independent reviewers (DJC,
130 APS, MHM, or KEW) using Covidence (<https://www.covidence.org/>). If a tiebreaker was needed,
131 a third reviewer was used, or a decision was made by consensus. This process was repeated
132 for full text screening. For the updated search, a single reviewer (MHM) performed the
133 screening. One of the reviewers independently extracted data from each included study using a
134 shared Google Sheet. Extracted data included study design, sampling method, study
135 population, diarrhea case definition, start and end dates, study catchment area matched to a
136 GADM polygon (<https://gadm.org/>) when possible, how the care seeking questions were
137 phrased, who the care was sought for (e.g., self or child), recall time in days, type of care facility
138 sought, number of individuals surveyed about diarrheal illness, whether they “would” and/or
139 “did” seek care, and when available, age and sex distribution of the surveyed population,
140 diarrhea symptoms, and diarrhea severity. When care seeking data were stratified by any of
141 these variables, we extracted the data at those stratifications. If a study reported “yes”, “no”, and
142 “maybe” for “would/will you seek care”, we coded “maybe” as “no”.

143

144 A single reviewer (SMH) evaluated the quality of the studies that met our inclusion criteria using
145 an adapted version of the appraisal tool for cross-sectional studies (AXIS) [18]. Specifically, we
146 assessed whether 1) the data presented in the study were internally consistent (i.e., numbers
147 for the proportion seeking care were identical throughout the abstract, text, and tables), 2) the
148 manuscript included a justification for their sample size, and 3) the response rate was reported
149 [0 = no, 1 = yes]. An overall quality score was created by summing across the three indicators
150 for total scores ranging from 0-3, where 0 was lowest quality and 3 was highest quality. Studies
151 were not excluded based on quality assessment criteria. We retained studies with internal
152 inconsistencies to avoid penalizing those that provided more detailed information or in-depth
153 discussions of results, where minor differences or ambiguities often arose. Studies with major

154 ambiguities were excluded in full text screening for lacking care seeking data (for example, if it
155 was unclear what the denominator was for the reported proportion seeking care).

156

157 **Data cleaning**

158 We manually reviewed studies by country, sorted by start date, to identify studies with
159 overlapping data (i.e., conducted among the same people in the same time period with the
160 same questions). Overlapping studies were excluded from the primary dataset, and reason for
161 selection of the retained study was recorded in the extraction sheet. We prioritized inclusion of
162 studies with 1) more representative sampling methods, 2) more consistent or clearly described
163 data, 3) more specific case definition, 4) larger sample size, 5) longer sampling period, 6) more
164 detailed age information, and 7) more detailed methods description, in that order.

165

166 Data were cleaned, checked for mistakes, and analyzed using R statistical software version
167 4.3.3 [19]. Extracted data were checked for correct location information (e.g., could be correctly
168 matched to a GADM shapefile, impossible values (e.g., number seeking care greater than
169 number surveyed, start date after end date), and inconsistent values (e.g., multiple different
170 sample sizes entered per study or stratification). Data were matched with World Bank 2023-
171 2024 Income Classifications [20] and regions by country.

172

173 We created three standardized categories for diarrhea case definition: 1) “diarrhea” included
174 explicit definitions of three or more loose stools in past 24 hours and diarrhea used broadly
175 without an explicit definition; 2) “severe diarrhea or cholera” included severe diarrhea, defined
176 as having danger signs, dehydration, ≥ 1 week duration, or death, and acute watery diarrhea of
177 any severity including deaths; 3) “gastroenteritis or other etiologies” included definitions of either
178 diarrhea or vomiting, and specific etiologies including rotavirus, Escherichia coli, Salmonella,
179 and giardia. A single study stratified care seeking by diarrhea cases that were “moderate to

180 severe” (diarrhea with sunken eyes, loss of skin turgor, dysentery, intravenous rehydration, or
181 hospitalization) and “less severe” (all other diarrhea) [21]; we categorized “moderate to severe”
182 as “severe diarrhea or cholera” and “less severe” as “diarrhea”.

183

184 In addition, we created standardized categories of the source of care sought (Table S1). For the
185 analysis of care seeking at hospitals or clinics, we defined “hospital or clinic” as any general
186 hospital or clinic or explicitly defined public hospitals or clinics. We excluded explicitly defined
187 private hospitals or clinics from this category because they are rarely part of surveillance
188 systems.

189

190 **Meta-analysis**

191 Observations, or units for analysis, were created by aggregating extracted data by study,
192 country, source of care sought, and potential sources of variation in care seeking (shown in
193 Table 1). When aggregating data from studies with multiple choice questions where there were
194 multiple different options within one category or source of care (e.g., multiple different hospitals
195 or clinics that could have been selected), we took the maximum value across those options.

196

197 We used logistic regression models with observation-level random intercepts to pool estimates
198 of the proportion of individuals with diarrhea that did or would seek care for diarrhea at hospitals
199 or clinics across all included studies, implemented using the `metaprop` function of the `meta`
200 package in R [22]. We used logistic regression models with observation-level random intercepts
201 and fixed effects to examine the effects of diarrhea case definition, whether the study was in an
202 urban or rural location, whether the study took place during or closely following a diarrhea
203 outbreak, as well as other components of study methodology on care seeking in univariate and
204 multivariate analyses, implemented using the `rma` function of the `metafor` package in R [23]. For
205 multivariate analysis, we included variables that we identified as significantly associated with

206 care seeking ($p < 0.05$) in univariate analysis but that were not significantly associated with
207 other included variables. In addition, we performed sub-group analyses by age group and
208 diarrhea case definition.

209

210 **Data and code availability**

211 All extracted data and source code for data cleaning and meta-analysis are available at
212 <https://github.com/wienslab/diarrhea-careseeking>.

213

214 **Role of the funding source**

215 This study was supported by the Bill and Melinda Gates Foundation (INV-044865) and the
216 National Institute of Allergy and Infectious Diseases (K22AI168389). The funders had no role in
217 study design, data collection and analysis, decision to publish, or preparation of the manuscript.

218

219 Results

220 **Study characteristics**

221 We retrieved 7069 studies from four databases, screened 5188 for eligibility following de-
222 duplication, and included 188 studies (Figure 1). Of these, 166 studies were non-overlapping
223 and asked participants whether they did seek care for a past diarrheal illness. For the primary
224 analysis, we stratified these study-level data by characteristics shown in Table 1, for a total of
225 211 observations pertaining to care seeking outside of participants' homes, and 177
226 observations pertaining to care seeking specifically at hospitals or clinics. The complete
227 analytical datasets can be found in Supplementary Data 1.

228

229 The studies included in the primary dataset were conducted in 62 countries across 10 World
230 Bank geographic regions, with most observations coming from East Africa (N = 60), South Asia
231 (N = 41), Latin America and the Caribbean (N = 25), Western Africa (N = 24), and East Asia and

232 the Pacific (N = 23). Twenty of the observations came from nationally representative studies,
233 and 53, 78, and 60 came from studies representative of first, second, and third administrative
234 divisions, respectively (Figure S1). Studies included sampling completion dates between 2000
235 and 2022, with most of the observations completed during 2005–2009 (N = 67) and 2010–2014
236 (N = 67) (Figure S2).

237

238 Most observations in the primary dataset were from studies that asked a caregiver about care
239 seeking for a child with diarrhea (N = 138, 65.4%) (Table 1). Others represented a combination
240 of children, other family members, or the respondent themselves (N = 59, 28.0%) and a small
241 proportion represented care seeking exclusively for the survey respondents (N = 14, 6.6%)
242 (Table 1). A small proportion of observations represented responses from caregivers of
243 individuals at a health facility (N = 11, 5.2%) (Table 1). More of the observations were from
244 exclusively rural areas (N = 83, 39.3%) than exclusively urban areas (N = 68, 32.2%), and two
245 (0.9%) came from internally displaced persons or refugees (Table 1). Seventeen (8.1%)
246 observations took place during or closely following a diarrheal disease outbreak (Table 1).

247

248 The majority of the observations came from surveys where respondents were asked about care
249 seeking for general diarrhea (N = 156, 73.8%), followed by either gastroenteritis or non-cholera
250 etiologies (N = 33, 15.6%), and severe diarrhea or cholera including deaths (N = 22, 10.4%)
251 (Table 1). Most observations focused on care seeking at any time during illness (N = 175,
252 82.9%) and most used recall periods of 14-30 days (N = 143, 67.8%) (Table 1).

253

254 **Trends in care seeking at hospitals or clinics in the unadjusted data**

255 Our primary interest was care-seeking at hospitals or clinics, where people can receive
256 treatment and are most often counted in disease surveillance efforts. Since care seeking
257 behavior among individuals at a clinic likely differs from care seeking behavior in the broader

258 community, we restricted the data in the analyses below to non-clinic-based studies unless
259 otherwise noted.

260

261 We observed variations in care-seeking by geographic region (Figure S3a) and country income
262 group (Figure S4a). These trends were inconsistent between case definitions, which also varied
263 by income group (Figure S3b, Figure S4b). Specifically, in high-income countries (HICs), the
264 most common case definitions were gastroenteritis or non-cholera etiologies, while severe
265 diarrhea and cholera were more often studied in low- and middle-income countries (LMICs).
266 Given a relatively low number of observations from high-income countries, we focus on care-
267 seeking behavior in LMICs in the main analyses.

268

269 We found that the reported proportion seeking care at a hospital or clinic in 145 observations
270 from LMICs was higher for observations from studies that surveyed participants about care
271 seeking at any time during illness (median of 34.7%; N = 125; IQR, 18.9% to 55.0%) compared
272 to their first source of care (median of 21.2%; N = 20; IQR, 10.0% to 33.5%) (Figure 2a) as well
273 as studies conducted during or following a diarrheal disease outbreak (median of 62.7%; N =
274 11; Interquartile Range (IQR), 44.2% to 65.5%) compared to those that were not (median of
275 31.0%; N = 134; IQR, 15.3% to 49.9%) (Figure 2c). Care seeking at hospitals or clinics was also
276 higher for severe diarrhea or cholera (median of 64.1%; N = 15; IQR, 45.4% to 70.0%)
277 compared to general diarrhea (median of 31.0%; N = 122; IQR, 15.3% to 48.2%) and
278 gastroenteritis or other etiologies (median of 33.3%; N = 8; IQR, 25.5% to 44.1%) (Figure 2b).
279 Although we found no significant correlation between overall study quality and reported care
280 seeking at hospitals or clinics (Spearman rho = 0.07; 95% Confidence Interval -0.09 to 0.23;
281 Figure S5a), studies with internal inconsistencies identified during study quality assessment
282 more often reported lower care seeking than those without (Figure S5b).

283

284 Studies that surveyed participants about their own care-seeking behavior reported lower levels
285 of care-seeking on average compared to those focused on children (Figure 2d). Yet, across 134
286 observations in LMICs that reported the age distribution of participants, we found no significant
287 correlation between the proportion of the study population under five years of age and the
288 proportion that sought care at hospitals or clinics (Spearman rho = 0.10; 95% Confidence
289 Interval -0.08 to 0.26; Figure 3a). Studies conducted only among children reported higher care
290 seeking at hospitals or clinics (median of 34.2%; N = 98; IQR, 17.5% to 55.0%) compared to
291 studies conducted only among adults or individuals over five (median of 22.4%; N = 22; IQR,
292 14.5% to 49.3%) (Figure 3c), but there was wide variation between studies and this trend held
293 for diarrhea and gastroenteritis but not for severe diarrhea or cholera (Figure 3b,d).

294

295 **Trends in care seeking at other sources outside the home in LMICs**

296 Hospitals and clinics were the most common source of care sought across studies in LMICs
297 (median of 32.3%; N = 145; IQR, 17.0% to 52.6%) (Figure 4a), though they were less commonly
298 the first source of care (median of 21.2%; N = 20; IQR, 10.0% to 33.5%) (Figure 4b). Seeking
299 care from community health workers (CHW) was reported more often as first sources of care
300 (median of 16.9%; N = 9; IQR, 4.5% to 18.9%) than the main or primary of care (median of
301 6.2%; N = 32; IQR, 1.7% to 14.7%) (Figure 4b). Seeking care from CHW and health posts were
302 also more frequently reported in observations from low-income compared to middle-income
303 countries (Figure S6) and for general diarrhea compared to severe diarrhea or cholera (Figure
304 S7). Traditional healers were rarely reported as a source of care (median of 4.0%; N = 48; IQR,
305 1.0% to 8.7%) (Figure 4). These less common sources were also less frequently included or
306 reported as options in survey questions than hospitals or clinics (Figure 4).

307

308 In studies that surveyed participants at a hospital or clinic, most participants reported
309 hospitals/clinics as their first source of care (median of 89.3%; N = 4; IQR, 74.8% to 97.7%)

310 (Figure S8a). In studies that surveyed participants about where they sought care before their
311 hospital or clinic visit, hospitals/clinics were the most common previous sources of care,
312 (median of 35.1%; N = 2; IQR, 31.2% to 39.0%), followed by pharmacies (median of 20.7%; N =
313 2; IQR, 18.7% to 20.4%), private hospitals/clinics (median of 10.6%; N = 2; IQR, 9.4% to
314 11.9%), and traditional healers (median of 5.6%; N = 2; IQR, 5.4% to 5.8%) (Figure S8b).

315

316 **Adjusted estimates of care seeking at hospitals or clinics**

317 Using a random-effects meta-analysis, we estimated that on average 32.8% (95% Confidence
318 Interval (CI) 28.1% to 37.9%) of individuals sought care at hospitals or clinics across 145
319 observations in LMICs (Figure S9). In addition, we predicted that care seeking for a future study
320 would be in the range of 3.3% to 87.5% (Figure S9). This wide prediction interval reflects the
321 substantial variation we found between studies in care seeking ($\tau^2 = 1.8$; $I^2 = 99.3$; 95% CI, 99.2
322 to 99.3), with adjusted study-level estimates of care seeking ranging from 0.7% (95% CI 0.5% to
323 0.9%) to 94.4% (95% CI 90.6% to 97.0%) (Figure S9).

324

325 We estimated that the proportion of people that would seek care during a hypothetical diarrheal
326 episode, as opposed to an actual previous episode, in LMICs was substantially higher, with
327 71.2% (95% CI 57.1% to 83.2%; prediction interval 5.8% to 99.1%) intending to seek care at a
328 hospital or clinic if they or their child had diarrhea across 29 observations (Figure S10). Care
329 seeking was on average lower in HICs, with 23.9% (95% CI 16.2% to 33.8%; prediction interval
330 2.5% to 79.6%) seeking care at a hospital or clinic when they or their child had a recent
331 diarrheal episode (Figure S11).

332

333 **Factors associated with variation in care seeking at hospitals or clinics in LMICs**

334 We next examined factors associated with variation in care seeking in LMICs. In univariate
335 analyses, we found that the odds of seeking health care at a hospital or clinic were higher for

336 severe diarrhea or cholera than general diarrhea, as well as during or after an outbreak and
337 when a recall period exceeded 30 days (Table S2). If the question was restricted to the first
338 source of care sought, the odds of seeking care at a hospital or clinic were lower (Table S2). We
339 found no significant differences in care seeking between observations conducted among only
340 children under five compared to those conducted only among individuals over five (OR 1.06
341 (95% CI 0.55 to 2.01)) (Table 2). In a subset of eight studies with results stratified by age,
342 differences in care seeking for individuals under five years of age compared to over five years
343 remained non-significant, though there was a trend towards higher care seeking for children
344 under five (OR 1.24 (95% CI 0.29 to 5.36)) (Table 2).

345

346 To examine potential confounding between the sources of variation in our dataset, we
347 additionally tested for associations between the variables related to study design and setting
348 specific attributes included in the univariate analyses above. We found that studies with severe
349 diarrhea or cholera case definitions were associated with longer recall periods and were more
350 likely to have been conducted during or after an outbreak compared with studies on general
351 diarrhea (Table S3). In sub-analysis, we found that there was no longer an effect of being in an
352 outbreak setting when subsetting to studies with severe diarrhea or cholera case definitions
353 (Table S4). In addition, there was no longer an effect of recall period when subsetting the data
354 to observations with general diarrhea case definitions (Table S4).

355

356 Since we could not separate the effects of case definition from recall period or outbreak context,
357 we examined the effects of case definition on care seeking at hospitals or clinics adjusting only
358 for the timing of care seeking (i.e., whether they were surveyed about their first source of care or
359 care at any time during illness). The adjusted odds of seeking care for severe diarrhea or
360 cholera remained significantly higher than for general diarrhea (OR 3.43 (95% CI 1.71 to 6.88))
361 (Table 3), and residual heterogeneity remained high ($\tau^2 = 1.57$; $I^2 = 99.5$; $R^2 = 10.2\%$).

362 Correspondingly, when we stratified the meta-analysis by case definition, we estimated that
363 58.6% (95% CI 39.9% to 75.2%; Prediction Interval 5.2% to 97.3%) of individuals sought care at
364 a hospital or clinic when they or their child had severe diarrhea or cholera (Figure S13)
365 compared to 29.9% (95% CI 25.3% to 35.1%; Prediction Interval 3.1% to 84.9%) for general
366 diarrhea (Figure S12) and 35.8% (95% CI 23.7% to 50.0%; Prediction Interval 6.5% to 81.8%)
367 for gastroenteritis and non-cholera etiologies (Figure S14).

368

369 Discussion

370 Here we estimated that most people with diarrhea (on average 67%) do not seek care at
371 hospitals or clinics where they can receive treatment when needed and be counted in typical
372 surveillance systems. We found substantial variation between studies, even after accounting for
373 sources of variation such as case definition and timing of care seeking. Hospitals and clinics
374 were the most common source of care investigated and reported, with pharmacies, CHW, and
375 health posts less commonly reported, and traditional healers very rarely cited as a source of
376 care. Studies that phrased questions in terms of hypothetical care seeking reported substantially
377 higher care seeking than those that asked about actual diarrheal episodes. We found no
378 significant differences in care seeking between children under five and older individuals, and the
379 odds of seeking care for severe diarrhea and cholera were significantly higher compared to
380 general diarrhea.

381

382 Our findings shed light on the extent to which we may underestimate diarrhea burden when we
383 rely solely on passive surveillance data, with the greatest underestimates occurring,
384 unsurprisingly, among individuals with mild or moderate symptoms. Additionally, our findings
385 suggest that the relationship between age and care-seeking behavior varies by location,
386 context, and etiology. While we found trends towards adults seeking care less often for
387 themselves than for children with diarrhea, we found that, on average, adults sought care for

388 severe diarrhea and cholera at similar rates as for children. Given these results and the
389 considerable heterogeneity across studies, we recommend that local data on care-seeking
390 behavior, stratified by age and either symptom severity or etiology, be used to adjust burden
391 estimates whenever possible. In the absence of local data or for global analyses, the case-
392 definition-stratified estimates from this review may provide useful approximations of missed
393 symptomatic cases.

394

395 Although we estimated lower average care-seeking rates than previous analyses of DHS data
396 (33% here vs. 45-51% in DHS data)—possibly due to our inclusion of a broader range of study
397 settings and populations—our findings of very low reported care-seeking at CHW and traditional
398 healers align with those studies [5–9]. This may reflect true patterns of care-seeking behavior,
399 but it could also indicate a research bias, as hospitals and clinics were more frequently included
400 in survey questionnaires than other sources of care. Additionally, only a minority of studies
401 reported results where multiple sources of care were selected, and individuals may be less likely
402 to report CHW or traditional healers as their sole or primary source of care. This is supported by
403 our finding that CHW were reported more frequently, and hospitals and clinics less frequently, in
404 studies that examined the first source of care rather than overall source of care. These
405 limitations could have led us and others to under-estimate the role of CHW and traditional
406 healers in diarrhea case management.

407

408 This study has several additional limitations. Only a small proportion of the included studies
409 were conducted among adults or individuals over five years old and care seeking in those
410 studies varied widely, which may have prevented us from identifying significant trends.
411 Methodology and questionnaires also varied between studies, which we may not have fully
412 captured in our analysis, and there were several variables we could not examine due to data
413 scarcity or potential confounding (e.g., income group, outbreak setting, and recall period).

414

415 We also estimated that more than twice as many people would seek care as actually did, but we
416 were not able to examine factors that shape care seeking behavior in detail. Decisions about
417 whether and where to seek care are complex, with tradeoffs and determining factors that vary
418 by location, time, and individuals. Existing theories for health service utilization describe a wide
419 range of factors influencing care seeking, including environmental and health system factors
420 (e.g., provider and treatment availability, transportation and access), population and socio-
421 demographic factors (e.g., age, gender, income, culture, community support, disease
422 prevalence), and individual behavior (e.g., perceptions about disease risk and severity, health
423 care costs, and benefits) [24]. The huge heterogeneity we found in care seeking, even after
424 accounting for different aspects of study methodology, may result from the propagation of
425 individual heterogeneity in decision making and contextual factors. An important area for future
426 research will be to examine these factors to identify ways in which care seeking could be
427 improved, especially for severe cases.

428

429 Given our findings, and their limitations, we propose several areas for future work. First, it will be
430 important for future studies to include questions about care seeking across ages and diarrhea
431 case definitions, include a broad range of sources of care including health posts and CHW, and
432 focus on actual rather than hypothetical care seeking where possible. In addition, studies that
433 explicitly examine the order that individuals seek care at different sources (for example, [25–27])
434 can help inform when and why there may be delays in seeking care at hospitals or clinics and
435 how this varies by age and disease severity. Finally, synthesizing qualitative studies that
436 examine reasons why individuals do or do not seek care could provide insight into non-hospital-
437 based care seeking as well as what needs to be done to reduce barriers for populations with the
438 least interaction with the health system.

439

440 In summary, our findings suggest that care-seeking estimates for broadly defined diarrhea may
441 not reflect true patterns, particularly for severe symptoms. Conclusions about care-seeking
442 behaviors across age groups were less straightforward, as we found trends but no significant
443 differences between children under five and older individuals. DHS data may reasonably
444 approximate general care-seeking patterns for non-severe, non-etiology-specific diarrhea for the
445 locations in which they were collected and in the absence of additional information. When
446 possible, assumptions about the relationship between age and care seeking should be setting-
447 specific, as adult care-seeking was not universally equivalent or lower.

448

449 Funding

450 This study was supported by the Bill and Melinda Gates Foundation (INV-044865 to ASA) and
451 the National Institute of Allergy and Infectious Diseases (K22AI168389 to KEW).

452

453 Conflicts of interest

454 The authors declare no conflicts of interest.

455

456 References

- 457 1. Troeger C, Blacker BF, Khalil IA, Rao PC, Cao S, Zimsen SR, et al. Estimates of the
458 global, regional, and national morbidity, mortality, and aetiologies of diarrhoea in 195
459 countries: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*
460 *Infectious Diseases*. 2018;18: 1211–1228. doi:10.1016/S1473-3099(18)30362-1
- 461 2. Troeger CE, Khalil IA, Blacker BF, Biehl MH, Albertson SB, Zimsen SRM, et al. Quantifying
462 risks and interventions that have affected the burden of diarrhoea among children younger
463 than 5 years: an analysis of the Global Burden of Disease Study 2017. *The Lancet*
464 *Infectious Diseases*. 2020;20: 37–59. doi:10.1016/S1473-3099(19)30401-3
- 465 3. Perez-Saez J, Zheng Q, Kaminsky J, Zou K, Demby MN, Alam C, et al. A decade of
466 shifting cholera burden in Africa and its implications for control: a statistical mapping
467 analysis. *medRxiv*; 2024. p. 2024.09.23.24314072. doi:10.1101/2024.09.23.24314072
- 468 4. Ali M, Nelson AR, Lopez AL, Sack DA. Updated global burden of cholera in endemic
469 countries. *PLOS Neglected Tropical Diseases*. 2015;9: e0003832.
470 doi:10.1371/journal.pntd.0003832

- 471 5. Page AL, Hustache S, Luquero FJ, Djibo A, Manzo ML, Grais RF. Health care seeking
472 behavior for diarrhea in children under 5 in rural Niger: results of a cross-sectional survey.
473 BMC Public Health. 2011;11: 389. doi:10.1186/1471-2458-11-389
- 474 6. Fissehaye T, Damte A, Fantahun A, Gebrekirstos K. Health care seeking behaviour of
475 mothers towards diarrheal disease of children less than 5 years in Mekelle city, North
476 Ethiopia. BMC Research Notes. 2018;11: 749. doi:10.1186/s13104-018-3850-3
- 477 7. Nasrin D, Wu Y, Blackwelder WC, Farag TH, Saha D, Sow SO, et al. Health care seeking
478 for childhood diarrhea in developing countries: evidence from seven sites in Africa and
479 Asia. Am J Trop Med Hyg. 2013;89: 3–12. doi:10.4269/ajtmh.12-0749
- 480 8. Nhampossa T, Mandomando I, Acacio S, Nhalungo D, Sacoora C, Nhacolo A, et al. Health
481 care utilization and attitudes survey in cases of moderate-to-severe diarrhea among
482 children ages 0–59 months in the district of Manhica, Southern Mozambique. Am J Trop
483 Med Hyg. 2013;89: 41–48. doi:10.4269/ajtmh.12-0754
- 484 9. Winter R, Wang W, Florey L, Pullum T. Levels and trends in care seeking for childhood
485 illness in USAID MCH priority countries. 2015 [cited 10 Jan 2023]. Available:
486 <https://dhsprogram.com/publications/publication-cr38-comparative-reports.cfm>
- 487 10. Yaya S, Odusina EK, Adjei NK. Health care seeking behaviour for children with acute
488 childhood illnesses and its relating factors in sub-Saharan Africa: evidence from 24
489 countries. Tropical Medicine and Health. 2021;49: 95. doi:10.1186/s41182-021-00385-1
- 490 11. Bogler L, Weber A-C, Ntambi J, Simen-Kapeu A, Zagre NM, Ekpini RE, et al. Health-care
491 seeking for childhood diseases by parental age in Western and Central Africa between
492 1995 and 2017: A descriptive analysis using DHS and MICS from 23 low- and middle-
493 income countries. J Glob Health. 2021;11: 13010. doi:10.7189/jogh.11.13010
- 494 12. Hodgins S, Pullum T, Dougherty L. Understanding where parents take their sick children
495 and why it matters: a multi-country analysis. Global Health: Science and Practice. 2013;1:
496 328–356. doi:10.9745/GHSP-D-13-00023
- 497 13. Geldsetzer P, Williams TC, Kirolos A, Mitchell S, Ratcliffe LA, Kohli-Lynch MK, et al. The
498 recognition of and care seeking behaviour for childhood illness in developing countries: a
499 systematic review. PLOS ONE. 2014;9: e93427. doi:10.1371/journal.pone.0093427
- 500 14. Abuzerr S, Nasser S, Yunesian M, Hadi M, Mahvi AH, Nabizadeh R, et al. Prevalence of
501 diarrheal illness and healthcare-seeking behavior by age-group and sex among the
502 population of Gaza strip: a community-based cross-sectional study. BMC Public Health.
503 2019;19: 704. doi:10.1186/s12889-019-7070-0
- 504 15. Chowdhury F, Khan IA, Patel S, Siddiq AU, Saha NC, Khan AI, et al. Diarrheal illness and
505 healthcare seeking behavior among a population at high risk for diarrhea in Dhaka,
506 Bangladesh. PLOS ONE. 2015;10: e0130105. doi:10.1371/journal.pone.0130105
- 507 16. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The
508 PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ.
509 2021;372: n71. doi:10.1136/bmj.n71

- 510 17. Wiens KE, Ashlynn SP, Costello DJ, Shipper AG, Lee EC, Azman AS. Care seeking for
511 diarrheal illness: a systematic review and meta-analysis. PROSPERO CRD42023402435.
512 2023 [cited 15 Nov 2023]. Available:
513 https://www.crd.york.ac.uk/prospéro/display_record.php?ID=CRD42023402435
- 514 18. Downes MJ, Brennan ML, Williams HC, Dean RS. Development of a critical appraisal tool
515 to assess the quality of cross-sectional studies (AXIS). *BMJ Open*. 2016;6: e011458.
516 doi:10.1136/bmjopen-2016-011458
- 517 19. R Core Team. R: a language and environment for statistical computing. R Foundation for
518 Statistical Computing, Vienna, Austria. 2024. Available: <https://www.R-project.org/>
- 519 20. World Bank Data Help Desk. World Bank Country and Lending Groups. 2024 [cited 6 May
520 2024]. Available: [https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-](https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups)
521 [world-bank-country-and-lending-groups](https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups)
- 522 21. Omore R, O'Reilly CE, Williamson J, Moke F, Were V, Farag TH, et al. Health care-
523 seeking behavior during childhood diarrheal illness: results of health care utilization and
524 attitudes surveys of caretakers in western Kenya, 2007-2010. *Am J Trop Med Hyg*.
525 2013;89: 29–40. doi:10.4269/ajtmh.12-0755
- 526 22. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical
527 tutorial. *BMJ Ment Health*. 2019;22: 153–160. doi:10.1136/ebmental-2019-300117
- 528 23. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *Journal of*
529 *Statistical Software*. 2010;36: 1–48. doi:10.18637/jss.v036.i03
- 530 24. Gliedt JA, Spector AL, Schneider MJ, Williams J, Young S. A description of theoretical
531 models for health service utilization: a scoping review of the literature. *Inquiry: A Journal of*
532 *Medical Care Organization, Provision and Financing*. 2023;60: 00469580231176855.
533 doi:10.1177/00469580231176855
- 534 25. Escobedo AA, Almirall P, Ávila I, Salazar Y, Alfonso M. Care-seeking behaviour and
535 diagnostic processes for symptomatic giardiasis in children attending an academic
536 paediatric hospital. *Pathogens and Global Health*. 2014;108: 271–278.
537 doi:10.1179/2047773214Y.0000000152
- 538 26. Newbrander W, Ickx P, Werner R, Mujadidi F. Compliance with referral of sick children: a
539 survey in five districts of Afghanistan. *BMC Pediatrics*. 2012;12: 46. doi:10.1186/1471-
540 2431-12-46
- 541 27. Soremekun S, Kasteng F, Lingam R, Vassall A, Kertho E, Settumba S, et al. Variation in
542 the quality and out-of-pocket cost of treatment for childhood malaria, diarrhoea, and
543 pneumonia: Community and facility based care in rural Uganda. *PLOS ONE*. 2018;13:
544 e0200543. doi:10.1371/journal.pone.0200543
- 545
- 546
- 547

548 Tables

549 **Table 1. Study characteristics.** Number of observations in the primary dataset with each characteristic.

550 A study has more than one observation if it reported results stratified by any of these characteristics.

Category	Characteristic	Number (N = 211)	Percent (%)
Timing of care seeking	Care sought at any time	175	82.9
	First source sought	29	13.7
	Care sought prior to current hospital visit	6	2.8
	Care sought within 24 hours of symptom onset	1	0.5
Relationship of diarrhea case to survey respondent	Child (of survey respondent)	138	65.4
	Self (survey respondent)	14	6.6
	Combination (child, self, or other family member)	59	28.0
Survey recall period	2-7 days before interview	13	6.2
	14-30 days before interview	143	67.8
	42-365 days before interview	18	8.5
	Not reported	37	17.5
Clinic-based surveillance	No	198	93.8
	Yes	13	6.2
During or recently after an outbreak that causes diarrhea	No	194	91.9
	Yes	17	8.1
Multiple choices for types of care sought	No	180	85.3
	Yes	31	14.7

Study location type	Rural	83	39.3
	Urban	68	32.2
	Urban and Rural	52	24.6
	Peri-Urban	5	2.4
	Internally Displaced Persons or Refugee Camp	2	0.9
	Urban and Peri-Urban	1	0.5
Country World Bank 2023-2024 income group [20]	Lower middle income	97	46.0
	Upper middle income	44	20.9
	Low income	43	20.4
	High income	27	12.8
Diarrhea case definition	Diarrhea	156	73.9
	Gastroenteritis or other etiologies	33	15.6
	Severe diarrhea or cholera	22	10.4

551
552
553
554
555
556
557
558

Table 2. Sub-analyses of age and care seeking at hospitals or clinics in LMICs. Odds that an individual seeks care for diarrheal illness for a child under five years of age compared to an adult or individual over five years of age. Results from two sub-analyses are shown: 1) studies only among children or only among adults, and 2) studies with care seeking stratified for children and adults. p-values are shown for the univariate mixed-effects models; n.s. indicates not significant ($p > 0.05$).

Sub-analysis	Age category	Odds ratio	p-value
Observations among only children (N = 98) or only adults (N = 22)	Over five	1 [Reference]	
	Under five	1.06 (0.55 - 2.01)	n.s.

Observations with care seeking stratified by age (n = 8)	Over five	1 [Reference]	
	Under five	1.24 (0.29 - 5.36)	n.s.

559

560

561 **Table 3. Factors associated with variation in care seeking at hospitals or clinics in LMICs.** Odds
 562 that an individual sought care for diarrheal illness for severe diarrhea or cholera and gastroenteritis or
 563 other etiologies compared to general diarrhea, adjusting for timing of care seeking. p-values are shown
 564 for the multivariate mixed-effects model; n.s. indicates not significant ($p > 0.05$).

Variable	Category	Odds ratio	p-value
Diarrhea case definition	Diarrhea	1 [Reference]	
	Severe diarrhea or cholera	3.43 (1.71 - 6.88)	< 0.001
	Gastroenteritis or other etiologies	1.15 (0.46 - 2.90)	n.s.
Timing of care seeking	Care sought at any time	1 [Reference]	
	First source sought	0.44 (0.24 - 0.80)	< 0.01

565

566

567

568 Figure legends

569 **Figure 1. PRISMA flow diagram.** Diagram illustrates the search, screening, and inclusion
 570 process for the systematic review, including databases searched, reasons for exclusion, and
 571 final studies included in the meta-analyses.

572

573 **Figure 2. Proportion that sought care at a hospital or clinic in LMICs by observation**
 574 **characteristics.** Proportion of respondents that reported seeking care for diarrheal illness at a
 575 hospital or clinic grouped by **A)** whether the study surveyed participants about care seeking at
 576 any time during illness or their first source of care, **B)** survey respondent category, **C)** whether

577 or not the study described an outbreak, and **D**) whether the care seeking was for the respondent
578 themselves, their child, or other, which often included any household member. Each point
579 represents an observation (N = 145). Boxes represent the median and interquartile range (IQR)
580 of the proportion for each group. Lines extend from the top and bottom of box to the largest
581 proportion value no further than $1.5 * IQR$ from the box.

582

583 **Figure 3. Relationship between age and care seeking at hospitals or clinics in LMICs. A-**
584 **B)** Relationship between the proportion of respondents that reported seeking care for diarrheal
585 illness at a hospital or clinic (y-axis) and the proportion of the study population under five years
586 of age (x-axis), **A**) overall and **B**) stratified by diarrhea case definition. Size of points is
587 proportional to the number of survey respondents per observation (N = 134). Spearman's rho is
588 shown with 95% confidence interval (CI) in green. Linear relationships are shown with purple
589 lines. **C-D)** Proportion seeking care grouped by studies that were exclusively among children
590 under five (N = 98) or exclusively among individuals over five (N = 22), **C**) overall and **D**)
591 stratified by diarrhea case definition.

592

593 **Figure 4. Proportion that sought care by source in LMICs.** Proportion of respondents that
594 reported seeking care for diarrheal illness at all categories of care sources, including any care
595 outside of the home, hospitals or clinics including public facilities, private hospitals or clinics,
596 pharmacies or kiosks, health posts, community healthcare workers (CHW), traditional healers,
597 and friends or family, for **A**) all studies and **B**) the subset of studies that asked survey
598 respondents about the first source of care they sought. Each point represents an observation.

Identification

Studies from databases/registers (**n = 7069**)

Embase (n = 3561)
 PubMed (n = 2372)
 Web of Science (n = 1123)
 Global Index Medicus (n = 13)

References removed (**n = 1881**)

Duplicates identified manually (n = 73)
 Duplicates identified by Covidence (n = 1808)

Screening

Studies screened (**n = 5188**)

Studies not relevant (**n = 4466**)

Studies assessed for eligibility (**n = 722**)

Studies excluded (**n = 534**)

Qualitative (n = 46)
 Case control (n = 5)
 Duplicate data (n = 11)
 Convenience sample (n = 24)
 Special population (n = 5)
 Illness not diarrhea (n = 93)
 No care seeking data (n = 78)
 Not primary research (n = 57)
 Full text not available (n = 71)
 No care seeking questions (n = 98)
 Sampling done before 2000 (n = 19)
 Sampling method not reported (n = 17)
 Not English, French or Spanish (n = 10)

Included

Studies included in review (**n = 188**)

Non-overlapping studies included in analysis (**n = 166**)





