

Hemodynamic Mechanisms Underlying Orthostatic Hypotension in Stroke Survivors: A Cross-Sectional Study

Kazuaki OYAKE, PT, PhD,^{1,2} Ayumi MOCHIDA, PT, MSc,^{2,3} Masakiyo TERASHI, PT, BSc,²
Mahiro HASEGAWA, PT,² Akari SAITO, PT, BSc,² Kunitsugu KONDO, MD, PhD,²
Yohei OTAKA, MD, PhD,⁴ Kimito MOMOSE, PT, PhD¹

¹ Department of Physical Therapy, School of Health Sciences, Shinshu University, Nagano, Japan

² Department of Rehabilitation Medicine, Tokyo Bay Rehabilitation Hospital, Chiba, Japan

³ 'Present address' Department of Physical Therapy, School of Rehabilitation Sciences, Health Sciences University of Hokkaido, Japan

⁴ Department of Rehabilitation Medicine, Fujita Health University School of Medicine, Aichi, Japan

Corresponding author:

Kazuaki Oyake, PT, PhD

Department of Physical Therapy, School of Health Sciences, Shinshu University

3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan

Tel: +81-263-37-2408

Fax: +81-263-37-2408

E-mail: k_oyake@shinshu-u.ac.jp

Declaration of interest: None.

Highlights

- Orthostatic hypotension is associated with a higher risk of adverse outcomes.
- Mechanisms of orthostatic hypotension were examined in stroke survivors.
- Orthostatic hypotension may be associated with impaired vasoconstriction.
- Findings may aid in managing orthostatic hypotension during stroke rehabilitation.

Abstract

Background: Orthostatic hypotension is an important consideration for stroke survivors, given its association with adverse outcomes, such as cardiovascular diseases and falls. Understanding the hemodynamic mechanisms underlying orthostatic hypotension is essential for selecting the appropriate treatment based on individual hemodynamic patterns. However, the relative contribution of changes in cardiac output and total peripheral resistance to orthostatic hypotension remains unclear in stroke survivors.

Objective: To determine whether orthostatic hypotension is more strongly associated with a marked decrease in cardiac output or an impaired increase in total peripheral resistance among individuals with stroke.

Methods: This cross-sectional study included 23 individuals with stroke (13 males, mean [SD] age 63.7 [12.1] years, mean time since stroke 85.1 [34.1] days) recruited from an intensive inpatient rehabilitation ward between June 2022 and November 2023. Participants underwent a head-up tilt test to assess orthostatic hypotension and associated changes in cardiac and total peripheral resistance indices. The head-up tilt test protocol consisted of a 5-min period in the supine position followed by a 5-min period with a 70° head-up tilt. Orthostatic hypotension was defined as a reduction in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg during the test.

Results: Orthostatic hypotension was identified in five participants (22%). During the head-up tilt test, these individuals demonstrated a significantly greater increase in the cardiac index ($p = 0.023$) and decrease in the total peripheral resistance index ($p = 0.002$) than those without orthostatic hypotension.

Conclusions: Our results suggest that an impaired increase in total peripheral resistance upon standing mainly contributes to orthostatic hypotension in individuals with stroke. These findings advance the understanding of the hemodynamic mechanisms underlying orthostatic hypotension in the stroke population and may guide the implementation of targeted therapeutic strategies.

Keywords: cardiac output, cerebrovascular disease, rehabilitation, tilt-table test, total peripheral resistance

Introduction

Stroke is one of the primary causes of disability, with an increased risk of recurrence [1], cardiovascular diseases [2], dementia [3], and falls [4]. Orthostatic hypotension is also associated with a higher risk of these adverse outcomes [5-8], with a prevalence of 13.0–52.1% among stroke survivors [9-14]. Appropriate rehabilitation interventions to prevent or treat orthostatic hypotension may reduce future risks in this population.

Orthostatic hypotension occurs due to a marked decrease in cardiac output and/or an impaired increase in total peripheral resistance upon standing [15]. The marked decrease in cardiac output reflects an excessive reduction in venous return, while the impaired increase in total peripheral resistance is associated with inadequate baroreflex-mediated compensatory vasoconstriction [15]. Treatment strategies differ based on the underlying mechanism. For instance, compression garments or fludrocortisone are potential treatments for marked cardiac output decrease, while midodrine may be prescribed for an impaired increase in total peripheral resistance [15]. Thus, understanding the hemodynamic mechanisms underlying orthostatic hypotension is essential for selecting the appropriate treatment based on individual hemodynamic patterns. However, to the best of our knowledge, no studies have investigated the relative contributions of cardiac output and total peripheral resistance changes to orthostatic hypotension in stroke

survivors.

We aimed to examine whether orthostatic hypotension is more strongly associated with a marked decrease in cardiac output or an impaired increase in total peripheral resistance in individuals with stroke. In this population, orthostatic hypotension may result from complex interactions between various conditions such as baroreflex dysfunction, muscle weakness, hypovolemia, and comorbidities [5, 12, 13, 16, 17]. Older adults who share similar potential causes of orthostatic hypotension with stroke survivors have been reported to show a decrease in total peripheral resistance upon standing [18, 19]. Thus, we hypothesized that the orthostatic hypotension would be more strongly associated with an impaired increase in total peripheral resistance than with changes in cardiac output in individuals with stroke.

Material and Methods

Study design

This study used a cross-sectional, observational design. The ethics committee of Shinshu University approved the study protocol on May 12, 2022 (approval number: 5514). All participants provided written informed consent before enrollment in the study. This study adhered to the principles of the Declaration of Helsinki of 1964, as revised in 2013.

Participants

We recruited participants from an

intensive inpatient rehabilitation ward between June 2022 and November 2023. The inclusion criteria were as follows: (1) age 40–90 years; (2) stroke onset within 180 days; and (3) ability to cooperate with study procedures. The exclusion criteria were as follows: (1) limited range of motion and/or pain affecting the head-up tilt test; (2) moderate to severe dementia as defined by a Mini-Mental State Examination score of < 21 [20, 21]; (3) implanted metal devices such as a pacemaker and joint arthroplasty; (4) bilateral hemiparetic stroke; (5) unstable medical conditions, such as arrhythmias and uncontrolled diabetes mellitus; and (6) any comorbid neurological disorders. Demographic and clinical data, including age, sex, stroke type, and comorbidities, were extracted from medical records.

Data collection

All assessments were performed in a quiet room at a comfortable temperature between 16:00 and 18:00. Participants were instructed to refrain from eating and consuming caffeinated products for at least 2 h and to avoid vigorous exercise for at least 12 h before the assessments [22].

Head-up tilt test

The participants rested in a supine position on a motorized tilt table (SPR-7001, SAKAI Medical Co., Ltd., Tokyo, Japan) for 10 min to establish stable baseline hemodynamics [22]. We then measured hemodynamic variables during a 5-min supine period, followed by a 5-min passive head-up tilt to 70° [22]. The test was

promptly terminated if a participant experienced severe symptoms, such as presyncope.

Throughout the head-up tilt test, the participants were instructed to regulate their breathing at a rate of 0.25 Hz (approximately 15 breaths per minute) using a computer metronome with audio cues. This procedure minimized the influence of respiratory fluctuations on heart rate variability assessments [23].

At the end of the test, participants were instructed to report the severity of orthostatic symptoms, such as dizziness and palpitations, experienced during the 5-min upright posture using a visual analog scale. Participants marked a 100-mm horizontal line with a pen, where 0 mm indicated no symptoms and 100 mm represented syncope or presyncope [24, 25].

Assessments of hemodynamic and heart rate variability responses to head-up tilt

We measured hemodynamic variables using a non-invasive impedance cardiography device (Task Force Monitor 3040i, CNSystems Medizintechnik GmbH., Graz, Austria). A digital cuff was placed on the middle finger of the non-paretic hand for continuous arterial blood pressure measurement. The non-paretic arm was supported by a sling to maintain the finger cuff at the level of the heart. Beat-to-beat systolic blood pressure, diastolic blood pressure, and mean arterial pressure were obtained using the vascular unloading technique [26] and automatically corrected to oscillometric blood pressure values from the

paretic arm.

Heart rate, R-R intervals, stroke volume, cardiac output, and total peripheral resistance were also measured on a beat-to-beat basis. Heart rate and R-R intervals were derived from six-lead electrocardiography. Stroke volume was calculated using an improved transthoracic impedance cardiography method [27], with three short-band electrodes placed on the participants: one on the neck and two below the thorax. Stroke volume was calculated using the following formula:

$$\text{Stroke volume} = V_{th} \times \text{LVET} \times (dZ/dt)_{\max}/Z_0,$$

where V_{th} is the electrical participating thoracic volume, LVET is the left ventricular ejection time, $(dZ/dt)_{\max}$ is the maximal rate of impedance reduction for a given heartbeat, and Z_0 is the base impedance. Cardiac output was calculated as the product of the heart rate and stroke volume. Total peripheral resistance was derived using Ohm's law:

$$\text{Total peripheral resistance} = (\text{mean arterial pressure}/\text{cardiac output}) \times 80.$$

We normalized stroke volume, cardiac output, and total peripheral resistance to the estimated body surface area to derive the stroke index, cardiac index, and total peripheral resistance index (TPRI), respectively. We excluded data acquired during postural changes to avoid motion artifacts. Data on systolic and diastolic blood pressure, mean arterial pressure, heart rate, stroke index, cardiac index, and TPRI were averaged every minute. Supine values were defined as the average values obtained from the final minute of the supine period.

Orthostatic hypotension was defined as a reduction in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg during the test [28, 29].

Heart rate variability analysis was performed to assess cardiac autonomic function. Heart rate variability parameters for the supine position were determined from the R-R intervals recorded during the 5-min supine period, while those for the upright position were derived from the 5-min upright period. We applied a threshold-based artifact correction algorithm with a medium filter to remove ectopic beats and artifacts [25, 30]. This filter identified R-R intervals that deviated by more than 0.25 s from the local average. The identified artifacts were replaced with interpolated values using cubic spline interpolation. These calculations were performed using the Kubios HRV standard software version 3.5 (Kubios Oy, Kuopio, Finland).

We calculated the frequency-domain variables of heart rate variability using fast Fourier transform analysis, which used a Welch periodogram method with a window width of 256 s and 50% window overlap. Low-frequency (0.04–0.15 Hz) and high-frequency (0.15–0.40 Hz) spectral components were collected as absolute values of power. The low-frequency component of heart rate variability represents a combination of sympathetic and parasympathetic activity and baroreflex function, whereas the high-frequency component of heart rate variability primarily reflects vagal tone [23, 31]. Furthermore, the low-to-high frequency ratio of heart rate

variability was used as an indicator of the sympathetic-parasympathetic balance, with a higher ratio reflecting greater sympathetic activity [23, 31].

Assessment of body composition

We measured body composition using a multi-frequency bioelectrical impedance analyzer (InBody S10; InBody Co., Ltd., Seoul, South Korea) with the participants in a supine position. Eight electrodes were attached to the thumbs, middle fingers, and ankles of both sides.

We collected data on the skeletal muscle mass index, total body water, and fat-free mass. The skeletal muscle mass index was calculated as the sum of the skeletal muscle mass in the upper and lower extremities divided by the height squared. We defined low muscle mass as a skeletal muscle mass index of < 7.0 kg/m² for male individuals and < 5.7 kg/m² for female individuals [32]. In addition, we calculated the ratio of total-body water to fat-free mass, as a lower ratio indicates hypovolemia induced by bed rest [33].

Assessment of functional outcomes

We assessed motor function and independence in performing activities of daily living as functional outcomes. To evaluate motor impairments in the paretic upper and lower extremities, we used the motor items of the Stroke Impairment Assessment Set [34]. The degree of independence in activities of daily living was measured using the Functional Independence Measure [35].

Statistical analyses

We calculated the required sample size using G*power version 3.1.9.2 (Heinrich-Heine-Universität, Düsseldorf, Germany). Based on previous studies comparing hemodynamic responses between middle-aged and older adults with and without orthostatic hypotension [18, 19], we assumed an effect size of 0.90 with an alpha level of 0.05 and a power of 0.80 for independent t-tests. Given the expected prevalence of orthostatic hypotension of 20% in stroke survivors [14], the required sample size was calculated to be 50 participants (10 with orthostatic hypotension and 40 without orthostatic hypotension).

The normality of distribution for all continuous variables was assessed using the Shapiro–Wilk test. Heart rate variability parameters showed significant deviation from normality ($p < 0.05$) and were logarithmically transformed prior to analysis. Characteristics of participants were compared between two groups (with and without orthostatic hypotension) using the unpaired t-test for continuous variables, the Mann–Whitney U test for ordinal variables, and Fisher’s exact test for dichotomous variables. To test our hypothesis, hemodynamic responses to the head-up tilt test were compared between the groups using two-way repeated-measures analysis of variance (ANOVA) with the group as the between-subject factor and time (supine and 1–5 minutes post-standing) as the within-subject factor. In addition, comparisons of heart rate variability responses to the head-up

tilt test between the groups were analyzed using a two-way repeated-measures ANOVA with the group as the between-subject factor and position (supine and upright positions) as the within-subject factor.

Based on our hypothesis that orthostatic hypotension in stroke survivors would be more strongly associated with an impaired increase in total peripheral resistance, we further investigated factors associated with orthostatic TPRI changes. We examined the associations between the orthostatic change in TPRI and the characteristics of participants, hemodynamic variables and heart rate variability parameters in the supine position, and the orthostatic changes in hemodynamic variables and heart rate variability parameters. These associations were analyzed using Pearson's product-moment correlation coefficient, the unpaired t-test, and Spearman's rank correlation coefficient based on variable types. Orthostatic changes in hemodynamic variables were calculated by subtracting the measurements in the supine position from those at the lowest TPRI during the upright period. Statistical analyses were performed using GraphPad Prism version 9.00 for Windows (GraphPad Software, San Diego, California, United States). Statistical significance was set at a $p < 0.05$.

Results

Participants

Fig. 1 presents the flow of participant enrollment. Of the 177 individuals with stroke assessed for eligibility, 154 were excluded, primarily because of moderate to severe dementia ($n = 97$). Owing to an unexpected equipment malfunction that could not be repaired within the study timeframe, only 23 participants were included in the analysis instead of the planned 50 participants. Table 1 presents the characteristics of the participants. All participants were in the subacute phase of stroke recovery [36] and participated in approximately 180 min of intensive daily rehabilitation.




Figure 1. Flow diagram of participant enrollment.

Table 1. Characteristics of participants.

Variable	Overall (<i>n</i> = 23)	With OH (<i>n</i> = 5)	Without OH (<i>n</i> = 18)	p-value
Age, years	63.7 (12.1)	64.6 (17.0)	63.5 (11.0)	0.862
Sex, male/female	13/10	4/1	9/9	0.339
Height, m	1.63 (0.08)	1.66 (0.07)	1.62 (0.08)	0.447
Weight, kg	66.6 (13.9)	61.2 (10.7)	67.3 (14.7)	0.405
Body mass index, kg/m ²	24.8 (4.9)	22.2 (2.8)	25.5 (5.1)	0.189
Stroke type, ischemia/hemorrhage	15/8	3/2	12/6	0.999
Side of motor paresis, right/left	8/15	1/4	7/11	0.621
Time since stroke, days	85.1 (34.1)	82.0 (33.8)	85.9 (35.2)	0.825
Antihypertensive medications				
Calcium channel blockers	13	2	11	0.618
Angiotensin II receptor blockers	4	1	3	0.999
Diuretics	4	0	4	0.539
Beta-blocker	2	1	1	0.395
Comorbidities				
Hypertension	19	3	16	0.194
Diabetes mellitus	3	0	4	0.539
Skeletal muscle mass index, kg/m ²	7.58 (1.07)	7.42 (1.37)	7.62 (1.02)	0.724
Low skeletal muscle mass, male/female	3/0	2	1	0.107
Ratio of total body-water to fat-free mass	0.741 (0.003)	0.712 (0.003)	0.746 (0.025)	0.690
Stroke Impairment Assessment Set motor score, points	16 (10–20)	10 (7–20)	17 (13–20)	0.599
Functional Independence Measure motor score, points	68 (54–78)	52 (49–73)	69 (58–79)	0.421
Functional Independence Measure cognitive score, points	31 (29–34)	31 (31–33)	31 (29–34)	0.966

Values are presented as the mean (SD), median (1st quartile–3rd quartile), or number. P-values indicate significant differences between the groups with and without OH.

OH, orthostatic hypotension

All the participants completed the head-up tilt test without any adverse orthostatic symptoms. The median (1st quartile–3rd quartile) visual analog scale score was 0 (0–8) mm. However, five participants (22%) met the criteria for orthostatic hypotension in this study. Two participants met both the systolic and diastolic blood pressure criteria, while three exhibited only diastolic orthostatic hypotension. The characteristics of participants did not differ significantly between the groups with and without orthostatic hypotension ($p > 0.05$, Table 1).

Hemodynamic and heart rate variability responses to the head-up tilt test

Fig. 2 shows the hemodynamic and heart rate variability responses to the head-up tilt test in both groups with and without orthostatic hypotension. Table 2 summarizes the changes in hemodynamic variables from supine to each minute of the 5-min upright period, stratified by orthostatic hypotension status.

The two-way repeated-measures ANOVA revealed significant interactions between group and time for systolic blood pressure ($F_{(5,105)} = 3.27$, $p = 0.009$; Fig. 2A), diastolic blood pressure ($F_{(5,105)} = 11.87$, $p < 0.001$; Fig. 2B), and mean arterial pressure ($F_{(5,105)} = 8.50$, $p < 0.001$; Fig. 2C). After standing, the group without orthostatic

hypotension showed increasing trends in these variables, whereas the group with orthostatic hypotension demonstrated decreasing trends (Table 2).

For heart rate, although neither the group \times time interaction ($F_{(5,105)} = 1.19$, $p = 0.318$) nor the main effect of the group ($F_{(1,21)} = 1.81$, $p = 0.194$) was significant, there was a significant main effect of time ($F_{(5,105)} = 29.70$, $p < 0.001$; Fig. 2D), with both groups showing increased values after standing (Table 2). The stroke index showed neither significant group \times time interaction ($F_{(5,105)} = 1.74$, $p = 0.132$) nor a main effect of the group ($F_{(1,21)} = 1.69$, $p = 0.208$; Fig. 2E).

Analysis of the cardiac index (Fig. 2F) and TPRI (Fig. 2G) revealed significant group \times time interactions (cardiac index: $F_{(5,105)} = 2.75$, $p = 0.023$; TPRI: $F_{(5,105)} = 4.19$, $p = 0.002$) and main effects of the group (cardiac index: $F_{(1,21)} = 5.22$, $p = 0.033$; TPRI: $F_{(1,21)} = 5.01$, $p = 0.036$). These results indicate that compared to the group without orthostatic hypotension, the group with orthostatic hypotension demonstrated significantly greater increases in the cardiac index and more pronounced decreases in the TPRI after standing (Table 2). In all participants with orthostatic hypotension, a higher cardiac index and a lower TPRI remained throughout the upright period than in the supine period.




Figure 2. Comparisons of the hemodynamic and heart rate variability responses to the head-up tilt test between the groups with and without orthostatic hypotension.

Hemodynamic variables include the (A) systolic blood pressure, (B) diastolic blood pressure, (C) mean arterial pressure, (D) heart rate, (E) stroke index, (F) cardiac index, and (G) TPRI. Heart rate variability parameters are the (H) lnLF, (I) lnHF, and (J) ln(LF/HF) ratio. The white squares and the blue diamonds represent the mean values in the groups with and without orthostatic hypotension, respectively. The error bars indicate the SD. Data at 0 on the x-axis corresponds to data at the final 1 min of the supine period. lnHF, logarithmic high-frequency component of heart rate variability; lnLF, logarithmic low-frequency component of heart rate variability; ln(LF/HF) ratio, logarithmic low-to-high frequency ratio of heart rate variability; TPRI, total peripheral resistance index.

For the logarithmic low-frequency component (lnLF), the logarithmic high-frequency component (lnHF), and the logarithmic low-to-high frequency ratio (ln(LF/HF) ratio), there were neither significant group \times time interactions (lnLF: $F_{(1,21)} = 0.78$, $p = 0.388$; lnHF: $F_{(1,21)} = 0.09$, $p = 0.767$; ln(LF/HF) ratio: $F_{(1,21)} = 0.37$, $p = 0.551$) nor main effects of group (lnLF: $F_{(1,21)} = 0.45$, $p = 0.511$; lnHF: $F_{(1,21)} = 1.94$, $p = 0.179$; ln(LF/HF) ratio: $F_{(1,21)} = 1.35$, $p = 0.259$).

Factors associated with an impaired orthostatic increase in TPRI

No significant associations were found between the orthostatic change in TPRI and the characteristics of participants ($p > 0.05$; Table 3). Correlations of orthostatic change in TPRI with hemodynamic variables and heart rate variability parameters during the head-up tilt test are shown in Fig. 3 and Table 4. We found that a greater orthostatic decrease in TPRI was associated with lower supine values of the stroke index ($r = 0.550$, 95% confidence interval [CI] = 0.178–0.784, $p =$

0.006; Fig. 3A), the cardiac index ($r = 0.570$, 95% CI = 0.208–0.796, $p = 0.004$; Fig. 3B), and a higher supine TPRI ($r = -0.613$, 95% CI = -0.818 to -0.269, $p = 0.002$; Fig. 3C). No other significant correlations were found between the orthostatic change in TPRI and supine hemodynamic variables or heart rate variability parameters ($p > 0.05$; Table 4).

Regarding orthostatic changes, a greater decrease in TPRI was significantly correlated with larger reductions in diastolic blood pressure ($r = 0.506$, 95% CI = 0.119–0.760, $p = 0.014$; Fig. 3D) and mean arterial pressure ($r = 0.532$, 95% CI = 0.154–0.775, $p = 0.009$; Fig. 3E). Additionally, a greater TPRI decrease was associated with larger increases in the stroke index ($r = -0.774$, 95% CI = -0.899 to -0.530, $p < 0.001$; Fig. 3F), cardiac index ($r = -0.824$, 95% CI = -0.923 to -0.624, $p < 0.001$; Fig. 3G), and ln(LF/HF) ratio ($r = -0.531$, 95% CI = -0.774 to -0.152, $p = 0.009$; Fig. 3H). No other significant correlations were found between the orthostatic change in TPRI and orthostatic changes in the remaining parameters ($p > 0.05$; Table 4).

Table 3. Associations between orthostatic change in total peripheral resistance index and the characteristics of participants.

Variable	Orthostatic change in TPRI	
	Statistics	p-value
Age	r = -0.089	0.687
Sex	t = 1.631	0.118
Height	r = -0.224	0.305
Weight	r = -0.214	0.328
Body mass index	r = -0.131	0.552
Stroke type	t = -0.647	0.524
Side of motor paresis	t = 0.475	0.639
Time since stroke	r = -0.230	0.291
Antihypertensive medications		
Calcium channel blockers	t = 0.406	0.689
Angiotensin II receptor blockers	t = 1.452	0.161
Diuretics	t = 1.589	0.127
Beta-blocker	t = 1.950	0.065
Comorbidities		
Hypertension	t = -1.331	0.198
Diabetes mellitus	t = 1.319	0.201
Skeletal muscle mass index	r = -0.277	0.202
Low muscle mass	t = 1.002	0.328
Ratio of total body-water to fat-free mass	r = -0.274	0.206
Stroke Impairment Assessment Set motor score	rho = -0.227	0.297
Functional Independence Measure motor score	rho = -0.071	0.748
Functional Independence Measure cognitive score	rho = -0.306	0.156

TPRI, total peripheral resistance index.




Figure 3. Correlations of orthostatic change in the total peripheral resistance index with hemodynamic variables and heart rate variability parameters during the head-up tilt test.

Scatter plots showing correlations between orthostatic change in TPRI and (A) supine stroke index, (B) supine cardiac index, (C) supine TPRI, (D) orthostatic change in diastolic blood pressure, (E) orthostatic change in mean arterial pressure, (F) orthostatic change in the stroke index, (G) orthostatic change in the cardiac index, and (H) orthostatic change in the ln(LF/HF) ratio. The red and blue diamonds represent participants with and without orthostatic hypotension, respectively. The solid black line represents the regression line, while the shaded area indicates the 95% confidence interval of the regression line. Vertical dashed lines in panels D-H indicate no change from supine to the upright position. Horizontal dashed lines indicate no change in TPRI from supine to the upright position.

ln(LF/HF) ratio, logarithmic low-to-high frequency ratio of heart rate variability; TPRI, total peripheral resistance index.

Table 4. Correlations of orthostatic change in total peripheral resistance index with hemodynamic variables and heart rate variability parameters during the head-up tilt test.

Variable	Orthostatic change in TPRI		
	r	95% CI	p-value
Supine systolic blood pressure	-0.086	-0.481 to 0.338	0.683
Supine diastolic blood pressure	-0.090	-0.484 to 0.335	0.683
Supine mean arterial pressure	-0.120	-0.507 to 0.307	0.584
Supine heart rate	0.147	-0.283 to 0.527	0.504
Supine stroke index	0.550	0.178–0.784	0.006
Supine cardiac index	0.571	0.208–0.796	0.004
Supine TPRI	-0.613	-0.818 to -0.269	0.002
Supine lnLF	0.249	-0.182 to 0.600	0.252
Supine lnHF	0.155	-0.274 to 0.533	0.479
Supine ln(LF/HF) ratio	0.114	-0.313 to 0.502	0.606
Orthostatic change in systolic blood pressure	0.343	-0.081 to 0.662	0.109
Orthostatic change in diastolic blood pressure	0.506	0.119–0.760	0.014
Orthostatic change in mean arterial pressure	0.532	0.154–0.775	0.009
Orthostatic change in heart rate	0.077	-0.346 to 0.474	0.767
Orthostatic change in stroke index	-0.774	-0.899 to -0.530	<0.001
Orthostatic change in cardiac index	-0.824	-0.923 to -0.624	<0.001
Orthostatic change in lnLF	-0.272	-0.615 to 0.158	0.210
Orthostatic change in lnHF	0.223	-0.208 to 0.582	0.306
Orthostatic change in ln(LF/HF) ratio	-0.531	-0.774 to -0.152	0.009

Values marked in bold indicate a significant correlation.

CI, confidence interval; lnHF, logarithmic high-frequency component of heart rate variability; ln(LF/HF) ratio, logarithmic low-to-high frequency ratio of heart rate variability; lnLF, logarithmic low-frequency component of heart rate variability; r, Pearson's product-moment correlation coefficient; TPRI, total peripheral resistance index.

Discussion

To the best of our knowledge, this study is the first to examine the hemodynamic mechanisms underlying orthostatic hypotension in stroke survivors. Compared to the group without orthostatic hypotension, the group with orthostatic hypotension demonstrated greater increases in the cardiac index and larger decreases in TPRI during the head-up tilt test. These results support our hypothesis that orthostatic hypotension may be more strongly associated with an impaired increase in total peripheral resistance than with changes in cardiac output in individuals with stroke. Our findings may provide clinicians with valuable insights into managing orthostatic hypotension during stroke rehabilitation.

The prevalence of orthostatic hypotension in this study population (22%) was similar to data from our previous study of inpatients with subacute stroke (24%) [14]. In addition, all participants remained asymptomatic during the head-tilt test, regardless of the presence or absence of orthostatic hypotension. A study also reported that many individuals with orthostatic hypotension were asymptomatic despite substantial reductions in systolic blood pressure and low upright systolic blood pressure [37]. These findings highlight the importance of measuring orthostatic changes in blood pressure rather than depending on symptoms alone to indicate orthostatic hypotension. Although habituation and cerebral autoregulation are possible explanations for the absence of orthostatic

symptoms, further research is needed to elucidate the mechanisms underlying this impaired symptom recognition [37].

The transition from the supine to the upright position causes blood pooling in the lower extremities and splanchnic veins, which may lead to reduced venous return, stroke volume, and cardiac output. In healthy adults, baroreflex-mediated compensation increases heart rate, cardiac contractility, and total peripheral resistance, ultimately raising the mean arterial pressure [29, 38]. Studies have reported statistically significant increases in blood pressure variables, heart rate, and total peripheral resistance, along with decreases in stroke volume and cardiac output, during this postural change in healthy individuals [39-41]. In contrast, our participants with orthostatic hypotension showed sustained higher cardiac index and lower TPRI values relative to the supine values throughout the upright period. These results suggest that inadequate baroreflex-mediated vasoconstriction mainly contributes to orthostatic hypotension in individuals with stroke. Additionally, we found the associations of orthostatic decrease in TPRI with orthostatic increases in the stroke index, cardiac index, and $\ln(\text{LF}/\text{HF})$ ratio as well as with orthostatic reductions in diastolic blood pressure and mean arterial pressure. These results suggest a compensatory mechanism in which the reduction in mean arterial pressure caused by decreased total peripheral resistance triggers enhanced cardiac sympathetic activity, leading to augmented stroke volume and cardiac output. Previous studies of middle-

aged and older adults have also reported that an impaired orthostatic increase in total peripheral resistance is primarily related to orthostatic hypotension [18, 19, 42].

Prior studies have established that impaired orthostatic increases in total peripheral resistance are linked to central neurodegenerative and peripheral autonomic disorders, including Parkinson's disease and peripheral neuropathies [15, 43]. However, none of our participants had these diagnosed conditions. While diabetes mellitus is known to cause peripheral autonomic neuropathy, we found no significant association between diabetes mellitus and orthostatic decreases in TPRI. Arterial stiffness emerges as another potential mechanism underlying impaired orthostatic increases in total peripheral resistance. Research has demonstrated that arterial stiffness is associated with a smaller increase or a decrease in the mean arterial pressure upon standing [38] and orthostatic hypotension [44]. A previous head-up tilt study revealed that individuals exhibiting smaller TPRI increases and cardiac index decreases showed higher pulse wave velocity, indicating increased arterial stiffness [45, 46]. These individuals also demonstrated lower supine stroke and cardiac indices, higher supine TPRI, and a greater orthostatic increase in the $\ln(LF/HF)$ ratio; however, the causality remained undetermined [46]. Although we did not include pulse wave velocity measurements, the results of correlation analyses suggest that the observed orthostatic TPRI decrease is associated with arterial stiffness. Given that both stretching and aerobic exercises

effectively reduce arterial stiffness [47, 48], future investigations should examine the effect of these exercise interventions on orthostatic changes in total peripheral resistance. Such research could clarify the mechanisms underlying impaired orthostatic total peripheral resistance increases and inform the development of targeted therapeutic strategies for orthostatic hypotension in stroke survivors.

Study limitations

This study had some limitations. First, owing to an unexpected equipment malfunction, we were only able to include 23 participants instead of the planned 50 participants, which may have limited our statistical power. In addition, although orthostatic hypotension in stroke survivors is potentially associated with older age, female sex, hypertension, diabetes mellitus, low skeletal muscle mass, hypovolemia, severe motor paresis, and increased dependence in activities of daily living [12, 17], we did not observe associations of these characteristics with orthostatic hypotension or TPRI changes. This lack of association may be attributed to our limited sample size. Nevertheless, our results provide valuable insights into selecting the appropriate treatment for orthostatic hypotension based on individual hemodynamic patterns. Second, all participants were subacute stroke survivors receiving daily intensive rehabilitation, which may limit the generalizability of our findings to individuals with acute stroke who present with unstable general conditions or those in the chronic phase of stroke recovery

after prolonged bed rest. Third, owing to the cross-sectional observational design of this study, we could not examine whether improvement in impaired orthostatic total peripheral resistance increases ameliorates orthostatic hypotension. Longitudinal studies are needed to investigate the temporal association between orthostatic hypotension and the orthostatic change in total peripheral resistance. Finally, this study did not include a control group. Further investigations should include an age- and sex-matched healthy control group for comparison with the stroke cohort. This approach would provide deeper insights into the pathogenesis of orthostatic hypotension and the impaired total peripheral resistance increase observed in individuals with stroke.

Conclusions

In individuals with stroke, orthostatic hypotension was characterized by an impaired increase in total peripheral resistance rather than a marked decrease in cardiac output. These findings suggest that inadequate baroreflex-mediated vasoconstriction is the primary mechanism underlying orthostatic hypotension in this population. Our results enhance the understanding of hemodynamic responses to postural change after stroke and provide guidance for selecting appropriate therapeutic strategies.

Acknowledgments

We would like to thank Editage (www.editage.com) for English language editing.

Funding

This work was supported by a grant from JSPS KAKENHI Grant Number JP21K17489 awarded to K.O. The funding source had no involvement in the study design; collection, analysis and interpretation of data; writing of the report; and the decision to submit the article for publication.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used Claude 3.5 Sonnet (Anthropic, San Francisco, CA, USA) for generating preliminary drafts and English editing assistance. After using this tool/service, the authors reviewed and edited the content as needed and took full responsibility for the content of the publication.

Data availability statement

The data underlying this article will be shared on reasonable request to the corresponding author.

References

- [1] Lin B, Zhang Z, Mei Y, Wang C, Xu H, Liu L, et al. Cumulative risk of stroke recurrence over the last 10 years: a systematic review and meta-analysis. *Neurol Sci.* 2021;42:61-71, <https://doi.org/10.1007/s10072-020-04797-5>.
- [2] Sposato LA, Lam M, Allen B, Richard L, Shariff SZ, Saposnik G. First-ever ischemic stroke and increased risk of incident heart disease in older adults. *Neurology.* 2020;94:e1559-e70, <https://doi.org/10.1212/wnl.00000000000009234>.
- [3] Kuźma E, Lourida I, Moore SF, Levine DA, Ukoumunne OC, Llewellyn DJ. Stroke and dementia risk: a systematic review and meta-analysis. *Alzheimers Dement.* 2018;14:1416-26, <https://doi.org/10.1016/j.jalz.2018.06.3061>.
- [4] Denissen S, Staring W, Kunkel D, Pickering RM, Lennon S, Geurts AC, et al. Interventions for preventing falls in people after stroke. *Cochrane Database Syst Rev.* 2019;10:CD008728, <https://doi.org/10.1002/14651858.CD008728.pub3>.
- [5] Mehta T, McClure LA, White CL, Taylor A, Benavente OR, Lakshminarayan K. Effect of postural hypotension on recurrent stroke: secondary prevention of small subcortical strokes (SPS3) study. *J Stroke Cerebrovasc Dis.* 2019;28:2124-31, <https://doi.org/10.1016/j.jstrokecerebrovasdis.2019.04.009>.
- [6] Ricci F, Fedorowski A, Radico F, Romanello M, Tatasciore A, Di Nicola M, et al. Cardiovascular morbidity and mortality related to orthostatic hypotension: a meta-analysis of prospective observational studies. *Eur Heart J.* 2015;36:1609-17, <https://doi.org/10.1093/eurheartj/ehv093>.
- [7] Xia X, Qiu C, Rizzuto D, Fratiglioni L, Dai L, Laukka EJ, et al. Role of orthostatic hypotension in the development of dementia in people with and without cardiovascular disease. 2023;80:1474-83, <https://doi.org/10.1161/HYPERTENSIONA.123.21210>.
- [8] Mol A, Bui Hoang PTS, Sharmin S, Reijnierse EM, van Wezel RJA, Meskers CGM, et al. Orthostatic hypotension and falls in older adults: a systematic review and meta-analysis. *J Am Med Dir Assoc.* 2019;20:589-597.e5, <https://doi.org/10.1016/j.jamda.2018.11.003>.
- [9] Panayiotou B, Reid J, Fotherby M, Crome P. Orthostatic haemodynamic responses in acute stroke. *Postgrad Med J.* 1999;75:213-8, <https://doi.org/10.1136/pgmj.75.882.213>.
- [10] Aries MJH, Bakker DC, Stewart RE, De Keyser J, Elting JWJ, Thien T, et al. Exaggerated postural blood pressure rise is related to a favorable outcome in patients with acute ischemic stroke. *Stroke.* 2012;43:92-6, <https://doi.org/10.1161/STROKEAHA.111.632349>.
- [11] Treger I, Shafir O, Keren O, Ring H. Cerebral blood flow velocity during postural changes on tilt table in stroke patients. *Eura Medicophys.* 2005;41:293-6.
- [12] Kong KH, Chuo AM. Incidence and outcome of orthostatic hypotension in stroke patients undergoing rehabilitation. *Arch Phys*

- Med Rehabil. 2003;84:559-62, <https://doi.org/10.1053/apmr.2003.50040>.
- [13] Phipps MS, Schmid AA, Kapoor JR, Peixoto AJ, Williams LS, Bravata DM. Orthostatic hypotension among outpatients with ischemic stroke. *J Neurol Sci.* 2012;314:62-5, <https://doi.org/10.1016/j.jns.2011.10.031>.
- [14] Oyake K, Tateishi T, Murayama J, Shijima N, Sekizuka T, Otaka Y, et al. Blood pressure reduction cut-points for orthostatic hypotension in stroke survivors using a sit-up test: a multicentre cross-sectional study. *J Hypertens.* 2023;41:83-9, <https://doi.org/10.1097/HJH.0000000000003305>.
- [15] Deegan BM, O'Connor M, Donnelly T, Carew S, Costelloe A, Sheehy T, et al. Orthostatic hypotension: a new classification system. *Europace.* 2007;9:937-41, Epub 2007/08/28. <https://doi.org/10.1093/europace/eum177>.
- [16] Tang A, Eng JJ, Krassioukov A. Application of the Sit-Up Test for orthostatic hypotension in individuals with stroke. *Auton Neurosci.* 2012;168:82-7, <https://doi.org/10.1016/j.autneu.2012.02.002>.
- [17] Ögren J, Mooe T, Irewall AL. Orthostatic hypotension in stroke/TIA patients: association with new events and the effect of the NAILED intervention. *PLOS ONE.* 2024;19(2):e0298435. <https://doi.org/10.1371/journal.pone.0298435>.
- [18] van Wijnen VK, Hove DT, Finucane C, Wieling W, van Roon AM, Ter Maaten JC, et al. Hemodynamic mechanisms underlying initial orthostatic hypotension, delayed recovery and orthostatic hypotension. *J Am Med Dir Assoc.* 2018;19:786-92, <https://doi.org/10.1016/j.jamda.2018.05.031>.
- [19] Hui G, Xiahuan C, Yanjun W, Wenyi L, Meilin L. Influencing factors and hemodynamic study of initial and sustained orthostatic hypotension in middle-aged and elderly patients. *J Clin Hypertens (Greenwich).* 2022;24:1491-7, <https://doi.org/10.1111/jch.14588>.
- [20] Folstein MF, Folstein SE, McHugh PR. 'Mini-mental state'. A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189-98, [https://doi.org/10.1016/0022-3956\(75\)90026-6](https://doi.org/10.1016/0022-3956(75)90026-6).
- [21] Perneczky R, Wagenpfeil S, Komossa K, Grimmer T, Diehl J, Kurz A. Mapping scores onto stages: Mini-Mental State Examination and clinical dementia rating. *Am J Geriatr Psychiatry.* 2006;14:139-44, <https://doi.org/10.1097/01.JGP.0000192478.82189.a8>.
- [22] Cheshire WP, Jr., Goldstein DS. Autonomic uprising: the tilt table test in autonomic medicine. *Clin Auton Res.* 2019;29:215-30, <https://doi.org/10.1007/s10286-019-00598-9>.
- [23] Laborde S, Mosley E, Thayer JF. Heart rate variability and cardiac vagal tone in psychophysiological research - recommendations for experiment planning, data analysis, and data reporting. *Front Psychol.* 2017;8:213, <https://doi.org/10.3389/fpsyg.2017.00213>.
- [24] Figueroa JJ, Singer W, Sandroni P, Sletten DM, Gehrking TL, Gehrking JA, et al. Effects of patient-controlled abdominal

- compression on standing systolic blood pressure in adults with orthostatic hypotension. *Arch Phys Med Rehabil.* 2015;96:505-10, <https://doi.org/10.1016/j.apmr.2014.10.012>.
- [25] Oyake K, Katai M, Yoneyama A, Ikegawa H, Kani S, Momose K. Comparisons of heart rate variability responses to head-up tilt with and without abdominal and lower-extremity compression in healthy young individuals: a randomized crossover study. *Front Physiol.* 2023;14:1269079, <https://doi.org/10.3389/fphys.2023.1269079>.
- [26] Jeleazcov C, Krajcinovic L, Münster T, Birkholz T, Fried R, Schüttler J, et al. Precision and accuracy of a new device (CNAPTM) for continuous non-invasive arterial pressure monitoring: assessment during general anaesthesia. *Br J Anaesth.* 2010;105:264-72, <https://doi.org/10.1093/bja/aeq143>.
- [27] Fortin J, Habenbacher W, Heller A, Hacker A, Grüllenberger R, Innerhofer J, et al. Non-invasive beat-to-beat cardiac output monitoring by an improved method of transthoracic bioimpedance measurement. *Comput Biol Med.* 2006;36:1185-203, <https://doi.org/10.1016/j.compbiomed.2005.06.001>.
- [28] Freeman R, Wieling W, Axelrod FB, Benditt DG, Benarroch E, Biaggioni I, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auton Res.* 2011;21:69-72, <https://doi.org/10.1007/s10286-011-0119-5>.
- [29] Wieling W, Kaufmann H, Claydon VE, van Wijnen VK, Harms MPM, Juraschek SP, et al. Diagnosis and treatment of orthostatic hypotension. *Lancet Neurol.* 2022;21:735-46, [https://doi.org/10.1016/s1474-4422\(22\)00169-7](https://doi.org/10.1016/s1474-4422(22)00169-7).
- [30] Baker J, Racosta JM, Balint B, Kimpinski K. Utility of time and frequency domain parameters of heart rate variability in the context of autonomic disorders characterized by orthostatic dysfunction. *J Clin Neurophysiol.* 2018;35:123-9, <https://doi.org/10.1097/wnp.0000000000000452>.
- [31] Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Front Public Health.* 2017;5:258, <https://doi.org/10.3389/fpubh.2017.00258>.
- [32] Chen LK, Woo J, Assantachai P, Auyeung TW, Chou MY, Iijima K, et al. Asian Working Group for Sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc.* 2020;21:300-7.e2, <https://doi.org/10.1016/j.jamda.2019.12.012>.
- [33] Ishizaki Y, Fukuoka H, Ishizaki T, Kino M, Higashino H, Ueda N, et al. Measurement of inferior vena cava diameter for evaluation of venous return in subjects on day 10 of a bed-rest experiment. *J Appl Physiol (1985).* 2004;96:2179-86, <https://doi.org/10.1152/japplphysiol.01097.2003>.
- [34] Chino N, Sonoda S, Domen K, Saitoh E, Kimura A. Stroke Impairment Assessment Set (SIAS). A new evaluation instrument for stroke patients. *Jpn J Rehabil Med.* 1994;31:119-25, <https://doi.org/10.2490/jjrm1963.31.119>.

- [35] Granger CV, Hamilton BB, Linacre JM, Heinemann AW, Wright BD. Performance profiles of the functional independence measure. *Am J Phys Med Rehabil.* 1993;72:84-9, <https://doi.org/10.1097/00002060-199304000-00005>.
- [36] Bernhardt J, Hayward KS, Kwakkel G, Ward NS, Wolf SL, Borschmann K, et al. Agreed definitions and a shared vision for new standards in stroke recovery research: the Stroke Recovery and Rehabilitation Roundtable taskforce. *Int J Stroke.* 2017;12:444-50, <https://doi.org/10.1177/1747493017711816>.
- [37] Freeman R, Illigens BMW, Lapusca R, Campagnolo M, Abuzinadah AR, Bonyhay I, et al. Symptom recognition is impaired in patients with orthostatic hypotension. *Hypertension.* 2020;75:1325-32, <https://doi.org/10.1161/HYPERTENSIONAHA.119.13619>.
- [38] Torjesen A, Cooper LL, Rong J, Larson MG, Hamburg NM, Levy D, et al. Relations of arterial stiffness with postural change in mean arterial pressure in middle-aged adults: the Framingham Heart Study. *Hypertension.* 2017;69:685-90, <https://doi.org/10.1161/HYPERTENSIONAHA.116.08116>
- [39] Critchley LA, Conway F, Anderson PJ, Tomlinson B, Critchley JA. Non-invasive continuous arterial pressure, heart rate and stroke volume measurements during graded head-up tilt in normal man. *Clin Auton Res.* 1997;7:97-101, <https://doi.org/10.1007/BF02267754>.
- [40] O'Leary DD, Kimmerly DS, Cechetto AD, Shoemaker JK. Differential effect of head-up tilt on cardiovagal and sympathetic baroreflex sensitivity in humans. *Exp Physiol.* 2003;88:769-74, <https://doi.org/10.1113/eph8802632>.
- [41] Oyake K, Murayama J, Tateishi T, Mochida A, Matsumoto M, Tsujikawa M, et al. Comparison of the sit-up test and head-up tilt test for assessing blood pressure and hemodynamic responses in healthy young individuals. *Blood Press Monit.* 2022;27:79-86, <https://doi.org/10.1097/mbp.0000000000000570>.
- [42] Cooke J, Carew S, Quinn C, O'Connor M, Curtin J, O'Connor C, et al. The prevalence and pathological correlates of orthostatic hypotension and its subtypes when measured using beat-to-beat technology in a sample of older adults living in the community. *Age Ageing.* 2013;42:709-14, <https://doi.org/10.1093/ageing/aft112>.
- [43] Fedorowski A, Ricci F, Hamrefors V, Sandau KE, Hwan Chung T, Muldowney JAS, et al. Orthostatic hypotension: management of a complex, but common, medical problem. *Circ Arrhythm Electrophysiol.* 2022;15:e010573, <https://doi.org/10.1161/CIRCEP.121.010573>.
- [44] Saz-Lara A, Cavero-Redondo I, Martínez-Vizcaíno V, Lucerón-Lucas-Torres M, Pascual-Morena C, Sequí-Domínguez I. Association between arterial stiffness and orthostatic hypotension: a systematic review and meta-analysis. *Front Physiol.* 2023;14:1164519, <https://doi.org/10.3389/fphys.2023.1164519>.
- [45] Laurent S, Cockcroft J, Van Bortel L,

Boutouyrie P, Giannattasio C, Hayoz D, et al. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J*. 2006;27:2588-605, <https://doi.org/10.1093/eurheartj/ehl254>.

[46] Tahvanainen AM, Tikkakoski AJ, Koskela JK, Nordhausen K, Viitala JM, Leskinen MH, et al. The type of the functional cardiovascular response to upright posture is associated with arterial stiffness: a cross-sectional study in 470 volunteers. *BMC Cardiovasc Disord*. 2016;16:101, <https://doi.org/10.1186/s12872-016-0281-8>.

[47] Kato M, Nihei Green F, Hotta K,

Tsukamoto T, Kurita Y, Kubo A, et al. The efficacy of stretching exercises on arterial stiffness in middle-aged and older adults: a meta-analysis of randomized and non-randomized controlled trials. *Int J Environ Res Public Health*. 2020;17:5643, <https://doi.org/10.3390/ijerph17165643>.

[48] Li G, Lv Y, Su Q, You Q, Yu L. The effect of aerobic exercise on pulse wave velocity in middle-aged and elderly people: a systematic review and meta-analysis of randomized controlled trials. *Front Cardiovasc Med*. 2022;9:960096, <https://doi.org/10.3389/fcvm.2022.960096>.