

1
2 Plasma proteomic analysis of intermuscular fat links muscle integrity with processing speed in
3 older adults.

4
5 Authors

6 Toshiko Tanaka^{1*}, Caterina Rosano^{2*}, Xiaoning Huang³, Qu Tian¹, Bennett A. Landman⁴, Ann Z
7 Moore¹, Iva Miljkovic², Andrew Perry⁵, Sadiya Khan³, Ravi Kalhan⁶, John Jeffrey Carr⁷, James
8 G. Terry⁷, Kristine Yaffe⁸, Keenan Walker⁹, Julián Candia¹, Luigi Ferrucci¹

- 9
10 1. Longitudinal Studies Section, Translational Gerontology Branch, NIH, NIA, 251
11 Bayview Boulevard, Baltimore MD, 21224, USA
12 2. Department of Epidemiology, School of Public Health, University of Pittsburgh, 3550
13 Terrace Street, Pittsburgh, PA, 15261, USA
14 3. Division of Cardiology, Department of Medicine, Feinberg School of Medicine,
15 Northwestern University, 676 N Saint Clair, Chicago, IL, 60611
16 4. Department of Computer Science, Vanderbilt University, 1211 Medical Center Drive,
17 Nashville, TN, 37232, USA
18 5. Vanderbilt Translational and Clinical Cardiovascular Research Center, Vanderbilt
19 University School of Medicine, 2525 West End Avenue, Nashville, TN, 37203, USA
20 6. Division of Pulmonary and Critical Care Medicine, Department of Medicine,
21 Northwestern University Feinberg School of Medicine, 676 North Saint Clair Street
22 Chicago, IL, 60611, USA
23 7. Department of Electrical and Computer Engineering, Vanderbilt University School of
24 Medicine, Nashville, 2301 Vanderbilt Place, TN, 37235, USA
25 8. UCSF Weill Institute for Neurosciences, University of California–San Francisco, San
26 Francisco, 1651 4th St, CA, 94158, USA
27 9. Laboratory of Behavioral Neuroscience, National Institute on Aging, Intramural Research
28 Program, Baltimore, 251 Bayview Boulevard, MD 21224, USA

29
30 *co-first authors

31
32 Corresponding author: Toshiko Tanaka PhD.
33 tanakato@mail.nih.gov
34 251 Bayview Boulevard, Baltimore, Maryland 21224
35
36
37
38
39
40
41
42
43
44
45
46

47

48 **Abstract**

49 **INTRODUCTION:** More intermuscular fat (IMF) has been associated with lower cognitive
50 performance and faster age-associated decline in cognitive function however, the mechanisms
51 driving this relationship have not been fully elucidated. We utilized proteomic analyses to
52 identify the molecular mediators of the association between IMF and cognition to gain further
53 insight into the mechanisms underlying this association.

54

55 **METHODS.** In this cross-sectional study, the plasma proteomic profile of IMF was assessed in
56 the Baltimore Longitudinal Study on Aging (BLSA; n=941, age=66.7±15.2) and validated in the
57 Coronary Artery Risk Development in Young Adults Study (CARDIA; n=2451, age=50.2±3.6).
58 The 7628 plasma proteins were assessed using an aptamer-based assay and tested for association
59 with IMF from the thigh (BLSA) and abdomen (CARDIA). Processing speed assessed by Digit
60 Symbol Substitution Test (DSST). Associations between the main exposures, outcome and
61 mediators were evaluated using linear regression, and mediating effects were assessed by causal
62 mediation analysis adjusting for age, sex, muscle area or muscle volume, self-reported race, and
63 years of education.

64

65 **RESULTS.** Higher IMF was associated with lower DSST performance both in the BLSA and
66 CARDIA studies. There were 722 plasma proteins associated with IMF in both the discovery and
67 replication cohorts (FDR-adjusted $p \leq 0.05$). Of the 722 IMF -associated proteins, 26 (24 unique
68 proteins) mediated the relationship between IMF and processing speed with mediation effects
69 ranging from 2.8 to 20.9% ($p \leq 0.05$). Overrepresentation analysis of the IMF-associated proteins

70 showed enrichment of proteins in synaptic function and organization, and growth factor binding
71 (FDR-adjusted $p \leq 0.05$).

72

73 **DISCUSSION.** There is a robust proteomic signature explaining, at least in part, the link of IMF
74 with DSST. This signature reflected neurological function and growth factor regulation, which
75 are both implicated in lower processing speed. Reducing IMF through behavioral or
76 pharmacological intervention may improve cognition through reduction in growth factor activity
77 and improvements in synaptic activity.

78

79

80

81 **Background**

82 Aging is associated with decline in cognitive function across different domains [1].
83 Slowing rates of decline in cognitive function is an important public health goal, as cognitive
84 decline is associated with unfavorable aging outcomes such as disability and dementia that could
85 influence one's quality of life [2]. There is growing evidence linking skeletal muscle health with
86 cognitive decline, thus understanding the biology linking muscle with cognitive function may
87 identify novel targets to slow the rates of cognitive decline with aging [3, 4].

88 Intermuscular fat (IMF) increases with older age, and it has been associated with worse
89 cognition in older adults, particularly with poorer processing speed [3-9]. It has been
90 hypothesized that neurodegenerative effects of IMF may be triggered by inflammatory and
91 cardio-metabolic disorders. When adipocytes reach their fat storage capacity, greater adipose
92 tissue accumulates ectopically, that is in organs that usually have little adiposity, such as the
93 skeletal muscle. As adipose tissue accumulates in skeletal muscle, both adipocytes and myocytes
94 may release pro-inflammatory cytokines and these in turn can increase risk of systemic
95 metabolic abnormalities such as insulin resistance and type 2 diabetes [10, 11]. It has also been
96 shown that myokines and adipokines (cytokines releases from myocytes and adipocytes,
97 respectively) have a direct signaling effects on the central nervous system, influencing amyloid
98 clearance and mitochondrial function in the brain [12-17]. However, evidence is primarily from
99 animal models or human studies at the extremes of muscle health (e.g. athletes or patient
100 populations); the molecules linking IMF with poorer cognitive function have not been directly
101 measured in community- dwelling older adults without overt disease.

102 To address this gap in knowledge, we aimed to characterize the plasma proteomic profile
103 of IMF and explore whether IMF-associated proteins mediate the relationship between IMF and

104 cognitive function, focusing on processing speed. By taking an agnostic approach to explore the
105 molecular pathways that are represented by the IMF-associated proteins, the results of this study
106 provide mechanistic insights into the link between IMF and cognitive function.

107

108 **Methods**

109 **Study Design and Cohort Description**

110 The discovery cohort, the Baltimore Longitudinal Study of Aging (BLSA), is a rolling
111 enrollment population-based study of participants residing predominantly in the Baltimore-
112 Washington DC area [18]. Briefly, eligibility criteria include the absence of major chronic
113 diseases (except for controlled hypertension), and cognitive or functional disabilities.
114 Participants are continually followed at varying time intervals based on their age, namely every 4
115 years for participants younger than 60 years, every 2 years between 60-79 years, and every year
116 for participants older than 80 years. Demographic characteristics including age, sex, self-reported
117 race, and years of education were obtained during a structured interview by BLSA staff. The
118 study protocol (Protocol number 03-AG-0325) was approved by the National Institutes of Health
119 Intramural Research Program Institutional Review Board and informed consent was obtained
120 from participants at each visit.

121 The validation cohort, Coronary Artery Risk Development in Young Adults (CARDIA)
122 study began in 1985 with the recruitment of 5,115 participants aged 18 to 30 years at field
123 centers located in Birmingham, AL, Chicago, IL, Minneapolis, MN, and Oakland, CA [19].
124 Recruitment was balanced for equal inclusion of black and white as well as female and male
125 participants, age (18–24, 25–30 years), and education (≤ 12 years, >12 years). The current study
126 includes data from 3,172 participants who agreed and received abdominal CTs at the year 25

127 (Y25) examination, of these, a total of 2,390 had DSST, proteomics and other key factors such as
128 age, sex, race, and education. All participants provided written informed consent, and
129 institutional review boards from each field center and the coordinating center approved the study
130 annually.

131

132 **Assessment of intermuscular fat**

133 IMF was quantified using CT scan of the thigh (BLSA) or abdomen (CARDIA). In the BLSA,
134 Imaging was performed by the Radiology Department of MedStar Harbor Hospital (Baltimore,
135 MD) using a Somatom Sensation 10CT scanner (Siemens, Malvern, PA). A two-stage deep
136 learning pipeline was implemented to estimate IMF from 2D thigh CT slices [20, 21]. From this
137 algorithm, the total muscle area, and IMF was calculated as the average of the right and left
138 thigh. In the CARDIA study, scans were performed using 64-channel multidetector GE CT
139 scanners (GE Healthcare Milwaukee, WI) at the Birmingham, AL, and Oakland, CA, centers and
140 Siemens CT scanners (Siemens, Erlangen, Germany) at the Chicago, IL, and Minneapolis, MN,
141 centers. The protocol has been previously published [22, 23].

142

143 **Cognitive function assessment**

144 In both cohorts, the Digit Symbol Substitution Test (DSST) was used to evaluate processing
145 speed [24]. During the test, participants are initially shown nine pairs of digits and symbols. In
146 the next sequence participants are provided a series of digits and asked to draw the matching
147 symbols for each digit. The score is the number of correct digit-symbol pairs identified during a
148 period of 90 seconds. The DSST values were analyzed as a z-score.

149

150 **Proteomic Assessment**

151 Proteomic assessment in fasting plasma samples were conducted using the 7k SomaScan assay
152 v4.1 (SomaLogic, Inc.; Boulder, CO, USA) in both BLSA and CARDIA, following a protocol
153 that was detailed elsewhere [25, 26]. The final normalized dataset (hybridization control, median
154 signal, plate-scale, inter-plate normalization) was used for the analysis. The final protein count
155 consisted of 7,268 target annotated human SOMAmers that were assessed and passed quality
156 control in both the BLSA and CARDIA study. Some proteins were targeted by multiple
157 SOMAmers, thus 6583 unique human proteins are represented in the panel. From this point
158 forward, each SOMAmer will be referred to proteins in the manuscript. Each protein relative
159 abundance (RFU) was log₂ transformed and scaled before analysis.

160

161

162 **Statistical Analysis**

163 All analyses were conducted using R version 4.2.2. The associations between IMF and
164 plasma proteins or DSST were assessed using linear regression models. For proteomic analysis
165 of IMF, the winsorized protein value was the dependent variable, and IMF the independent
166 variable in the regression model. The models were adjusted for age, sex, muscle area (or muscle
167 volume for CARDIA), self-reported race, and years of education. For the proteomic analysis, the
168 p-value was adjusted using false discovery rate (FDR), and an FDR adjusted $p \leq 0.05$ was
169 considered statistically significant. Proteins that were significantly associated with IMF in the
170 BLSA were tested for validation in CARDIA study. Significant validation was considered at
171 FDR adjusted $p \leq 0.05$. To test whether the IMF-associated proteins mediate the association
172 between IMF and DSST, a causal mediation analysis was conducted using R regmedint package

173 (version 1.0.1)[27]. For each model, IMF was the main exposure variable, DSST the dependent
174 variable, and individual protein was the mediator. Mediation was considered significant when a
175 significant indirect association for IMF on DSST was observed ($p \leq 0.05$).

176 Over-representation analysis (ORA) was conducted to determine whether any pathways
177 were enriched among the IMF-associated proteins. Using the proteins measured by the
178 SomaScan platform as background, we used the 716 IMF-associated proteins to explore
179 enrichment of GO biological processes (BP), cellular component (CC), and molecular function
180 (MF) libraries. Enrichment was considered significant at FDR adjusted $p \leq 0.05$. The over-
181 representation analysis was performed using R ClusterProfiler package (version 4.6.2).

182 183 **Results**

184 *Association between intermuscular fat and DSST.* The average age of the discovery cohort was
185 older than the validation cohort (**Table 1**; 66.7 ± 15.2 vs 50.2 ± 3.6). The age difference is reflected
186 in processing speed, with higher DSST scores in the CARDIA study compared to BLSA. Despite
187 these differences, IMF was found to be statistically significant in both studies. In the BLSA,
188 there was significant negative relationship between IMF and DSST (β [95%CI]=-0.11 [-0.15,-
189 0.05]) indicating that higher IMF was associated with slower DSST. Similarly, in the CARDIA
190 study a negative relationship was also observed (β [95%CI]=-0.09[-0.12,-0.05]).

191
192 *Proteomics of intermuscular fat* – In the discovery cohort, of 7268 SOMAmers (6375 proteins)
193 assessed, 782 SOMAmers (710 proteins) were significantly associated with IMF (Supplemental
194 Table 1; Figure 1A). There were 321 SOMAmers (297 proteins) found in higher abundance with
195 greater IMF, and 461 SOMAmers (413 proteins) in lower abundance with higher IMF. These 782

196 SOMAmers were tested for association with IMF in the CARDIA study, and 722 (654 proteins;
197 92.3%) were found to be significantly associated. For all proteins, the direction of association
198 was consistent, with high correlation ($r=0.95$, Figure 1B) between the beta estimates from the
199 regression models from the two studies for the 782 SOMAmers that were tested for replication.

200 An ORA analysis based on GO terms for the 654 proteins represented by the 722 IMF-
201 associated SOMAmers, identified multiple pathways (Figure 1C). The GO-biological process
202 (BP) terms encompass two broad categories of pathways that represent synaptic biology (
203 “regulation of synapse assembly”, “regulation of synapse structure and activity”, “axon
204 development”) and inflammation (“taxis” and “chemotaxis”). The GO-cellular component (CC)
205 terms complement the results from synaptic function from the BP that include CC terms such as
206 “intrinsic component of synaptic membrane, GABA-ergic synapse” and “axonal growth cone”,
207 as well as indicator of extracellular matrix. Finally, the GO-molecular function (MF) identified
208 included “glycosaminoglycan binding” and “growth factor binding”, and terms representing
209 general signal transduction activities.

210
211 *Proteomic mediators of IMF and DSST.* Of the 722 IMF-associated SOMAmers, 26 SOMAmers
212 representing 24 proteins significantly mediated the relationship between IMF and DSST in both
213 BLSA and CARDIA studies (Table S2; Figure 2). The proportion mediation ranges were 7.2-
214 20.9% in BLSA and 2.8-8.9% in CARDIA. For all but one protein, the directions of indirect
215 effects was negative in both the BLSA and CARDIA studies. Among the 24 mediating proteins,
216 14 were found in lower abundance with higher IMF, and associated with higher DSST.
217 Conversely, 10 proteins were found in higher abundance with greater IMF and associated with

218 lower DSST. The ORA analysis of the 24 proteins that mediate the relationship represent
219 pathways showed no significantly enriched pathways.

220

221 **Discussion**

222 In this study, we explored the plasma proteomic profiles of IMF showed that there is a
223 distinct proteomic signature of 654 proteins in two independent cohorts of older adults living in
224 the community. The protein signature was highly consistent despite IMF being measured in
225 skeletal muscle at different locations. A subset of 24 IMF-associated proteins explain, at least in
226 part, the association between higher IMF and slower processing speed. Taking an agnostic
227 approach, enrichment analyses of the proteins identified indicate that pathways such as synaptic
228 function and growth factor activity may be the link between IMF and processing speed.

229 The study results confirm the growing evidence supporting the associations between
230 higher IMF and lower cognitive function, particularly with slower processing speed. Since the
231 initial reporting inverse associations of measures of IMF using peripheral quantitative computer
232 tomography (pQCT) with measured of psychomotor function, visual learning, and overall
233 cognitive function [28], similar associations have been reported in different populations
234 particularly with processing speed [3, 4, 9] and with other indicators of brain health such as
235 neuroimaging markers [29, 30]. Our study explored used proteomic to explore the mechanisms
236 that may link higher IMF with slower processing speed.

237 One intriguing insight from our analysis was that the proteome associated with IMF was
238 enriched with proteins involved in synaptic organization and function. The four proteins
239 (SLITRK3, NOTCH1, C1QL3, EPHA4) that significantly mediated the association were found
240 in lower abundance with higher IMF and lower processing speed. These proteins are expressed

241 widely across the central nervous system, with some also found in muscle tissue [31-34]. They
242 play crucial roles in brain function by influencing various processes: SLITRK3 regulates neurite
243 outgrowth [32]; C1QL3 participate in neuronal organization as trans-synaptic cell adhesion
244 complexes [33]; NOTCH is involved in neuronal stem cell maintenance and activation [31, 34];
245 and EPH4 is crucial for synaptic function and integrity through regulation of bidirectional
246 communication between neurons and astrocytes [35]. SLITRK3 knockout mice have impaired
247 postsynaptic neurotransmission resulting in multiple abnormal hypermobile movements[36].
248 C1QL3 knockout animals display fewer excitatory neurons compared to wildtype and exhibited
249 stunted fear conditioning as well as a modest deficit in motor learning [37]. In humans, several
250 lines of evidence support the link between these proteins with cognitive function. First, lower
251 abundance of soluble NOTCH1 protein was reported in patients with AD, further protein
252 abundances were positively correlated with MMSE [38]. Mendelian randomization study
253 supported the causal association between SLITRK3 with general cognitive function [39]. Taken
254 together, our data suggest lower levels of proteins involved in synaptic function and integrity
255 simultaneously manifest as lower integrity in the skeletal muscle (higher IMF) and central
256 nervous system (slower processing speed). Our results of causal modeling analyses indicate a
257 potential direction of these associations, whereby higher IMF may affect processing speed by
258 lowering the circulating levels of neuroprotective proteins.

259 Among the proteins that were found in higher abundance in relation to higher IMF and
260 lower DSST (e.g. with potential neurodegenerative effects), the myokine High-Temperature
261 Requirement A1 (HTRA1) is notable, for its proteolytic effects on extracellular matrix
262 remodeling skeletal muscle, brain, and adipose tissue[40, 41]. HTRA1 has also inhibitory effects
263 on the transforming growth factor- β superfamily. In a translational proteomic study of several

264 thousand proteins[42]that HTRA1 was positively correlated with brain amyloid. Several
265 HTRA1 mutations have been associated with cerebral small vessel disease in humans [43], but
266 these studies did not test associations with cognitive function. We and others have shown that
267 higher cerebral small vessel disease is strongly associated with lower processing speed; thus it is
268 possible that higher IMF leads to higher levels of the myokine HTRA1, this in turn drives
269 cerebral small vessel disease, manifesting as slower processing speed.

270 The finding that proteins related to IGF are related with IMF and cognition has been
271 previously shown. Of note, IGF binding protein 2 (IGFBP-2) was positively associated with
272 skeletal muscle mass and strength in prior proteomic studies in South Africa[44], and
273 Germany[45] . Conversely, its role in relation with cognition is less clear. In a recent study of
274 older adults[46], IGFBP2 was associated with smaller hippocampal volume among amyloid
275 negative adults, but not amyloid positive, indicating the influence on IGFBP2 on hippocampal
276 volume varies depending on severity of brain pathology; conversely, others found a positive
277 association with tau[48] . The association of IGFBP2 with cognitive status also varies across
278 studies, presumably because the role of IGFBP-2 varies depending on developmental stages[47,
279 48]. Differences in age, and/or in brain pathology may explain why in our study IGFBP-2 had an
280 inverse association with DSST in CARDIA (younger) but not in BLSA (older).

281 Prior studies have linked IMF with factors that influence cardiometabolic health, in
282 particular serum cytokine levels and adipokines [11]. Both muscle and the fat that infiltrates
283 muscle have secretory capabilities including pro- and anti- inflammatory cytokines that have
284 been associated with cognitive health [49]. Moreover, cytokines released by skeletal muscles and
285 adipocytes with neuronal signaling properties can cross the blood brain barrier [50].
286 Consequently, some studies explored these proteins as possible mechanism linking IMF with

287 cognitive function. In the Geelong osteoporosis study (GOS), TNF α explained the association
288 between muscle density measured in the radius and psychomotor function, but the mediation was
289 not established for the muscle density measured in the tibia [28]. Further, other inflammatory
290 markers, IL-6, IL-10, and IL-13 did not explain the association in the GOS study. In other
291 studies, while no formal mediation analysis was conducted, the association between IMF and
292 cognitive function were robust to adjustment for cardiometabolic conditions, including
293 hypertension, diabetes, obesity, measures of inflammation (leptin, adiponectin, IL-6) [4]. In the
294 current report, IL-6 was moderately associated with IMF in the BLSA, and CRP was negatively
295 associated with IMF in both BLSA and CARDIA studies but were not significant mediators.
296 Intriguingly, enrichment analysis did not show inflammation as a major mediating pathway
297 linking IMF with processing speed.

298 While our findings provide strong biological evidence linking IMF with processing
299 speed, there are several important limitations. First, this is a cross-sectional study, thus we cannot
300 make causal inferences on the associations we observe. For example, it is possible that higher
301 IMF may drive lower levels of proteins, which in turn compromise synaptic integrity in the
302 brain, ultimately manifesting as slower processing speed. Alternatively, there is another factor
303 that alters homeostasis of proteins engaged in synapsis remodeling among older adults with
304 higher IMF. Further, there may be a bi-directional relationship – those with lower cognitive
305 function and processing speed may have lowered synaptic activity leading to decreased muscle
306 quality. While our study uses direct measurements of IMF using CT scans, the levels of IMF are
307 correlated with other measures of adiposity thus we cannot exclude the possibility that the
308 correlations we observed are not driven by fat infiltrations in the muscle per se, but through
309 overall adiposity. The analysis was not adjusted for measures of adiposity due to the correlation

310 between the two traits, and overall adiposity may be part of the casual pathways linking
311 intermuscular fat with processing speed. Our study has several strengths. First, our results were
312 consistent across two studies that have different demographics from the standpoint of age, self-
313 reported race, and geographical distributions. In both studies, IMF was measured using CT-scan
314 providing us with an objective measure. Finally, while the proteomic assessment did not measure
315 all circulating proteins, the analysis targeted large number of proteins using the same assessment
316 method. Future studies should explore validating these findings in longitudinal studies to
317 determine the temporal relationship between proteins abundances, IMF, and processing speed.

318 By leveraging data on circulating proteins, we provide strong evidence suggesting that
319 IMF affects processing speed through inflammation, regulating of growth factor activity, and
320 synaptic organization, function and activity. These data support the interplay between the
321 muscular quality and the brain and suggest that the muscle is an important target for prevention
322 of cognitive decline associated with aging.

323

324 **References**

325 [1] Harada CN, Natelson Love MC, Triebel KL. Normal cognitive aging. *Clin Geriatr Med*.
326 2013;29:737-52.

327 [2] Amieva H, Meillon C, Proust-Lima C, Dartigues JF. Is Low Psychomotor Speed a Marker of
328 Brain Vulnerability in Late Life? Digit Symbol Substitution Test in the Prediction of Alzheimer,
329 Parkinson, Stroke, Disability, and Depression. *Dement Geriatr Cogn Disord*. 2019;47:297-305.

330 [3] Tian Q, Lee PR, Yang Q, Moore AZ, Landman BA, Resnick SM, et al. The mediation roles of
331 intermuscular fat and inflammation in muscle mitochondrial associations with cognition and
332 mobility. *J Cachexia Sarcopenia Muscle*. 2024;15:138-48.

- 333 [4] Rosano C, Newman A, Santanasto A, Zhu X, Goodpaster B, Miljkovic I. Increase in skeletal
334 muscular adiposity and cognitive decline in a biracial cohort of older men and women. *J Am*
335 *Geriatr Soc.* 2023;71:2759-68.
- 336 [5] Booranasuksakul U, Macdonald IA, Stephan BCM, Siervo M. Body Composition,
337 Sarcopenic Obesity, and Cognitive Function in Older Adults: Findings From the National Health
338 and Nutrition Examination Survey (NHANES) 1999-2002 and 2011-2014. *J Am Nutr Assoc.*
339 2024:1-14.
- 340 [6] Spauwen PJ, Murphy RA, Jónsson PV, Sigurdsson S, Garcia ME, Eiriksdottir G, et al.
341 Associations of fat and muscle tissue with cognitive status in older adults: the AGES-Reykjavik
342 Study. *Age Ageing.* 2017;46:250-7.
- 343 [7] Uchida K, Sugimoto T, Tange C, Nishita Y, Shimokata H, Saji N, et al. Association between
344 Reduction of Muscle Mass and Faster Declines in Global Cognition among Older People: A 4-
345 Year Prospective Cohort Study. *J Nutr Health Aging.* 2023;27:932-9.
- 346 [8] Wang W, Luo Y, Zhuang Z, Song Z, Huang N, Li Y, et al. Total and regional fat-to-muscle
347 mass ratio and risks of incident all-cause dementia, Alzheimer's disease, and vascular dementia. *J*
348 *Cachexia Sarcopenia Muscle.* 2022;13:2447-55.
- 349 [9] Acevedo-Fontanez AI, Cvejkus RK, Zmuda JM, Kuipers AL, Barinas-Mitchell E, Sekikawa
350 A, et al. Skeletal muscle adiposity is a novel risk factor for poor cognition in African Caribbean
351 women. *Obesity (Silver Spring).* 2023;31:2398-406.
- 352 [10] Correa-de-Araujo R, Addison O, Miljkovic I, Goodpaster BH, Bergman BC, Clark RV, et al.
353 Myosteatosis in the Context of Skeletal Muscle Function Deficit: An Interdisciplinary Workshop
354 at the National Institute on Aging. *Front Physiol.* 2020;11:963.

- 355 [11] Miljkovic I, Vella CA, Allison M. Computed Tomography-Derived Myosteatorosis and
356 Metabolic Disorders. *Diabetes Metab J.* 2021;45:482-91.
- 357 [12] Arjunan A, Song J. Pharmacological and physiological roles of adipokines and myokines in
358 metabolic-related dementia. *Biomed Pharmacother.* 2023;163:114847.
- 359 [13] Burtcher J, Millet GP, Place N, Kayser B, Zanou N. The Muscle-Brain Axis and
360 Neurodegenerative Diseases: The Key Role of Mitochondria in Exercise-Induced
361 Neuroprotection. *Int J Mol Sci.* 2021;22.
- 362 [14] Heo J, Noble EE, Call JA. The role of exerkinines on brain mitochondria: a mini-review. *J*
363 *Appl Physiol (1985).* 2023;134:28-35.
- 364 [15] Jo D, Yoon G, Kim OY, Song J. A new paradigm in sarcopenia: Cognitive impairment
365 caused by imbalanced myokine secretion and vascular dysfunction. *Biomed Pharmacother.*
366 2022;147:112636.
- 367 [16] Lee TH, Formolo DA, Kong T, Lau SW, Ho CS, Leung RYH, et al. Potential exerkinines for
368 physical exercise-elicited pro-cognitive effects: Insight from clinical and animal research. *Int*
369 *Rev Neurobiol.* 2019;147:361-95.
- 370 [17] Oudbier SJ, Goh J, Looijaard S, Reijnierse EM, Meskers CGM, Maier AB.
371 Pathophysiological Mechanisms Explaining the Association Between Low Skeletal Muscle Mass
372 and Cognitive Function. *J Gerontol A Biol Sci Med Sci.* 2022;77:1959-68.
- 373 [18] Shock NW, Greulich RC, Andres R, Arenberg D, Costa. P, Lakatta. E, et al. Normal Human
374 Aging: The Baltimore Study of Aging. 1984.
- 375 [19] Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DR, Jr., et al.
376 CARDIA: study design, recruitment, and some characteristics of the examined subjects. *J Clin*
377 *Epidemiol.* 1988;41:1105-16.

- 378 [20] Yang Q, Yu X, Lee HH, Tang Y, Bao S, Gravenstein KS, et al. Label efficient segmentation
379 of single slice thigh CT with two-stage pseudo labels. *J Med Imaging (Bellingham)*.
380 2022;9:052405.
- 381 [21] Yang Q, Yu X, Lee HH, Tang Y, Bao S, Gravenstein KS, et al. Quantification of muscle,
382 bones, and fat on single slice thigh CT. *Proc SPIE Int Soc Opt Eng*. 2022;12032.
- 383 [22] Terry JG, Shay CM, Schreiner PJ, Jacobs DR, Jr., Sanchez OA, Reis JP, et al. Intermuscular
384 Adipose Tissue and Subclinical Coronary Artery Calcification in Midlife: The CARDIA Study
385 (Coronary Artery Risk Development in Young Adults). *Arterioscler Thromb Vasc Biol*.
386 2017;37:2370-8.
- 387 [23] Terry JG, Hartley KG, Steffen LM, Nair S, Alman AC, Wellons MF, et al. Association of
388 smoking with abdominal adipose deposition and muscle composition in Coronary Artery Risk
389 Development in Young Adults (CARDIA) participants at mid-life: A population-based cohort
390 study. *PLoS Med*. 2020;17:e1003223.
- 391 [24] Royer FL, Gilmore GC, Gruhn JJ. Normative data for the Symbol Digit Substitution Task. *J*
392 *Clin Psychol*. 1981;37:608-14.
- 393 [25] Candia J, Daya GN, Tanaka T, Ferrucci L, Walker KA. Assessment of variability in the
394 plasma 7k SomaScan proteomics assay. *Sci Rep*. 2022;12:17147.
- 395 [26] Choi B, Liu GY, Sheng Q, Amancherla K, Perry A, Huang X, et al. Proteomic Biomarkers
396 of Quantitative Interstitial Abnormalities in COPD Gene and CARDIA Lung Study. *Am J Respir*
397 *Crit Care Med*. 2024;209:1091-100.
- 398 [27] Li KYaY. regmedint: Regression-Based Causal Mediation Analysis with Interaction and
399 Effect Modification Terms }. 2022.

- 400 [28] Sui SX, Williams LJ, Holloway-Kew KL, Hyde NK, Anderson KB, Tembo MC, et al.
401 Skeletal Muscle Density and Cognitive Function: A Cross-Sectional Study in Men. *Calcif Tissue*
402 *Int.* 2021;108:165-75.
- 403 [29] Beck D, de Lange AG, Alnaes D, Maximov, II, Pedersen ML, Leinhard OD, et al. Adipose
404 tissue distribution from body MRI is associated with cross-sectional and longitudinal brain age in
405 adults. *Neuroimage Clin.* 2022;33:102949.
- 406 [30] Gurholt TP, Kaufmann T, Frei O, Alnaes D, Haukvik UK, van der Meer D, et al. Population-
407 based body-brain mapping links brain morphology with anthropometrics and body composition.
408 *Transl Psychiatry.* 2021;11:295.
- 409 [31] Ables JL, Decarolis NA, Johnson MA, Rivera PD, Gao Z, Cooper DC, et al. Notch1 is
410 required for maintenance of the reservoir of adult hippocampal stem cells. *J Neurosci.*
411 2010;30:10484-92.
- 412 [32] Aruga J, Mikoshiba K. Identification and characterization of Slitrk, a novel neuronal
413 transmembrane protein family controlling neurite outgrowth. *Mol Cell Neurosci.* 2003;24:117-
414 29.
- 415 [33] Sticco MJ, Pena Palomino PA, Lukacsovich D, Thompson BL, Foldy C, Ressler S, et al.
416 C1QL3 promotes cell-cell adhesion by mediating complex formation between ADGRB3/BAI3
417 and neuronal pentraxins. *FASEB J.* 2021;35:e21194.
- 418 [34] Wang R, Zhao C, Li J, Li Y, Liu Y, Dong H, et al. Notch1 promotes mouse spinal neural
419 stem and progenitor cells proliferation via p-p38-pax6 induced cyclin D1 activation. *Exp Cell*
420 *Res.* 2018;373:80-90.
- 421 [35] Murai KK, Pasquale EB. Eph receptors and ephrins in neuron-astrocyte communication at
422 synapses. *Glia.* 2011;59:1567-78.

- 423 [36] Efthymiou S, Han W, Ilyas M, Li J, Yu Y, Scala M, et al. Human mutations in SLITRK3
424 implicated in GABAergic synapse development in mice. *Front Mol Neurosci.* 2024;17:1222935.
- 425 [37] Martinelli DC, Chew KS, Rohlmann A, Lum MY, Ressler S, Hattar S, et al. Expression of
426 C1ql3 in Discrete Neuronal Populations Controls Efferent Synapse Numbers and Diverse
427 Behaviors. *Neuron.* 2016;91:1034-51.
- 428 [38] Cho SJ, Yun SM, Jo C, Jeong J, Park MH, Han C, et al. Altered expression of Notch1 in
429 Alzheimer's disease. *PLoS One.* 2019;14:e0224941.
- 430 [39] Tin A, Fohner AE, Yang Q, Brody JA, Davies G, Yao J, et al. Identification of circulating
431 proteins associated with general cognitive function among middle-aged and older adults.
432 *Commun Biol.* 2023;6:1117.
- 433 [40] Tai YK, Iversen JN, Chan KKW, Fong CHH, Abdul Razar RB, Ramanan S, et al. Secretome
434 from Magnetically Stimulated Muscle Exhibits Anticancer Potency: Novel Preconditioning
435 Methodology Highlighting HTRA1 Action. *Cells.* 2024;13.
- 436 [41] Tiaden AN, Bahrenberg G, Mirsaidi A, Glanz S, Bluher M, Richards PJ. Novel Function of
437 Serine Protease HTRA1 in Inhibiting Adipogenic Differentiation of Human Mesenchymal Stem
438 Cells via MAP Kinase-Mediated MMP Upregulation. *Stem Cells.* 2016;34:1601-14.
- 439 [42] Bai B, Wang X, Li Y, Chen PC, Yu K, Dey KK, et al. Deep Multilayer Brain Proteomics
440 Identifies Molecular Networks in Alzheimer's Disease Progression. *Neuron.* 2020;105:975-91 e7.
- 441 [43] Uemura M, Nozaki H, Kato T, Koyama A, Sakai N, Ando S, et al. HTRA1-Related Cerebral
442 Small Vessel Disease: A Review of the Literature. *Front Neurol.* 2020;11:545.
- 443 [44] Dlamini SN, Norris SA, Mendham AE, Mtintsilana A, Ward KA, Olsson T, et al. Targeted
444 proteomics of appendicular skeletal muscle mass and handgrip strength in black South Africans:
445 a cross-sectional study. *Scientific Reports.* 2022;12:9512.

- 446 [45] Huemer MT, Bauer A, Petrera A, Scholz M, Hauck SM, Drey M, et al. Proteomic profiling
447 of low muscle and high fat mass: a machine learning approach in the KORA S4/FF4 study. *J*
448 *Cachexia Sarcopenia Muscle*. 2021;12:1011-23.
- 449 [46] Lane EM, Hohman TJ, Jefferson AL. Insulin-like growth factor binding protein-2
450 interactions with Alzheimer's disease biomarkers. *Brain Imaging Behav*. 2017;11:1779-86.
- 451 [47] Khan S. IGFBP-2 Signaling in the Brain: From Brain Development to Higher Order Brain
452 Functions. *Front Endocrinol (Lausanne)*. 2019;10:822.
- 453 [48] Bonham LW, Geier EG, Steele NZR, Holland D, Miller BL, Dale AM, et al. Insulin-Like
454 Growth Factor Binding Protein 2 Is Associated With Biomarkers of Alzheimer's Disease
455 Pathology and Shows Differential Expression in Transgenic Mice. *Front Neurosci*. 2018;12:476.
- 456 [49] Zoico E, Rossi A, Di Francesco V, Sepe A, Oliosio D, Pizzini F, et al. Adipose tissue
457 infiltration in skeletal muscle of healthy elderly men: relationships with body composition,
458 insulin resistance, and inflammation at the systemic and tissue level. *J Gerontol A Biol Sci Med*
459 *Sci*. 2010;65:295-9.
- 460 [50] Geng J, Wang L, Zhang L, Qin C, Song Y, Ma Y, et al. Blood-Brain Barrier Disruption
461 Induced Cognitive Impairment Is Associated With Increase of Inflammatory Cytokine. *Front*
462 *Aging Neurosci*. 2018;10:129.

463

464 **Acknowledgement:** The BLSA was supported by the Intramural Research Program of the NIH,
465 National Institute on Aging

466

467 **Conflict of interest:** The authors have no conflicts to declare.

468

469 **Funding Sources:** The Coronary Artery Risk Development in Young Adults Study (CARDIA) is
470 conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in
471 collaboration with the University of Alabama at Birmingham (75N92023D00002 &
472 75N92023D00005), Northwestern University (75N92023D00004), University of Minnesota
473 (75N92023D00006), and Kaiser Foundation Research Institute (75N92023D00003). This
474 manuscript has been reviewed by CARDIA for scientific content. CARDIA proteomics data was
475 funded through the CARDIA Lung Study (R01 HL122477).

476
477 **Consent Statement:** All participants of the study in this report provided informed consent

478
479 **Data availability:** Data for this study can be accessed at the BLSA data coordinating center
480 (<https://www.blsa.nih.gov>), and CARDIA coordinating center (www.cardia.dopm.uab.edu).

481
482 **Key Words:** Proteomics; Intermuscular fat; Processing speed; Epidemiology; Aging

483
484 **Figure Legends:**




485 Figure 1. Plasma proteomic profile of intermuscular fat. Associations between intermuscular fat
486 (IMF) and plasma proteins were tested in participants of the BLSA (n=941) and validated in the
487 CARDIA study (n=2445). (A) In the discovery BLSA study, of the 7268 SOMAmers assessed,
488 782 were associated with IMF. (B) Of the 782 SOMAmers, 722 were validated in the CARDIA
489 study. The associations between the beta estimates from the regression between IMF and
490 SOMAmers in the BLSA and CARDIA study was high (r=0.96). (C) Enrichment analysis of GO
491 biological process (BP), cellular component (CC), and molecular functions (MF) of the 722


492 SOMAmers (representing 654 unique proteins) show enrichment in neurological pathways and
493 growth factor binding.

494

495 Figure 2. Association between intermuscular fat and processing speed mediation by plasma
496 proteins. The association between degree of intermuscular fat (IMF) and processing speed
497 assessed using DSST was significantly mediated by 24 proteins in both the BLSA and CARDIA
498 studies. Except for IGFBP2, all proteins show negative indirect effects. Red: There were 14
499 proteins that had negative associations with IMF (lower levels for higher IMF) and positive
500 association with DSST, indicating a potential neuroprotective role. * IGFBP2 was the only
501 protein with a negative association with DSST in CARDIA. Blue: there were 10 proteins with
502 positive associations with IMF (higher levels for higher IMF) and negative associations with
503 DSST, indicating a potential neurodegenerative role. Created with BioRender.com

504





Intermuscular
Fat


-

+

ADAMTS13
ALB
APLP1
C1QL3
EPHA4
IGFALS
IGFBP2*

IGLON5
NOTCH1
OMG
SCO1
SERPINC1
SLITRK3
TNFSF12

+



Processing
Speed

-

CSTB
DNAJB9
HTRA1
PLBD1
PLS3

PROK2
RIC8A
SUMO2
TRAPPC3
UBE2G2