

Supplemental Material

Supplemental Text

Supplemental Tables

Supplemental Table S1 - Fit indexes of the unidimensional models for irritability, outbursts and impairment Confirmatory Factor Analysis Models

Supplemental Table S2 – Factor Loadings and Category Threshold parameters for items in the Irritable Mood section

Supplemental Table S3 - Factor Loadings and Category Threshold parameters for items in the Temper Outbursts section

Supplemental Table S4 - Factor Loadings and Category Threshold parameters for items in the Impairment section

Supplemental Table S5 – Latent Class Model Fit Indexes and Entropy for each four latent class solutions for irritable mood, outbursts and impairment items

Supplemental Table S6 - Fit indexes of the unidimensional, correlated and bifactor models of irritable mood and temper outbursts

Supplemental Table S7 - Irritable mood, outburst and impairment items and their categories of response as described in DMDD section of DAWBA questionnaire

Supplemental Table S8 - Left Censored regression comparing DMDD groups in SDQ scores

Supplemental Table S9 - Psychiatric comorbidities in Disruptive Mood Dysregulation Disorder defined using different combination rules for the irritable mood and temper outburst domains

Supplemental Figures

Supplemental Figure S1 – Latent Class Clinical Indicator endorsement distribution between classes with low and high levels of irritable mood, outbursts and impairment

Supplemental Figure S2 – Receiver Operating Curve for predicting latent classes of subjects with high levels of irritable mood, outbursts and impairment

Supplemental Figure S3 – Comparison between Strengths and Difficulties Questionnaire Scores between Typically Developing Pre-Adolescents/Early Adolescents, Pre-Adolescents/Early Adolescents with other disorders, with irritable mood, outbursts and combined Disruptive Mood Dysregulation Disorder.

Supplemental Figure S4 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder and Matched controls for psychiatric comorbidity using the OR rule.

Supplemental Figure S5 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder and Matched controls for psychiatric comorbidity using the AND rule.

Supplemental code

Supplemental code 1 – Symptomatic thresholds

Supplemental code 2 – Syndromic thresholds

Supplemental code 3 – Clinical operationalization

Supplemental Text

Fit Indexes for Confirmatory Factor Analysis

For all CFA unidimensional models, we used delta parameterization and weighted least-square parameters using a diagonal weight matrix with standard errors and with mean and variance-adjusted chi-square test statistics (WLSMV) estimators. The measures of goodness of fit were assessed through the following fit indices: chi-square, SRMR (standardized root mean square residual), CFI (comparative fit index), TLI (Tucker-Lewis Index) and RMSEA, (root mean square error of approximation). To demonstrate good fit to the data, an estimated model should have a SRMR near or below 0.08, a RMSEA of near or below 0.06 and CFI and TLI near or above 0.95 (Hu & Bentler, 1999). Details about the estimators and the model fit used in this study can be found in Supplemental Table S1, which provides the fit indexes of the unidimensional models for irritability, outbursts and impairment items.

References

- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6(1), 1–55. <https://doi.org/10.1080/10705519909540118>

Supplemental Table S1 - Fit indices of the unidimensional irritable mood, outbursts and impairment Confirmatory Factor Analysis Models

Fit indexes	Irritable Mood*	Outbursts**	Impairment
FP	27	44	16
χ^2	3254.236 (df=28; p<0.001)	3352.504 (df=91; p<0.001)	2865.135 (df=6; p<0.001)
RMSEA	0.057	0.057	0.032
CI 90%	0.040- 0.074	0.047-0.068	0.000-0.088
CFI	0.989	0.968	1.000
TLI	0.984	0.961	0.999
SRMR	0.057	0.085	0.019

Note: FP: Free Parameters; χ^2 : Robust Chi Square Difference Test; df: degrees of freedom; RMSEA: Root Mean Square Error of Approximation; CI: Confidence Interval; CFI, Comparative Fit Index; TLI, Tucker-Lewis Index; SRMR: Standardized Root Mean Square Residual. * Items p1y12b-p1y12c allowed to correlated due to similar item content/context; ** Items p1y4b-p1y4c and p1y3b-p1y3c allowed to correlated due to correlated due to similar item content/context

Supplemental Table S2 – Factor Loadings and Category Threshold parameters for items in the Irritable Mood section

Items (item code in DAWBA)	Factor Loadings	T1	T2	T3	T4	Location (mean)	Location rank
<i>Frequency/duration</i>							
Frequency of irritable/angry mood		Occasionally	1-2/week	\geq 3/week	Every day		
	0.495			0.078	1.143	0.611	4
Long duration of irritability		<1hr	Few hours	Most/all of the day			
	0.420	0.053	0.98	1.726		0.920	6
<i>Characteristics</i>							
Easily irritated	0.905	-0.138	0.775			0.319	3
Intense irritability	0.910	0.281	0.973			0.627	5
<i>Settings</i>							
At home	0.624	-2.123	0.303			-0.910	1
In the classroom	0.292	0.471	1.521			0.996	7

With peers	0.423	0.438	1.608		1.023	8
			A medium			
Irritability evident to others		A little	amount	A great deal		
	0.437	-1.345	-0.095	1.256	-0.061	2

Note: T1= First threshold; T2 = Second threshold; T3 = Third threshold; T4 = Fourth threshold.

Supplemental Table S3 - Factor Loadings and Category Threshold parameters for items in the Temper Outbursts section

Items (item code in DAWBA)	Factor Loadings	T1	T2	T3	T4	Location (mean)	Location rank
<i>Frequency/duration</i>							
Frequency of outbursts		Occasionally	1-2/week	≥ 3/week	Every day		
	0.565			-0.068	1.129	-0.424	5
<i>Characteristics</i>							
		A little	A lot				
Slamming doors	0.525	-0.893	0.333			-0.274	6
Shouting	0.550	-1.004	0.062			-0.472	4
Swearing	0.693	-1.129	-0.015			-0.572	3
Saying mean things to others	0.743	-0.548	0.259			-0.143	7
Saying negative things about himself	0.481	0.633	1.316			0.981	10
Physical aggression to others	0.619	0.050	1.086			0.576	9
Deliberate self-harm	0.527	1.211	2.030			1.626	14
Breaking things	0.572	0.562	1.471			1.023	11

<i>Settings</i>		A little	A lot		
At home	0.722	-2.597	0.068	-1.264	1
In the classroom	0.351	0.604	1.628	1.123	13
With peers	0.372	0.448	1.585	1.024	12
<i>Triggers</i>					
Recognizable triggers		Perhaps	Definitely		
	-0.193	-0.815	-0.403	-0.610	2
Easily triggered		A little	A lot		
	0.664	-0.109	0.784	0.347	9

Note: T1 = First threshold; T2= Second threshold; T3= Third threshold; T4 = Fourth threshold.

Supplemental Table S4 - Factor Loadings and Category Threshold parameters for items in the Impairment section

Items	Factor Loadings	T1	T2	T3	Location (mean)	Location rank
		A little	A medium amount	A great deal		
Impact on family life	0.709	0.018	1.036	2.109	1.054	1
Impact on friendship	0.903	0.545	1.434	2.376	1.452	3
Impact on learning	0.825	0.601	1.238	2.443	1.427	2
Impact on leisure	0.826	0.672	1.521	2.757	1.650	4

Note: T1 = First threshold; T2 = Second threshold; T3 = Third threshold.

Table S5 – Latent Class Model Fit Indexes and Entropy for each of the four latent class solutions for irritable mood, outbursts and impairment items

Irritable Mood

Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-1.296.125	120	2.634.154	2.611.943	2.641.154	324.66270	-
Model 2	-1.191.353	112	2472.502	2424.906	2487.502	115.11877	0.886
Model 3	-1173.152	104	2483.992	2411.012	2506.992	78.71695	0.98
Model 4	-1167.479	96	2520.538	2422.173	2551.538	67.37120	0.948

Outbursts

Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-1501.228	247	3050.347	3024.963	3058.347	335.1590	-
Model 2	-1435.849	238	2973.468	2919.527	2990.468	204.4022	0.665
Model 3	-1424.724	229	3005.096	2922.597	3031.096	182.1518	0.634
Model 4	-1412.391	220	3034.308	2923.251	3069.308	157.4852	0.719

Impairment

Model	log-likelihood	resid. Df	BIC	aBIC	cAIC	likelihood-ratio	Entropy
Model 1	-1015.8047	11	2055.555	2042.863	2059.555	307.99668133	-

Model 2	-872.9925	6	1799.863	1771.306	1808.863	22.37226443	0.789
Model 3	-866.3003	1	1816.411	1771.988	1830.411	8.98796565	0.661
Model 4	-861.8353	-4	1837.413	1777.125	1856.413	0.05789396	0.738

Note: BIC, Bayesian Information Criterion; aBIC, adjusted Bayesian Information Criteria; cAIC,

Consistent Akaike Information Criterion; df, degrees of freedom

Supplemental Table S6 - Fit indices of the unidimensional, correlated and bifactor models of irritable mood and temper outbursts (n=398)

Fit indexes	Unidimensional	Correlated with two dimensions	Bifactor*
FP	31	32	46
χ^2	241.4	231.625	124.387
RMSEA	0.066	0.064	0.041
CI 90%	0.056-0.076	0.054-0.074	0.028- 0.054
CFI	0.926	0.930	0.975
TLI	0.847	0.917	0.965
SRMR	0.136	0.134	0.103

Note: FP: Free Parameters; χ^2 : Robust Chi Square Difference Test; df: degrees of freedom; RMSEA: Root Mean Square Error of Approximation; CI: Confidence Interval; CFI, Comparative Fit Index; TLI, Tucker-Lewis Index; SRMR: Standardized Root Mean Square Residual;

* The bifactor model resulted in negative residual variances and items with negative factor loadings in specific domains and, therefore, this model was not considered further in the analysis.

Supplemental Table S7 - Irritable mood, outburst and impairment items and their response categories as described in DMDD section of the DAWBA questionnaire

DAWBA Questionnaire Items	Response categories				
Irritable Mood Section					
Frequency of irritable/angry mood (p1y1)	Never	Occasionally	1-2/week	≥ 3/week	Every day
Easily irritated (p1y8)	No	A little	A lot		
Intense irritability (p1y9)	No	A little	A lot		
Long duration of irritability (p1y10)	No more than a few minutes	Less than an hour	A few hours	A medium amount	
Irritability evident to others (p1y11)	Not at all	A little	A great deal		
At home (p1y12a)	No	A little	A lot		
In the classroom (p1y12b)	No	A little	A lot		
With peers (p1y12c)	No	A little	A lot		
Angry weeks (irritable most of the day, nearly every day) * (p1y13)	No	Yes			
Outbursts Section					
Frequency of outbursts (p1y2)	Never	Occasionally	1-2/week	≥ 3/week	Every day

Slamming doors (p1y3a)	No	A little	A lot		
Shouting (p1y3b)	No	A little	A lot		
Swearing (p1y3c)	No	A little	A lot		
Saying mean things to others (p1y3d)	No	A little	A lot		
Saying negative things about himself (p1y3e)	No	A little	A lot		
Physical aggression to others (p1y3f)	No	A little	A lot		
Deliberate self-harm (p1y3g)	No	A little	A lot		
Breaking things (p1y3h)	No	A little	A lot		
At home (p1y4a)	No	A little	A lot		
In the classroom (p1y4b)	No	A little	A lot		
With peers (p1y4c)	No	A little	A lot		
Recognizable triggers (p1y5)	No	Perhaps	Definitely		
Easily triggered (p1y6)	No	A little	A lot		
Outburst-free gap in the last year ** (p1y7)	Less than a day	Less than a week	Less than a month	1-3 months	More than 3 months

Impairment Section					
			A medium		
Impact on family life (p1y19a)	Not at all	A little	amount	A great deal	

			A medium	
Impact on friendship (p1y19b)	Not at all	A little	amount	A great deal
			A medium	
Impact on learning (p1y19c)	Not at all	A little	amount	A great deal
			A medium	
Impact on leisure (p1y19d)	Not at all	A little	amount	A great deal

Note:

* The item “angry weeks” was not included because the item “long duration of irritability” already contains the response option that captures “irritable most of the day, nearly every day”.

** Item excluded of the analysis. The rationale for excluding this item is that it is unclear whether we would expect this item to be monotonically related to the overall latent construct given short periods of irritability with large gaps could also inform episodes of irritability (a marker of severity and bipolar disorder in children).

Supplemental Table S8 - Left Censored regression comparing DMDD groups in SDQ

scores

	Contrast with TDC			Contrast with Other DSM-5 Disorders		
	MD	SE	p-value	MD	SE	p-value
SDQ: Total						
Other disorders	9.54	0.35	<0.001	-	-	-
Irritable mood only	12.12	0.94	<0.001	2.57	1.07	0.016
Outbursts only	13.42	1.23	<0.001	3.87	1.37	0.005
Combined	15.56	0.89	<0.001	6.02	1.02	<0.001
SDQ: Emotional						
Other disorders	2.75	0.16	<0.001	-	-	-
Irritable mood only	2.94	0.42	<0.001	0.19	0.49	0.694
Outbursts only	2.46	0.55	<0.001	-0.28	0.63	0.657
Combined	3.31	0.40	<0.001	0.56	0.04	0.234
SDQ: Hyperkinetic						
Other disorders	3.65	0.21	<0.001	-	-	-
Irritable mood only	4.50	0.54	<0.001	0.85	0.53	0.113
Outbursts only	5.38	0.71	<0.001	1.73	0.68	0.011

Combined	5.35	0.52	<0.001	1.69	0.04	<0.001
----------	------	------	--------	------	------	--------

SDQ: Conduct

Other disorders	2.81	0.162	<0.001	-	-	-
-----------------	------	-------	--------	---	---	---

Irritable mood only	3.96	0.416	<0.001	1.16	0.46	0.111
---------------------	------	-------	--------	------	------	-------

Outbursts only	5.13	0.547	<0.001	2.33	0.59	<0.001
----------------	------	-------	--------	------	------	--------

Combined	5.71	0.396	<0.001	2.91	0.39	<0.001
----------	------	-------	--------	------	------	--------

SDQ: Peer relationships

Other disorders	2.42	0.17	<0.001	-	-	-
-----------------	------	------	--------	---	---	---

Irritable mood only	3.11	0.43	<0.001	0.69	0.48	0.146
---------------------	------	------	--------	------	------	-------

Outbursts only	2.82	0.57	<0.001	0.40	0.62	0.514
----------------	------	------	--------	------	------	-------

Combined	3.67	0.41	<0.001	1.25	0.46	<0.001
----------	------	------	--------	------	------	--------

SDQ: Impact

Other disorders	4.09	0.21	<0.001	-	-	-
-----------------	------	------	--------	---	---	---

Irritable mood only	5.35	0.47	<0.001	1.22	0.45	0.006
---------------------	------	------	--------	------	------	-------

Outbursts only	4.51	0.63	<0.001	0.42	0.59	0.711
----------------	------	------	--------	------	------	-------

Combined	6.07	0.45	<0.001	1.94	0.43	<0.001
----------	------	------	--------	------	------	--------

Note: TDC, Typically Developing Comparisons; SDQ, Strengths and Difficulties

Questionnaire; MD, Mean Difference.

Supplemental Table S9 – Psychiatric comorbidities in Disruptive Mood Dysregulation Disorder defined using different combination rules for the irritable mood and temper outburst domains


Psychiatric Comorbidities	Current DMDD (n=86)	Irritable Mood (n= 40)	Outbursts (n=22)	AND Rule (n=44)	OR Rule (n=107)
	Prevalence Estimation (%)				
<i>Any disorder (except DMDD)</i>	62.8	60.0	56.5	79.5	67.3
<i>Any Anxiety Disorder</i>	23.3	22.5	13.0	31.8	24.3
Generalized Anxiety Disorder	5.8	7.5	0	9.1	6.5
Separation Anxiety Disorder	7	2.5	4.3	13.6	7.5
Agoraphobia	3.5	2.5	0	4.5	2.8
Panic Disorder	1.2	0	0	2.3	0.9
Social Anxiety Disorder	3.5	10.0	0	4.5	5.6
Specific Phobia	11.6	2.5	8.7	15.9	9.3
Other Specified Anxiety Disorder	1.2	0	0	0	0
<i>Any Mood Disorder</i>	11.6	17.5	4.3	15.9	14.0
Major Depressive Disorder	8.1	12.5	4.3	11.4	10.3
Bipolar Disorder/Mania	3.5	7.5	0	2.3	3.7
Other Specified Depressive Disorder	1.2	2.3	0	2.5	1.9

<i>Any Attention-Deficit/Hyperactivity Disorder</i>	44.2	30.0	34.8	50.0	39.3
Combined	29.1	17.5	17.4	31.8	23.4
Hyperactive/impulsive	3.5	0	13.0	4.5	4.7
Inattentive	9.3	7.5	0	11.4	7.5
Another Specified ADHD	2.3	5	4.3	2.3	3.7
<i>Any Disruptive Behavior Disorder</i>	20.9	20.0	34.8	38.6	30.8
Oppositional Defiant Disorder	0	12.5	21.7	11.4	14.0
Conduct Disorder	18.6	10.0	17.4	27.3	18.7
Other Disruptive Behavior Disorder	2.3	0	4.3	4.5	2.8
<i>Other Disorders</i>					
Obsessive-Compulsive Disorder	1.2	2.5	0	2.3	1.9
Posttraumatic Stress Disorder	3.5	0	0	4.5	1.9
Autism Spectrum Disorder	3.5	0	4.3	6.8	3.7
Eating Disorders	1.2	0	0	2.3	0.9
<i>Current DMDD</i>	-	37.5	39.1	84.1	57.0


Note: DMDD, Disruptive Mood Dysregulation Disorder

Supplemental Figure S1 – Latent Class Clinical Indicator endorsement distribution between classes with low and high levels of irritable mood, outbursts and impairment

Irritable Mood



Outbursts




Impairment




Note: Graphs represents the chance of endorsement of each irritability, outburst and impairment clinical indicators between the two latent classes.


Supplemental Figure S2 – Receiver Operating Curve for predicting latent classes of subjects with high levels of irritable mood, outbursts and impairment



Supplemental Figure S3 – Comparison of Strengths and Difficulties Questionnaire Scores among Typically Developing Pre-Adolescents/Early Adolescents, Pre-Adolescents/Early Adolescents with non-DMDD disorders, with irritable mood only, outbursts only and both irritable mood and outbursts (combined).



Supplemental Figure S4 – Comparison of Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder, defined using the OR rule, and subjects matched for psychiatric comorbidity.



Supplemental Figure S5 – Comparison between Strengths and Difficulties Questionnaire Scores between Disruptive Mood Dysregulation Disorder, defined using the AND rule, and subjects matched for psychiatric comorbidity.

