

Supplementary Information

Cerebrospinal fluid p-tau217 performs better than p-tau181 as a biomarker of Alzheimer's disease

Supplementary Table 1. Spearman correlations between CSF p-tau, p-tau/t-tau and regional [¹⁸F]flortaucipir.

	Braak I/II ROI	Braak III/IV ROI	Braak V/VI ROI	Inferior temporal cortex
All (n=184)				
p-tau217	0.752 (<0.001)	0.734 (<0.001)	0.620 (<0.001)	0.698 (<0.001)
p-tau181	0.706 (<0.001)	0.687 (<0.001)	0.572 (<0.001)	0.650 (<0.001)
p-tau217/t-tau	0.778 (<0.001)	0.774 (<0.001)	0.660 (<0.001)	0.740 (<0.001)
p-tau181/t-tau	0.741 (<0.001)	0.752 (<0.001)	0.625 (<0.001)	0.720 (<0.001)
CU (n=63)				
p-tau217	0.410 (0.001)	0.212 (0.095)	0.011 (0.932)	0.212 (0.096)
p-tau181	0.330 (0.008)	0.208 (0.102)	0.017 (0.895)	0.214 (0.093)
p-tau217/t-tau	0.472 (<0.001)	0.234 (0.065)	0.048 (0.710)	0.240 (0.058)
p-tau181/t-tau	0.379 (0.002)	0.264 (0.036)	0.090 (0.481)	0.301 (0.016)
MCI-AD (n=28)				
p-tau217	0.663 (<0.001)	0.535 (0.003)	0.460 (0.014)	0.483 (0.009)
p-tau181	0.550 (0.002)	0.324 (0.093)	0.302 (0.118)	0.276 (0.155)
p-tau217/t-tau	0.620 (<0.001)	0.599 (0.001)	0.472 (0.011)	0.560 (0.002)
p-tau181/t-tau	0.534 (0.002)	0.403 (0.033)	0.345 (0.072)	0.361 (0.059)
AD dementia (n=40)				
p-tau217	0.288 (0.071)	0.552 (<0.001)	0.535 (<0.001)	0.549 (<0.001)
p-tau181	0.300 (0.060)	0.400 (0.011)	0.358 (0.023)	0.399 (0.011)
p-tau217/t-tau	0.238 (0.138)	0.760 (<0.001)	0.742 (<0.001)	0.768 (<0.001)
p-tau181/t-tau	0.206 (0.202)	0.399 (0.011)	0.318 (0.046)	0.420 (0.007)

Data are shown as Spearman correlation coefficients (p-value). Significant results are shown in bold. AD, Alzheimer's disease; CU, cognitively unimpaired; MCI, mild cognitive impairment; ROI, region of interest.

Supplementary Table 2. Demographic and clinical characteristics of validation cohort.

	Amyloid Positive Mild AD n=32
Age, <i>years</i>	73 (8)
Sex F/M, <i>n</i>	13/19
Education, <i>years</i>	15 (2)
MMSE	23 (2)
p-tau217, <i>pg/ml</i>	918 (773)
p-tau181, <i>pg/ml</i>	442 (313)

Data are shown as mean (SD) unless otherwise specified.

AD, Alzheimer's disease dementia; F, female; M, male; MMSE, Mini Mental State Examinations.

Supplementary Table 3. Performance of the p-tau217 and p-tau181 assays.

Sample	Mean concentration, pg/ml	SD	N	% CV
p-tau217				
10 pg/mL QC	9.97	0.59	22	5.95
50 pg/mL QC	44.31	2.06	22	4.66
High Control Sample	737.99	40.96	22	5.55
Medium Control Sample	328.72	17.81	22	5.42
Low Control Sample	59.69	4.93	22	8.26
p-tau181				
10 pg/mL QC	9.49	0.24	22	2.53
50 pg/mL QC	45.84	1.05	22	2.29
High Control Sample	379.61	21.25	22	5.60
Medium Control Sample	190.27	7.29	22	3.83
Low Control Sample	71.19	3.92	22	5.51

CV, coefficient of variation; QC, quality control; SD, standard deviation.