

		Included patients n = 41	Excluded patients due to incomplete specimen collection n= 5	p
Age^a		53.9 (12.6)	47.5 (12.0)	0.78
BMI^a		25.8 (4.3)	29.7 (9.8)	0.11
Gender^b	Male	28 (68.3)	2 (40)	0.21
	Female	13 (31.7)	3 (60)	
Highest level of education achieved^{b#}	Less than year 12	23 (56.1)	2 (40)	0.19
	Year 12 or higher	17 (41.5)	2 (40)	
Country of birth^{b#}	Australia	29 (70.7)	3 (60)	0.67
	Overseas	12 (29.3)	1 (20)	

Supplementary Table 1.

Descriptive characteristics. Data are expressed as ^amean (standard deviation) and ^bfrequency (percentage). *BMI = body mass index (kg/m²). # denotes missing data

		Included patients n = 41	Excluded patients due to incomplete specimen collection n= 5	P
Hepatitis C genotype	Unknown	2 (4.8)	4 (80)	<0.01
	1a	10 (24.3)	-	
	1b	12 (29.3)	-	
	3	16 (39.0)	1 (20)	
	4	1 (2.4)	-	
Hepatitis B co-infection		3 (7.3)	0	0.53

Supplementary Table 2.

Hepatitis C genotype and hepatitis B co-infection of patients with depression (study group) and without depression (control group). Data are expressed as frequency (percentage)

	Included patients n = 41	Excluded patients due to incomplete specimen collection n= 5	P
Comorbid cardiac disease	1 (2.4)	2 (40)	0.028
Comorbid diabetes	4 (9.8)	0	1.0
Comorbid arthritis	13 (31.7)	1 (20)	1.0
Comorbid thyroid disease	2 (4.9)	0	1.0
Past diagnosis of major depression#	23 (56.1)	1 (20)	0.18
Past diagnosis of bipolar affective disorder#	1 (2.4)	0	1.0
Past diagnosis of schizophrenia#	5 (12.2)	0	1.0
Past diagnosis of anxiety#	11 (26.8)	1 (20)	1.0
Past suicide attempt#	4 (9.7)	0	1.0

Supplementary Table 3.

Comorbidities and psychiatric history. Data are expressed as frequency (percentage). # denotes missing data