

		Antibody N	Antibody P/B
Sex, n (%)	Males	133 (22.09)	22 (17.60)
	Females	469 (77.91)	103 (82.40)
Age median (IQR)		45.71 (34.14-54.13)	41.95 (33.57-50.25)
Age groups (y), n (%)	22-30	103 (17.11)	22 (17.60)
	31-40	126 (20.93)	38 (30.40)
	41-50	164 (27.24)	35 (28.00)
	51-60	175 (29.07)	25 (20.00)
	61-77	34 (5.65)	5 (4.00)
Nasal/oral swabs, positive, n (%)		0 (0)	1 (0.8)
Exposure risk occupations, n (%)	High	270 (44.85)	65 (52.00)
	Medium	228 (37.87)	49 (39.20)
	Low	104 (17.28)	11 (8.80)
Professional information, n (%)	Medical doctor	143 (23.75)	47 (37.60)
	Nurse	180 (29.90)	32 (25.60)
	Obstetrician	43 (7.14)	10 (8.00)
	Healthcare operator	58 (9.63)	11 (8.80)
	Administration staff	56 (9.30)	8 (6.40)
	Laboratory technician	14 (2.33)	2 (1.60)
	Radiology technician	8 (1.33)	1 (0.80)
	Other	100 (16.61)	14 (11.2)
Professional risk in the last three months, yes, n (%)	Performance of invasive maneuvers (e.g. aerosol / bronchoalveolar lavage / intubation)	107 (17.77)	26 (20.80)
	Performance of Covid19 swabs on a patient	53 (8.80)	8 (6.40)
Epidemiological risk (out of the workplace), n (%)	Unknown contacts with positive/suspect Covid-19	464 (77.59)	89 (71.77)
	Contact with symptomatic/positive Covid-19	65 (10.87)	14 (11.29)
	Contact with symptomatic/negative Covid-	19 (3.18)	4 (3.23)

	19		
	Contact with asymptomatic/positive Covid-19	32 (5.35)	10 (8.06)
	Contact with symptomatic/suspect without Covid-19 confirmation	18 (3.01)	7 (5.65)
Personal medical history, yes, n (%)	Flu vaccine (2019-2020)	206 (34.22)	47 (37.60)
	Chronic conditions	223 (37.04)	39 (31.20)
	Chronic drugs	208 (34.55)	35 (28.00)
	Symptoms at 3 months, ≥ 1	444 (73.75%)	98 (78.40)
	Symptoms at 3 days, ≥ 1	242 (40.20)	55 (44.00)

Supplementary Table 3: Description of samples characteristics with regard to the antibody test. In the antibody columns: N=negative; P/B= Positive/Borderline. IQR=interquartile range; n (%)= number in percentage