

Table S1. Anxiety

<i>Study</i>	<i>Condition</i>	<i>Measure</i>	<i>Cut-off used</i>	<i>Sample</i>	<i>Prevalence</i>			<i>Timing</i>
					<i>Overall</i>	<i>High exposure</i>	<i>Low exposure</i>	
Xiao et al 2020 [1]	COVID-19	ZSRAS	Not used	HE only	Not given	Not given	Not given	During
C-H. Liu et al 2020 [2]	COVID-19	ZSRAS	>50	HE and LE	12.5%	Not given	Not given	During
Z. Liu et al 2020 [3]	COVID-19	ZSRAS	>49	HE and LE	16.0%	18.6%	14.0%	During
J. Huang et al 2020 [4]	COVID-19	ZSRAS	Not given	HE only	Not relevant	23.04%	Not relevant	During
Liang et al 2020 [5]	COVID-19	ZSRAS	Not used	HE and LE	Not given	Not given	Not given	During
Lai et al 2020 [6]	COVID-19	GAD-7	>4	HE and LE	44.6%	51.3%	39.4%	During
Zhu et al 2020 [7]	COVID-19	GAD-7	>7	HE and LE	24.36%	23.66%	24.83%	During
Sun et al 2020 [8]	COVID-19	SCL-90-A	Not used	HE and LE	Not given	Not given	Not given	During
Ji et al 2017 [9]	Ebola	SCL-90-R-A	Not used	HE and LE	Not given	Not given	Not given	During
McAlonan et al 2005 [10]	SARS	DASS-21-A	Not given	HE and LE	Not given	Not given	Not given	During & 1 year
R. Chen et al 2006 [11]	SARS	ZSRAS	>59*	HE Only	Not relevant	Not given	Not relevant	During, T1
Iancu et al 2005 [12]	SARS	MSAS	Not used	Undifferentiated	Not given	Not given	Not given	During
C.S. Chen et al 2005 [13]	SARS	SCL-90-R-A	Not used	HE and LE	Not given	Not given	Not given	During
Poon et al 2004 [14]	SARS	STAI	Not used	HE and LE	Not given	Not given	Not given	During

ZSRAS = Zung Self-rating Anxiety Scale; MSAS = Modified Spielberger Anxiety Scale; SCL-90-A = Symptom Checklist 90 Anxiety Subscale; SCL-90-R-A = Symptom Checklist 90 Revised Anxiety Subscale; DASS-21 = Depression, Anxiety and Stress Scale

Table S1. All studies measuring anxiety symptoms for healthcare responders for high risk epidemic and pandemic disease outbreak using validated scales

Table S2. PTSD symptoms

<i>Study</i>	<i>Condition</i>	<i>Measure</i>	<i>Cut-off used</i>	<i>Sample</i>	<i>Prevalence</i>			<i>Timing</i>
					<i>Overall</i>	<i>High exposure</i>	<i>Low exposure</i>	
J. Huang et al 2020 [4]	COVID-19	PTSD-SS	Not given	HE only	Not relevant	27.39%	Not relevant	During
Lai et al 2020 [6]	COVID-19	IES-R	>8	HE and LE	71.5%	76.0%	68.0%	During
Zhu et al 2020 [7]	COVID-19	IES-R	>33	HE and LE	30.56%	33.43%	28.65%	During
Xiao et al 2020 [1]	COVID-19	SASR	Not given	HE only	Not given	Not given	Not given	During
Matsuishi et al 2020 [15]	H1N1	IES-R	Not given	HE and LE	Not given	Not given	Not given	During
Lee et al 2018 [16]	MERS	IES-R	>24	HE and LE	51.5%	Not given	Not given	During
Bukhari et al 2016 [17]	MERS	IES	Idiosyncratic	HE and LE	Not given	Not given	Not given	During
Chan & Huak 2004 (doctors) [18]	SARS	IES	>29	HE and LE	Not given	18.8%	19.8%	During
Chan & Huak 2004 (nurses) [18]	SARS	IES	>29	HE and LE	Not given	19.4%	19.5%	
Chong et al 2004 [19]	SARS	IES	Not given	HE and LE	Not given	Not given	Not given	During
Koh et al 2005 [20]	SARS	IES	>16	HE and LE	Not given	Not given	Not given	During
C.S. Chen et al 2005 [13]	SARS	IES-R	>36	HE and LE	11%	17%	2%	During
Ho et al 2005 [21]	SARS	CIES-R	Not given		Not given	Not given	Not given	During
Su et al 2007 [22]	SARS	DTS-C	>22	HE and LE	28.43%	32.86%	18.75%	During
Sim et al 2004 [23]	SARS	IES	Idiosyncratic	HE and LE	9.4%	7.2%	10.6%	During
Styra et al 2008 [24]	SARS	IES-R	>19	HE and LE	Not given	Not given	Not given	During
Lin et al 2007 [25]	SARS	DTS-C	>40	HE and LE	19.3%	21.7%	13.0%	During
Tham et al 2004 / Phua et al 2005 [26, 27]	SARS	IES	>25	Undifferentiated	17.7%	Not relevant	Not relevant	6 months after
Maunder et al 2006 [28]	SARS	IES	>25	HE and LE	Not given	13.8%	8.4%	1-2 years after
Lancee et al 2008 [29]	SARS	CAPS	DX	HE only	Not relevant	2%	Not relevant	1-2 years after
McAlonan et al 2005 [10]	SARS	IES-R	Not given	HE and LE	Not given	Not given	Not given	During & 1 year
Wu et al 2008/2009 [30] [31]	SARS	IES-R	>19	HE and LE	10.1%	Not given	Not given	3 years after

PTSD-SS = PTSD Self Report Scale; IES = Impact of Events Scale; IES-R = Impact of Events Scale – Revised; SASR = Stanford Acute Stress Reaction; CIES-R = Chinese Impact of Event Scale-Revised; Chinese version of the Davidson Trauma Scale (DTS-C); CAPS = Clinician Administered PTSD Scale

Table 2. All studies measuring PTSD symptoms for healthcare responders for high risk epidemic and pandemic disease outbreak using validated measures

Table S3. Depression

<i>Study</i>	<i>Condition</i>	<i>Measure</i>	<i>Cut-off used</i>	<i>Sample</i>	<i>Prevalence</i>			<i>Timing</i>
					<i>Overall</i>	<i>High exposure</i>	<i>Low exposure</i>	
Zhu et al 2020 [7]	COVID-19	PHQ-9	>9	HE and LE	13.55%	13.73%	13.43%	During
Lai et al 2020 [6]	COVID-19	PHQ-9	>4	HE and LE	50.4%	58.4%	44.6%	During
Chung & Yeung 2020 [32]	COVID-19	PHQ-9	>9	Undifferentiated	34.8%	Not given	Not given	During
Liang et al 2020 [5]	COVID-19	ZSRAD	Not used	HE and LE	Not given	Not given	Not given	During
Z. Liu et al 2020 [3]	COVID-19	ZSRAD	>49	HE and LE	34.6%	37.2%	32.5%	During
R. Chen et al 2006 [11]	SARS	ZSRAD	>59*	HE Only	Not given	Not given	Not given	During, T1
Su et al 2007 [22]	SARS	BDI	>9	HE and LE	27.45%	38.57%	3.13%	During
McAlonan et al 2005 [10]	SARS	DASS-21	Not given	HE and LE	Not given	Not given	Not given	During & 1 year
Wu et al 2008 [30]	SARS	CES-D	>15	HE and LE	22.8%	Not given	Not given	3 years after

PHQ-9 = Patient Health Questionnaire 9; DASS-21 = Depression, Anxiety and Stress Scale; ZSRAD = Zung Self-rating Depression Scale; BDI = Beck Depression Inventory

Table 3. All studies measuring depression symptoms for healthcare responders for high risk epidemic and pandemic disease outbreak using validated measures

Table S4. Anxiety averages

Study	Condition	Measure	Summary stat	Sample	Average (SD if mean)			Timing
					Overall	High exposure	Low exposure	
Xiao et al 2020 [1]	COVID-19	ZSRAS	Mean	HE only	Not relevant	55.26 (14.18)	Not relevant	During
C-H. Liu et al 2020 [2]	COVID-19	ZSRAS	Mean	HE and LE	39.56 (8.91)	41.11 (9.79)	38.83 (8.38)	During
Z. Liu et al 2020 [3]	COVID-19	ZSRAS	Not given	HE and LE	Not given	Not given	Not given	During
J. Huang et al 2020 [4]	COVID-19	ZSRAS	Mean	HE only	42.91 (10.89)	Not given	Not given	During
Liang et al 2020 [5]	COVID-19	ZSRAS	Mean	HE and LE	Not given	28.09 (4.98)	29.36 (4.46)	During
Lai et al 2020 [6]	COVID-19	GAD-7	Median	HE and LE	4.0	5.0	3.0	During
Zhu et al 2020 [7]	COVID-19	GAD-7	Not given	HE and LE	Not given	Not given	Not given	During
Sun et al 2020 [8]	COVID-19	SCL-90-A	Mean	HE and LE	1.50 (0.79)	Not given	Not given	During
Ji et al 2017 [9] (Chinese staff)				HE only				During
Ji et al 2017 [9] (Sierra Leone staff)	Ebola	SCL-90-R-A	Mean		Not relevant	1.83 (0.23)	Not relevant	
Leone staff)	Ebola	SCL-90-R-A	Mean	HE only				During
Li et al 2015 [33]	Ebola	SCL-90-R-A	Mean	HE and LE	Not relevant	1.84 (0.89)	1.76 (0.75)	
McAlonan et al 2005 [10]	SARS	DASS-21	Mean	HE and LE	0.27 (0.42)	1.16 (0.52)	1.65 (0.99)	
R. Chen et al 2006 [11]	SARS	ZSRAS	Mean	HE and LE	Not given	4.9 (4.3)	2.1 (2.5)	During & 1 year
Su et al 2007 [22]	SARS	STAI	Mean	HE only	Not relevant	61 (12.62)	Not relevant	During, T1
Iancu et al 2005 [12]	SARS	MSAS	Mean	HE and LE	Not given	48.4 (10.6)	43.1 (8.1)	During
Iancu et al 2005 [12]	SARS	MSAS	Mean	Undifferentiated	13.5 (3.5)	Not relevant	Not relevant	During
C.S. Chen et al 2005 [13]	SARS	SCL-90-R-A	Mean	HE and LE	0.54 (0.54)	0.67 (0.75)	0.57 (0.46)	During
Poon et al 2004 [14]	SARS	STAI	Mean	HE and LE	Not given	52.6 (10.5)	49.8 (10.1)	During
Chan & Huak 2004 (doctors) [18]	SARS	GHQ-28 A	Mean	HE and LE	Not given	0.9 (1.8)	0.8 (1.4)	During
Chan & Huak 2004 (nurses) [18]	SARS	GHQ-28 A	Mean	HE and LE	Not given	0.7 (1.1)	0.6 (1.2)	During
Sim et al 2004 [23]	SARS	GHQ-28 A	Mean	HE and LE	Not given	0.8 (1.6)	0.7 (1.6)	During

ZSRAS = Zung Self-rating Anxiety Scale; MSAS = Modified Spielberger Anxiety Scale; SCL-90-A = Symptom Checklist 90 Anxiety Subscale; SCL-90-R-A = Symptom Checklist 90 Revised Anxiety Subscale; STAI = Spielberger Trait Anxiety Inventory.

Table S4. All studies reporting averages of validated anxiety symptom scales for healthcare responders for high risk epidemic and pandemic disease outbreaks

Table S5. PTSD symptom averages

<i>Study</i>	<i>Condition</i>	<i>Measure</i>	<i>Summary stat</i>	<i>Sample</i>	<i>Average (SD if mean)</i>			<i>Timing</i>
					<i>Overall</i>	<i>High exposure</i>	<i>Low exposure</i>	
J. Huang et al 2020 [4]	COVID-19	PTSD-SS	Mean	HE only	42.92 (17.88)	Not given	Not given	During
Lai et al 2020 [6]	COVID-19	IES-R	Median	HE and LE	20	22.5	17.0	During
Zhu et al 2020 [7]	COVID-19	IES-R	Not given	HE and LE	Not given	Not given	Not given	During
Xiao et al 2020 [1]	COVID-19	SASR	Mean	HE only	Not relevant	77.59 (29.53)	Not relevant	During
Matsuishi et al 2020 [15]	H1N1	IES-R	Mean	HE and LE	2.49 (6.63)	3.1 (8.0)	2.2 (5.9)	During
Lee et al 2018 [16]	MERS	IES-R	Mean	HE and LE	26.3 (19.09)	30.02 (19.55)	22.33 (17.70)	During
Bukhari et al 2016 [17]	MERS	IES	Not given	HE and LE	Not given	Not given	Not given	During
Chan & Huak 2004 (doctors) [18]	SARS	IES	Mean	HE and LE	Not given	15.1 (15.8)	15.4 (15.4)	During
Chan & Huak 2004 (nurses) [18]	SARS	IES	Mean	HE and LE	Not given	16.8 (12.9)	16.0 (14.6)	During
Koh et al 2005 [20]	SARS	IES	Median	HE and LE	16	Not given	Not given	During
C.S. Chen et al 2005 [13]	SARS	IES-R	Mean	HE and LE	17.8 (12.4)	19.1 (14.2)	13.7 (9.1)	During
Ho et al 2005 [21]	SARS	CIES-R	Not given	Undifferentiated	Not given	Not given	Not given	During
Chong et al 2004 [19]	SARS	IES	Mean	HE and LE	34.8 (19.7)	30.6 (15.8)	35.1 (20.0)	During
Styra et al 2008 [24]	SARS	IES-R	Not given	HE and LE	Not given	Not given	Not given	During
Su et al 2007 [22]	SARS	DTS-C	Mean	HE and LE	Not given	17.6 (11.9)	15.2 (12.5)	During
Lin et al 2007 [25]	SARS	DTS-C	Mean	HE and LE	Not given	26.03 (3.09)	16.04 (3.26)	During
Sim et al	SARS	IES-R	Mean	HE and LE	Not given	Not given	Not given	9 weeks after
Tham et al 2004 (doctors) [26]	SARS	IES	Median	Undifferentiated	9.5	Not relevant	Not relevant	6 months after
Tham et al 2004 (nurses) [26]	SARS	IES	Median	Undifferentiated	15	Not relevant	Not relevant	6 months after
McAlonan et al 2005 [10]	SARS	IES-R	Mean	HE and LE	Not given	17.2 (14.2)	9.3 (11.2)	During & 1 year
Maunder et al 2006 [28]	SARS	IES	Median	HE and LE	Not given	11	7	1-2 years after
Lancee et al 2008 [29]	SARS	IES	Mean	HE only	Not relevant	12.8 (10.3)	Not relevant	1-2 years after
Wu et al 2008/2009 [30] [31]	SARS	IES-R	Not given	HE and LE	Not given	Not given	Not given	3 years after

PTSD-SS = PTSD Self Report Scale; IES = Impact of Events Scale; IES-R = Impact of Events Scale – Revised; SASR = Stanford Acute Stress Reaction; CIES-R = Chinese Impact of Event Scale-Revised; Chinese version of the Davidson Trauma Scale (DTS-C)

Table S5. All studies reporting averages of validated PTSD symptom scales for healthcare responders for high risk epidemic and pandemic disease outbreaks

Table S6. Depression averages

<i>Study</i>	<i>Condition</i>	<i>Measure</i>	<i>Summary stat</i>	<i>Sample</i>	<i>Average (SD)</i>			<i>Timing</i>
					<i>Overall</i>	<i>High exposure</i>	<i>Low exposure</i>	
Zhu et al 2020 [7]	COVID-19	PHQ-9	Not given	HE and LE	Not given	Not given	Not given	During
Lai et al 2020 [6]	COVID-19	PHQ-9	Median	HE and LE	5.0	6.0	4.0	During
Chung & Yeung 2020 [32] ^a	COVID-19	PHQ-9	Mean	Undifferentiated	7.6 (No SD)	Not given	Not given	During
Liang et al [5] ^e	COVID-19	ZSRAD	Mean	HE and LE	Not given	30.05 (8.16)	34.09 (8.97)	During
Z. Liu et al 2020 [3]	COVID-19	ZSRAD	Not given	HE and LE	Not given	Not given	Not given	During
Sun et al [8] ^b	COVID-19	SCL-90-A	Mean	HE and LE	1.53 (0.79)	Not given	Not given	During
Ji et al 2017 [9] (Chinese staff)				HE only				During
Ji et al 2017 [9] (Sierra Leone staff)	Ebola	SCL-90-R-A	Mean		Not relevant	1.22 (0.31)	Not relevant	
			Mean	HE only				During
Chan & Huak 2004 (doctors) [18]	SARS	GHQ-28 D	Mean	HE and LE	Not given	0.6 (1.1)	0.4 (0.9)	During
Chan & Huak 2004 (nurses) [18]	SARS	GHQ-28 D	Mean	HE and LE	Not given	0.3 (0.6)	0.4 (1.0)	During
McAlonan et al 2005 [10]	SARS	DASS-21	Mean	HE and LE	Not given	4.9 (4.8)	2.2 (2.6)	During & 1 year
C.S. Chen et al 2005 [13]	SARS	SCL-90-R-A	Mean	HE and LE	0.65 (0.62)	0.64 (0.7)	0.55 (0.47)	During
R. Chen et al 2006 [11]	SARS	ZSRAD	Mean	HE Only	61	Not given	Not given	During, T1
Sim et al 2004 [23] ^d	SARS	GHQ-28 D	Mean	HE and LE	Not given	0.1 (0.5)	0.2 (0.9)	During
Su et al 2007 [22] ^c	SARS	BDI	Mean	HE and LE	Not given	10.2 (6.6)	6.7 (6.6)	During

PHQ-9 = Patient Health Questionnaire 9; ZSRAD = Zung Self-rating Depression Scale; SCL-90-A Symptom Checklist 90 Depression subscale; BDI = Beck Depression Inventory

Table S6. All studies reporting averages of validated anxiety depression scales for healthcare responders for high risk epidemic and pandemic disease outbreaks

References

1. Xiao H, Zhang Y, Kong D, Li S, Yang N. The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China. *Med Sci Monit.* 2020;26:e923549-1-e923549-8.
2. Liu C, Yang Y, Zhang XM, Xu X, Dou Q-L, Zhang W-W. The prevalence and influencing factors for anxiety in medical workers fighting COVID-19 in China: A cross-sectional survey. preprint. *Psychiatry and Clinical Psychology*; 2020. doi:10.1101/2020.03.05.20032003.
3. Liu Z, Han B, Jiang R, Huang Y, Ma C, Wen J, et al. Mental Health Status of Doctors and Nurses During COVID-19 Epidemic in China. SSRN Scholarly Paper. Rochester, NY: Social Science Research Network; 2020. <https://papers.ssrn.com/abstract=3551329>. Accessed 28 Mar 2020.
4. Huang JZ, Han MF, Luo TD, Ren AK, Zhou XP. Mental health survey of 230 medical staff in a tertiary infectious disease hospital for COVID-19. [Chinese]. *Zhonghua lao dong wei sheng zhi ye bing za zhi = Zhonghua laodong weisheng zhiyebing zazhi = Chinese journal of industrial hygiene and occupational diseases.* 2020;38:E001.
5. Liang Y, Chen M, Zheng X, Liu J. Screening for Chinese medical staff mental health by SDS and SAS during the outbreak of COVID-19. *Journal of Psychosomatic Research.* 2020;133:110102.
6. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open.* 2020;3:e203976–e203976.
7. Zhu Z, Xu S, Wang H, Liu Z, Wu J, Li G, et al. COVID-19 in Wuhan: Immediate Psychological Impact on 5062 Health Workers. *medRxiv.* 2020;:2020.02.20.20025338.
8. Sun N, Xing J, Xu J, Geng LS, Li QY. Study of the mental health status of medical personnel dealing with new coronavirus pneumonia. *medRxiv.* 2020;:2020.03.04.20030973.
9. Ji D, Ji Y-J, Duan X-Z, Li W-G, Sun Z-Q, Song X-A, et al. Prevalence of psychological symptoms among Ebola survivors and healthcare workers during the 2014-2015 Ebola outbreak in Sierra Leone: a cross-sectional study. *Oncotarget.* 2017;8:12784–91.
10. McAlonan GM, Lee AM, Cheung V, Wong JWS, Chua SE. “Psychological distress and negative appraisals in survivors of severe acute respiratory syndrome (SARS)” and “Severe acute respiratory syndrome (SARS) in Hong Kong in 2003: Stress and psychological impact among frontline healthcare workers”: Comment. *Psychological Medicine.* 2005;35:459–60.
11. Chen R, Chou K-R, Huang Y-J, Wang T-S, Liu S-Y, Ho L-Y. Effects of a SARS prevention programme in Taiwan on nursing staff’s anxiety, depression and sleep quality: A longitudinal survey. *Journal of Nursing Studies.* 2006;43:215–25.
12. Iancu I, Strous R, Poreh A, Kotler M, Chelben Y. Psychiatric Inpatients’ Reactions to the SARS Epidemic: An Israeli Survey. *Journal of Psychiatry and Related Sciences.* 2005;42:258–62.
13. Chen C-S, Wu H-Y, Yang P, Yen C-F. Psychological Distress of Nurses in Taiwan Who Worked During the Outbreak of SARS. *PS.* 2005;56:76–9.

14. Poon E, Liu KS, Cheong DL, Lee CK, Yam LYC, Tang WN. Impact of severe respiratory syndrome on anxiety levels of front-line health care workers. *Hong Kong Med J*. 2004;10:325–30.
15. Matsuishi K, Kawazoe A, Imai H, Ito A, Mouri K, Kitamura N, et al. Psychological impact of the pandemic (H1N1) 2009 on general hospital workers in Kobe. *Psychiatry and Clinical Neurosciences*. 2012;66:353–60.
16. Lee SM, Kang WS, Cho A-R, Kim T, Park JK. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. *Comprehensive Psychiatry*. 2018;:123–7.
17. Bukhari EE, Temsah MH, Aleyadhy AA, Alrabiaa AA, Alhboob AA, Jamal AA, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak perceptions of risk and stress evaluation in nurses. *The Journal of Infection in Developing Countries*. 2016;10:845–50.
18. Chan AOM, Huak CY. Psychological impact of the 2003 severe acute respiratory syndrome outbreak on health care workers in a medium size regional general hospital in Singapore. *Occup Med (Lond)*. 2004;54:190–6.
19. Chong M-Y, Wang W-C, Hsieh W-C, Lee C-Y, Chiu N-M, Yeh W-C, et al. Psychological impact of severe acute respiratory syndrome on health workers in a tertiary hospital. *The British Journal of Psychiatry*. 2004;185:127–33.
20. Koh D, Lim MK, Chia SE, Ko SM, Qian F, Ng V, et al. Risk Perception and Impact of Severe Acute Respiratory Syndrome (SARS) on Work and Personal Lives of Healthcare Workers in Singapore What Can We Learn? *Medical Care*. 2005;43:676–82.
21. Ho SMY, Kwong-Lo RSY, Mak CWY, Wong JS. Fear of Severe Acute Respiratory Syndrome (SARS) Among Health Care Workers. *Journal of Consulting and Clinical Psychology*. 2005;73:344–9.
22. Su T-P, Lien T-C, Yang C-Y, Su YL, Wang J-H, Tsai S-L, et al. Prevalence of psychiatric morbidity and psychological adaptation of the nurses in a structured SARS caring unit during outbreak: A prospective and periodic assessment study in Taiwan. *Journal of Psychiatric Research*. 2007;41:119–30.
23. Sim K, Chong PN, Chan YH, Soon WSW. Severe Acute Respiratory Syndrome-Related Psychiatric and Posttraumatic Morbidities and Coping Responses in Medical Staff Within a Primary Health Care Setting in Singapore. *Journal of Clinical Psychiatry*. 2004;65:1120–7.
24. Styra R, Hawryluck L, Robinson S, Kasapinovic S, Fones C, Gold WL. Impact on health care workers employed in high-risk areas during the Toronto SARS outbreak. *Journal of Psychosomatic Research*. 2008;64:177–83.
25. Lin C-Y, Peng Y-C, Wu Y-H, Chang J, Chan C-H, Yang D-Y. The psychological effect of severe acute respiratory syndrome on emergency department staff. *Emergency Medicine Journal*. 2007;24:12–7.
26. Tham K-Y, Tan Y, Loh O, Tan W, Ong M, Tang H. Psychiatric morbidity among emergency department doctors and nurses after the SARS outbreak. *Annals of the Academy of Medicine, Singapore*. 2004;33:S78-9.
27. Phua DH, Tang HK, Tham KY. Coping Responses of Emergency Physicians and Nurses to the 2003 Severe Acute Respiratory Syndrome Outbreak. *Academic Emergency Medicine*. 2005;12:322–8.

28. Maunder RG, Lancee WJ, Balderson KE, Bennett JP, Borgundvaag B, Evans S, et al. Long-term Psychological and Occupational Effects of Providing Hospital Healthcare during SARS Outbreak. *Emerg Infect Dis.* 2006;12:1924–32.
29. Lancee WJ, Maunder RG, Goldbloom DS. Prevalence of Psychiatric Disorders Among Toronto Hospital Workers One to Two Years After the SARS Outbreak. *PS.* 2008;59:91–5.
30. Wu P, Liu X, Fang Y, Fan B, Fuller CJ, Guan Z, et al. Alcohol Abuse/Dependence Symptoms Among Hospital Employees Exposed to a SARS Outbreak. *Alcohol Alcohol.* 2008;43:706–12.
31. Wu P, Fang Y, Guan Z, Fan B, Kong J, Yao Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: Exposure, risk perception, and altruistic acceptance of risk. *Journal of Psychiatry.* 2009;54:302–11.
32. Chung J, Yeung W. Staff Mental Health Self-Assessment During the COVID-19 Outbreak. *East Asian Arch Psychiatry.* 2020;30:34.
33. Li L, Wan C, Ding R, Liu Y, Chen J, Wu Z, et al. Mental distress among Liberian medical staff working at the China Ebola Treatment Unit: a cross sectional study. *Health and Quality of Life Outcomes.* 2015;13:156.