

Supplementary file

Models – main analysis

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 1)

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.000	0.007	-0.06	0.949	0.985	1.014	
Aged 50-79 (%)	1.031	0.019	1.69	0.091	0.995	1.069	*
Aged 80+ (%)	0.917	0.064	-1.25	0.212	0.800	1.051	
Population density (10,000 pop/km ²)	1.157	0.188	0.90	0.368	0.842	1.591	
BAME (%)	1.014	0.004	3.77	<0.001	1.007	1.022	***
Constant	0.000	0.000	-17.01	<0.001	0.000	0.000	***
Constant	-1.999	0.126	.b	.b	-2.245	-1.752	
Mean dependent var		110.170	SD dependent var			88.959	
Pseudo r-squared		0.027	Number of obs			147.000	
Chi-square		40.100	Prob > chi2			0.000	
Akaike crit. (AIC)		1468.881	Bayesian crit. (BIC)			1489.814	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 2)

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.018	0.009	2.13	0.033	1.001	1.036	**
Aged 50-79 (%)	1.002	0.017	0.14	0.890	0.969	1.037	
Aged 80+ (%)	0.936	0.067	-0.93	0.355	0.813	1.077	
Population density (10,000 pop/km ²)	1.183	0.203	0.98	0.326	0.846	1.655	
Deprivation (%)	1.024	0.009	2.70	0.007	1.006	1.042	***
Constant	0.000	0.000	-15.40	<0.001	0.000	0.000	***
Constant	-1.945	0.125	.b	.b	-2.190	-1.701	
Mean dependent var		110.170	SD dependent var			88.959	
Pseudo r-squared		0.022	Number of obs			147.000	
Chi-square		33.158	Prob > chi2			0.000	
Akaike crit. (AIC)		1475.823	Bayesian crit. (BIC)			1496.756	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 3)

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.010	0.008	1.24	0.215	0.994	1.027	
Aged 50-79 (%)	1.030	0.018	1.64	0.100	0.994	1.067	
Aged 80+ (%)	0.937	0.064	-0.95	0.342	0.820	1.071	
Population density (10,000 pop/km ²)	1.068	0.172	0.41	0.683	0.779	1.465	
BAME (%)	1.014	0.004	3.74	<0.001	1.007	1.021	***
Deprivation (%)	1.022	0.009	2.64	0.008	1.006	1.039	***
Constant	0.000	0.000	-16.61	<0.001	0.000	0.000	***
Constant	-2.048	0.127	.b	.b	-2.296	-1.801	
Mean dependent var		110.170	SD dependent var			88.959	
Pseudo r-squared		0.031	Number of obs			147.000	
Chi-square		46.991	Prob > chi2			0.000	
Akaike crit. (AIC)		1463.991	Bayesian crit. (BIC)			1487.914	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Statistical model

Equation for negative binomial regression Model 3

$$\log(Y_i) = \beta_0 + \beta_1 Duration_i + \beta_2 Age50_79_i + \beta_3 Age80plus_i + \beta_4 PopDensity_i + \beta_5 BAME_i + \beta_6 Deprivation_i + \log(population_i)$$

Where Y_i is the number of COVID19 deaths for local authority i . *Duration* is the number of days from when each local authority had at least 10 laboratory confirmed COVID19 cases to the 22nd April 2020. *Age50_79* is the percent of the population aged 50-79 years. *Age80plus* is the percent of the population aged 80+ years. *PopDensity* is the number of people per square kilometre expressed in terms of 10,000 people/km². *BAME* is the percent of the population who classify their ethnic group as: Black, Asian, Mixed or Other. *Deprivation* is the percent of the population experiencing deprivation relating to low income. *Population* is an offset variable with a parameter estimate constrained to 1.

Robustness test – excluding local authorities in London

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England – excluding local authorities in London

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.013	0.009	1.48	0.139	0.996	1.030	
Aged 50-79 (%)	1.068	0.026	2.72	0.006	1.019	1.120	***
Aged 80+ (%)	0.844	0.067	-2.14	0.032	0.723	0.986	**
Population density (10,000 pop/km ²)	2.266	0.993	1.87	0.062	0.959	5.350	*
BAME (%)	1.013	0.006	2.11	0.035	1.001	1.025	**
Deprivation (%)	1.023	0.011	2.14	0.032	1.002	1.044	**
Constant	0.000	0.000	-14.82	<0.001	0.000	0.000	***
Constant	-2.054	0.144	.b	.b	-2.337	-1.771	
Mean dependent var		107.200	SD dependent var			97.701	
Pseudo r-squared		0.027	Number of obs			115.000	
Chi-square		30.783	Prob > chi2			0.000	
Akaike crit. (AIC)		1131.544	Bayesian crit. (BIC)			1153.503	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Robustness test – using linear regression model

Multivariable linear regression model for COVID19 mortality rate per 100,000 population for local authorities in England

	Coef.	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	0.378	0.234	1.61	0.109	-0.085	0.841	
Aged 50-79 (%)	0.904	0.501	1.80	0.073	-0.086	1.894	*
Aged 80+ (%)	-1.563	1.973	-0.79	0.429	-5.464	2.338	
Population density (10,000 pop/km ²)	0.879	4.463	0.20	0.844	-7.946	9.703	
BAME (%)	0.464	0.103	4.49	<0.001	0.260	0.669	***
Deprivation (%)	0.683	0.245	2.79	0.006	0.199	1.167	***
Constant	-22.554	16.053	-1.41	0.162	-54.292	9.184	
Mean dependent var		30.346	SD dependent var			12.820	
R-squared		0.313	Number of obs			147.000	
F-test		10.630	Prob > F			0.000	
Akaike crit. (AIC)		1124.974	Bayesian crit. (BIC)			1145.907	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic

Robustness test – association between the proportion of the population from Black ethnic backgrounds and COVID19 mortality

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.013	0.009	1.56	0.120	0.997	1.030	
Aged 50-79 (%)	1.009	0.017	0.54	0.589	0.976	1.044	
Aged 80+ (%)	0.960	0.068	-0.58	0.561	0.836	1.102	
Population density (10,000 pop/km ²)	1.007	0.173	0.04	0.969	0.719	1.410	
Proportion Black ethnicity (%)	1.026	0.009	2.73	0.006	1.007	1.044	***
Deprivation (%)	1.024	0.009	2.80	0.005	1.007	1.041	***
Constant	0.000	0.000	-16.13	0.000	0.000	0.000	***
Constant	-2.001	0.126	.b	.b	-2.247	-1.755	
Mean dependent var		110.170	SD dependent var			88.959	
Pseudo r-squared		0.027	Number of obs			147.000	
Chi-square		40.671	Prob > chi2			0.000	
Akaike crit. (AIC)		1470.311	Bayesian crit. (BIC)			1494.234	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

IRR = incident rate ratio

Robustness tests – using ONS mortality figures up to 10th April 2020

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 1) – using ONS mortality data

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.018	0.008	2.13	0.033	1.001	1.035	**
Aged 50-79 (%)	1.020	0.020	0.98	0.325	0.981	1.060	
Aged 80+ (%)	0.970	0.074	-0.40	0.693	0.836	1.126	
Population density (10,000 pop/km ²)	1.409	0.255	1.90	0.058	0.988	2.008	*
BAME (%)	1.017	0.004	4.30	<0.001	1.009	1.025	***
Constant	0.000	0.000	-18.24	<0.001	0.000	0.000	***
Constant	-1.849	0.131	.b	.b	-2.105	-1.592	
Mean dependent var		67.068	SD dependent var			52.276	
Pseudo r-squared		0.060	Number of obs			147.000	
Chi-square		84.585	Prob > chi2			0.000	
Akaike crit. (AIC)		1340.725	Bayesian crit. (BIC)			1361.658	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 2) – using ONS mortality data

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.038	0.010	3.81	<0.001	1.018	1.059	***
Aged 50-79 (%)	0.987	0.019	-0.69	0.491	0.950	1.025	
Aged 80+ (%)	0.984	0.079	-0.20	0.841	0.840	1.152	
Population density (10,000 pop/km ²)	1.531	0.302	2.16	0.031	1.040	2.254	**
Deprivation (%)	1.021	0.010	2.12	0.034	1.002	1.041	**
Constant	0.000	0.000	-15.36	<0.001	0.000	0.000	***
Constant	-1.736	0.127	.b	.b	-1.985	-1.486	
Mean dependent var		67.068	SD dependent var			52.276	
Pseudo r-squared		0.050	Number of obs			147.000	
Chi-square		70.985	Prob > chi2			0.000	
Akaike crit. (AIC)		1354.325	Bayesian crit. (BIC)			1375.258	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio

Multivariable negative binomial regression model for COVID19 mortality rate for local authorities in England (Model 3) – using ONS mortality data

	IRR	St.Err.	t-value	p-value	[95% Conf	Interval]	Sig
Duration (days)	1.028	0.010	2.96	0.003	1.009	1.047	***
Aged 50-79 (%)	1.019	0.020	0.98	0.329	0.981	1.059	
Aged 80+ (%)	0.988	0.075	-0.15	0.878	0.853	1.146	
Population density (10,000 pop/km ²)	1.302	0.236	1.46	0.146	0.913	1.856	
BAME (%)	1.017	0.004	4.34	<0.001	1.009	1.025	***
Deprivation (%)	1.020	0.009	2.20	0.028	1.002	1.039	**
Constant	0.000	0.000	-17.11	<0.001	0.000	0.000	***
Constant	-1.883	0.131	.b	.b	-2.140	-1.626	
Mean dependent var		67.068	SD dependent var			52.276	
Pseudo r-squared		0.063	Number of obs			147.000	
Chi-square		89.382	Prob > chi2			0.000	
Akaike crit. (AIC)		1337.928	Bayesian crit. (BIC)			1361.852	

*** $p < 0.01$, ** $p < 0.05$, * $p < 0.1$

BAME = Black, Asian and Minority Ethnic; IRR = incident rate ratio