Supplemental Tables

Supplementary Table 1: Number of blood culture sets collected per patient with "not tested" SARS-CoV-2 status

| | Jan-Feb 2019 (N=29019) | Mar 1-15, 2019 (N=7601) | Mar 16-31, 2019 (N=7688) | Jan-Feb 2020 (N=28593) | Mar 1-15, 2020 (N=5942) | Mar 16-31, 2020 (N=1896) | Total (N=80739 |
|-------------------|------------------------------|-------------------------------|--------------------------------|------------------------------|-------------------------------|--------------------------------|-------------------|
| BC per patient | | | | | | | |
| 1 | $1612 \\ (5.6\%)$ | 378 (5.0%) | 433~(5.6%) | $1689 \\ (5.9\%)$ | 349~(5.9%) | 166 (8.8%) | 4627 $(5.7%)$ |
| 2 | 9253 (31.9%) | $2302 \\ (30.3\%)$ | 2334 (30.4%) | 9438 (33.0%) | $2016 \\ (33.9\%)$ | 685 (36.1%) | 26028 (32.2%) |
| 3-5 | 5834 (20.1%) | 1508 (19.8%) | 1479 (19.2%) | 6238 (21.8%) | 1071 (18.0%) | 363 (19.1%) | 16493 (20.4%) |
| 6-10 | 6010 (20.7%) | 1638 (21.5%) | 1598 (20.8%) | 6023 (21.1%) | 1309 (22.0%) | 339 (17.9%) | 16917 (21.0%) |
| 11-30 | 5214 (18.0%) | 1540 (20.3%) | 1607 (20.9%) | 4695 (16.4%) | 1128 (19.0%) | 331 (17.5%) | 14515 (18.0%) |
| >30 | $1096 \\ (3.8\%)$ | 235 (3.1%) | 237 (3.1%) | 510 (1.8%) | 69 (1.2%) | 12~(0.6%) | 2159 (2.7%) |

| | Negative $(N=2986)$ | Positive $(N=4478)$ | Total (N= 7464) | p value |
|----------------|---------------------|---------------------|--------------------|---------------|
| BC per patient | | | | $< 0.001^{1}$ |
| 1 | 114 (3.8%) | 127 (2.8%) | 241 (3.2%) | |
| 2 | 896(30.0%) | 2194 (49.0%) | 3090(41.4%) | |
| 3-5 | 687(23.0%) | 1189(26.6%) | 1876(25.1%) | |
| 6-10 | 637~(21.3%) | 829 (18.5%) | 1466 (19.6%) | |
| 11-30 | 597(20.0%) | 139(3.1%) | 736(9.9%) | |
| >30 | 55 (1.8%) | 0 (0.0%) | 55~(0.7%) | |

Supplementary Table 2: Number of BC sets collected per patient tested for SARS-CoV-2

1. Pearson's Chi-squared test

| | Not tested | Negative | Positive | Total |
|---|----------------------|--------------------------|--------------------|---------------------|
| | (N=6260) | (N=245) | (N=201) | (N=6706) |
| Organism Name | | | | |
| Abiotrophia defectiva | 3~(0.0%) | 0 (0.0%) | 0 (0.0%) | 3 (0.0%) |
| Achromobacter xylosoxidans | 6(0.1%) | 0(0.0%) | 0(0.0%) | 6(0.1%) |
| Achromobacter xylosoxidans subsp | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| xylosoxidans | () | - (, •) | - (, •) | () |
| Acinetobacter baumannii | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6(0.1%) |
| Acinetobacter baumannii complex | 15(0.2%) | 0 (0.0%) | 0 (0.0%) | 15 (0.2%) |
| Acinetobacter baumannii/nosocomialis group | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Acinetobacter junii | $\frac{1}{3}(0.0\%)$ | 0 (0.0%) | 0 (0.0%) | 3(0.0%) |
| Acinetobacter pittii | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Acinetobacter radioresistens | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Acinetobacter species | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Acinetobacter ursingii | 5(0.1%) | 0 (0.0%) | 0(0.0%) | 5(0.1%) |
| Actinobaculum schaalii | 1 (0.0%) | 0 (0.0%) | 0(0.0%) | 1 (0.0%) |
| Actinomyces meyeri | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Actinomyces neuii | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Actinomyces odontolyticus | 6 (0.1%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 6 (0.1%) |
| Actinomyces oris | 3 (0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 3 (0.0%) |
| Actinomyces radicidentis | 1 (0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Actinomyces species | 2(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 2(0.0\%) |
| Actinomyces species Actinomyces turicensis | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 1 (0.5%) | 2(0.0%) 2(0.0\%) |
| Actinomyces turicensis Aerococcus urinae | 1(0.0%) 1(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 1(0.5%) 1(0.5%) | 2(0.0%) 2(0.0\%) |
| Aerococcus viridans | 4 (0.1%) | · · · · · | · · · · | · · · · |
| | · / | 0 (0.0%) | 0 (0.0%) | 4 (0.1%) |
| Alcaligenes species | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Alloiococcus species | 3(0.0%) | 0 (0.0%) 0 (0.0%) | 0 (0.0%) | 3(0.0%) |
| Anaerococcus prevotii | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Anaerococcus vaginalis | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Arcanobacterium haemolyticum | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Arthrobacter cumminsii | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Atopobium parvulum | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1(0.0%) |
| Bacillus cereus | 9(0.1%) | 1 (0.4%) | 0 (0.0%) | 10 (0.1%) |
| Bacillus megaterium | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1(0.0%) |
| Bacillus non-anthracis | 25 (0.4%) | 0 (0.0%) | 9(4.5%) | 34 (0.5%) |
| Bacillus species | 28 (0.4%) | 0 (0.0%) | 0 (0.0%) | 28 (0.4%) |
| Bacteroides fragilis | 39(0.6%) | 1 (0.4%) | 0 (0.0%) | 40 (0.6%) |
| Bacteroides nordii | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Bacteroides ovatus | 5(0.1%) | 0 (0.0%) | 0 (0.0%) | 5(0.1%) |
| Bacteroides thetaiotaomicron | 9(0.1%) | 2(0.8%) | 0 (0.0%) | 11 (0.2%) |
| Bacteroides uniformis | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Bacteroides vulgatus | 1 (0.0%) | 2 (0.8%) | 0 (0.0%) | 3 (0.0%) |
| Bifidobacterium longum | 1 (0.0%) | 0~(0.0%) | 0 (0.0%) | 1 (0.0%) |
| $Brevibacterium\ casei$ | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0%) |
| $Brevibacterium\ species$ | 3~(0.0%) | 0~(0.0%) | 0~(0.0%) | 3 (0.0%) |
| Burkholderia cepacia complex | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Burkholderia multivorans | 14 (0.2%) | 0~(0.0%) | $0 \ (0.0\%)$ | 14 (0.2%) |
| $Campylobacter\ coli$ | 2~(0.0%) | 0~(0.0%) | 0~(0.0%) | 2~(0.0%) |
| Campylobacter fetus | 2 (0.0%) | 0~(0.0%) | 0~(0.0%) | 2~(0.0%) |
| Candida albicans | 75~(1.2%) | 3~(1.2%) | 3~(1.5%) | 81~(1.2%) |
| Candida auris | 5~(0.1%) | 0~(0.0%) | 0~(0.0%) | 5~(0.1%) |
| Candida bracarensis | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |

Supplementary Table 3: Microorganisms isolated per blood culture set by SARS-CoV-2 status

| | Not tested $(N=6260)$ | Negative (N=245) | Positive (N=201) | Total $(N=6706)$ |
|---|-----------------------|--------------------------|--------------------------|----------------------|
| Candida dubliniensis | 4 (0.1%) | 0 (0.0%) | 0 (0.0%) | 4 (0.1%) |
| Candida glabrata | 72(1.2%) | 5(0.0%) 5(2.0%) | 1 (0.5%) | 78 (1.2%) |
| Candida guilliermondii | 2(0.0%) | 0(0.0%) | 0 (0.0%) | 2 (0.0%) |
| Candida krusei | 2(0.0%) 2(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 2(0.0%) 2(0.0%) |
| Candida lusitaniae | 8 (0.1%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | $\frac{2}{8}(0.0\%)$ |
| Candida magnoliae | 1 (0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Candida orthopsilosis | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Candida parapsilosis complex | 26 (0.4%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 0(0.0%) 0(0.0%) | 1(0.07) 26 (0.4% |
| Candida parapsilosis complex Candida parapsilosis | 14 (0.2%) | 1 (0.4%) | 0(0.0%) 0(0.0%) | 15 (0.2%) |
| Candida pararugosa | 14(0.2%) 1(0.0%) | 1(0.4%) 0(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | · · · |
| - • | () | · · · · · | · / | 1 (0.0%) |
| Candida tropicalis | 27 (0.4%) | 1 (0.4%) | 0 (0.0%) | 28 (0.4%) |
| Capnocytophaga sputigena | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Cellulomonas species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Citrobacter amalonaticus | 3(0.0%) | 0 (0.0%) | 0 (0.0%) | 3 (0.0%) |
| Citrobacter braakii | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Citrobacter freundii | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Citrobacter freundii complex | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1%) |
| Citrobacter koseri | 6 (0.1%) | 2(0.8%) | 0 (0.0%) | 8 (0.1% |
| Clostridium baratii | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Clostridium bifermentans | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Clostridium butyricum | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Clostridium cadaveris | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| $Clostridium\ clostridio forme$ | 7~(0.1%) | 0~(0.0%) | 0~(0.0%) | 7 (0.1% |
| Clostridium innocuum | 1~(0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0% |
| Clostridium limosum | 2~(0.0%) | 0~(0.0%) | 0~(0.0%) | 2 (0.0% |
| $Clostridium \ perfringens$ | 5~(0.1%) | 0~(0.0%) | 0~(0.0%) | 5 (0.1% |
| Clostridium ramosum | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0% |
| Clostridium septicum | 4 (0.1%) | 0~(0.0%) | 0~(0.0%) | 4 (0.1% |
| Clostridium species | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0% |
| Comamonas kerstersii | 2 (0.0%) | 0~(0.0%) | 0~(0.0%) | 2 (0.0% |
| Comamonas testosteroni | 0 (0.0%) | 0~(0.0%) | 1 (0.5%) | 1 (0.0% |
| Corynebacterium afermentans | 8 (0.1%) | 0~(0.0%) | 0~(0.0%) | 8 (0.1% |
| Corynebacterium amycolatum | 6(0.1%) | 0(0.0%) | 0(0.0%) | 6(0.1%) |
| Corynebacterium aurimucosum | 6(0.1%) | 0(0.0%) | 0(0.0%) | 6(0.1%) |
| Corynebacterium confusum | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Corynebacterium coyleae | 5(0.1%) | 0(0.0%) | 0(0.0%) | 5(0.1%) |
| Corynebacterium jeikeium | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1 (0.0% |
| Corynebacterium matruchotii | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Corynebacterium minutissimum | 3(0.0%) | 0 (0.0%) | 0 (0.0%) | 3 (0.0% |
| Corynebacterium propinquum | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Corynebacterium pseudodiphtheriticum | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Corynebacterium striatum | 7 (0.1%) | 4(1.6%) | 0 (0.0%) | 11 (0.2% |
| Corynebacterium tuberculostearicum | 1(0.0%) | 0 (0.0%) | 0(0.0%) | 1 (0.0% |
| Corynebacterium tuscaniense | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Corynebacterium tuscamense Corynebacterium xerosis | 1(0.0%) 1(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Corynebacterium xerosis | 1(0.0%) 89 (1.4%) | 3(1.2%) | 5(0.0%) 5(2.5%) | 97 (1.4) |
| | · · · · | · / | · / | |
| Corynebacterium species, not <i>C. jeikeium</i> | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Cutibacterium (Propionibacterium) acnes | 51 (0.8%) | 4(1.6%) | 0 (0.0%) | 55 (0.8% |
| Dermabacter hominis | 12 (0.2%) | 0 (0.0%) | 0 (0.0%) | 12 (0.2%) |
| Desulfovibrio species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Dialister pneumosintes | 1~(0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0% |

| | Not tested (N=6260) | Negative (N=245) | Positive (N=201) | Total (N=6706) |
|---|------------------------|---------------------------------------|---------------------|-------------------|
| Dolosigranulum pigrum | 1 (0.0%) | 0 (0.0%) | 1 (0.5%) | 2(0.0%) |
| Dolosigranulum species | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Eggerthella lenta | 3(0.0%) | 1(0.4%) | 0(0.0%) | 4(0.1%) |
| Eikenella corrodens | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Eikenella species | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Elizabethkingia meningoseptica | 4(0.1%) | 0(0.0%) | 0(0.0%) | 4(0.1%) |
| Eneterobacter | 16(0.3%) | 1(0.4%) | 1(0.5%) | 18(0.3%) |
| Enterobacter cloacae ssp. cloacae | 13(0.2%) | 0(0.0%) | 0(0.0%) | 13(0.2%) |
| Enterobacter cloacae | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Enterobacter cloacae complex | 79(1.3%) | 0 (0.0%) | 5(2.5%) | 84~(1.3%) |
| Enterobacter cloacae complex (absence of KPC | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1%) |
| resistance gene) | | , , , , , , , , , , , , , , , , , , , | . , | . , |
| Enterococcus avium | 3 (0.0%) | 0 (0.0%) | 0 (0.0%) | 3~(0.0%) |
| Enterococcus casseliflavus | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Enterococcus faecalis | 232~(3.7%) | 4(1.6%) | 2(1.0%) | 238 |
| | | | | (3.5%) |
| Enterococcus faecalis, Vancomycin-Resistant | 13 (0.2%) | 0 (0.0%) | 0 (0.0%) | 13(0.2%) |
| Enterococcus faecalis, Vancomycin-Susceptible | 22(0.4%) | 0(0.0%) | 0(0.0%) | 22(0.3%) |
| Enterococcus faecium | 38(0.6%) | 3(1.2%) | 0(0.0%) | 41 (0.6%) |
| Enterococcus faecium, Vancomycin-Resistant | 101 (1.6%) | 11 (4.5%) | 1 (0.5%) | 113 (1.7%) |
| Enterococcus faecium, Vancomycin-Susceptible | 5(0.1%) | 0 (0.0%) | 0 (0.0%) | 5(0.1%) |
| Enterococcus gallinarum | 17(0.3%) | 1(0.4%) | 0(0.0%) | 18(0.3%) |
| Enterococcus hirae | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1 (0.0%) |
| Enterococcus italicus | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Enterococcus raffinosus | 3(0.0%) | 0(0.0%) | 0(0.0%) | 3(0.0%) |
| Enterococcus species | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Escherichia coli | 721 (11.5%) | 25 (10.2%) | 8 (4.0%) | 754 $(11.2%)$ |
| Escherichia coli - ESBL producer | 30~(0.5%) | 0 (0.0%) | 0 (0.0%) | 30(0.4%) |
| Escherichia coli (absence of KPC resistance | 123~(2.0%) | 4(1.6%) | 2(1.0%) | 129 |
| gene) | | | | (1.9%) |
| Facklamia hominis | 5(0.1%) | 0 (0.0%) | 0 (0.0%) | 5(0.1%) |
| Finegoldia magna | 13~(0.2%) | 0~(0.0%) | 0~(0.0%) | 13~(0.2%) |
| Flavonifractor plautii | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1~(0.0%) |
| Fusobacterium mortiferum | 1~(0.0%) | 0~(0.0%) | 0~(0.0%) | 1~(0.0%) |
| Fusobacterium nucleatum | 9~(0.1%) | 0~(0.0%) | 0~(0.0%) | 9~(0.1%) |
| Fusobacterium species | 3(0.0%) | 0~(0.0%) | 2~(1.0%) | 5~(0.1%) |
| Fusobacterium varium | 1~(0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0%) |
| Gemella haemolysans | 3(0.0%) | 0~(0.0%) | 0~(0.0%) | 3~(0.0%) |
| $Gemella\ morbillorum$ | 3(0.0%) | 0~(0.0%) | 0~(0.0%) | 3~(0.0%) |
| $Geotrichum\ capitatum$ | 7~(0.1%) | 0~(0.0%) | 0~(0.0%) | 7~(0.1%) |
| $Globicatella\ sanguinis$ | 1 (0.0%) | 0~(0.0%) | $1 \ (0.5\%)$ | 2~(0.0%) |
| Gordonia aichiensis | 1~(0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0%) |
| $Granulicatella \ adiacens$ | 11~(0.2%) | 0~(0.0%) | $1 \ (0.5\%)$ | $12 \ (0.2\%)$ |
| Haemophilus influenzae | 24 (0.4%) | 2 (0.8%) | 0 (0.0%) | 26 (0.4%) |
| Haemophilus parainfluenzae | 8 (0.1%) | $0 \ (0.0\%)$ | 0 (0.0%) | 8 (0.1%) |
| Halomonas species | 1 (0.0%) | 0~(0.0%) | $0 \ (0.0\%)$ | 1 (0.0%) |
| Helcococcus sueciensis | 1 (0.0%) | 0~(0.0%) | $0 \ (0.0\%)$ | 1 (0.0%) |
| Klebsiella aerogenes | 28~(0.4%) | $0 \ (0.0\%)$ | 0 (0.0%) | 28 (0.4%) |
| Klebsiella oxytoca | 21~(0.3%) | 0 (0.0%) | $0 \ (0.0\%)$ | $21 \ (0.3\%)$ |

| | Not tested $(N=6260)$ | Negative (N=245) | Positive (N=201) | $\begin{array}{c} \text{Total} \\ (N=6706) \end{array}$ |
|---|-----------------------|--------------------------|---------------------|---|
| Klebsiella oxytoca, absence of KPC resistance | 2 (0.0%) | 0 (0.0%) | 0 (0.0%) | 2 (0.0%) |
| gene | | | | |
| Klebsiella pneumoniae | 221 (3.5%) | 14 (5.7%) | 6 (3.0%) | 241 (3.6%) |
| Klebsiella pneumoniae complex | 15~(0.2%) | 0 (0.0%) | $0 \ (0.0\%)$ | 15 (0.2%) |
| Klebsiella pneumoniae subsp pneumoniae | 71 (1.1%) | 2 (0.8%) | $0 \ (0.0\%)$ | 73 (1.1%) |
| Klebsiella pneumoniae, ESBL producer | 4(0.1%) | 1(0.4%) | 0(0.0%) | 5(0.1%) |
| Klebsiella pneumoniae, absence of KPC | 46(0.7%) | 2(0.8%) | 0(0.0%) | 48 (0.7%) |
| resistance gene | | | | |
| Klebsiella pneumoniae, Carbapenem resistant | 1 (0.0%) | 0~(0.0%) | 0 (0.0%) | 1 (0.0%) |
| Klebsiella pneumoniae, Carbapenem resistant | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1%) |
| Klebsiella variicola | 7(0.1%) | 0(0.0%) | 0(0.0%) | 7(0.1%) |
| Kluyvera cryocrescens | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Kluyvera species | 3(0.0%) | 0(0.0%) | 0(0.0%) | 3(0.0%) |
| Kocuria kristinae | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Kocuria marina | 0(0.0%) | 0(0.0%) | 1(0.5%) | 1 (0.0%) |
| Kocuria rhizophila | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1(0.0%) |
| Kocuria species | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Kocuria varians | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Lactobacillus delbruekii | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Lactobacillus fermentum | 5(0.1%) | 0 (0.0%) | 0(0.0%) | $\frac{2}{5}(0.0\%)$ |
| Lactobacillus gasseri | 2(0.0%) | 0 (0.0%) | 0(0.0%) 0(0.0%) | 2(0.0%) |
| Lactobacillus paracasei | 2(0.0%) 2(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 2(0.0%) 2(0.0\%) |
| Lactobacillus rhamnosus | 4 (0.1%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 4 (0.1%) |
| Lactobacillus salivarius | 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Lactobacillus | 6 (0.1%) | 0(0.0%) 0(0.0%) | 2(1.0%) | 8 (0.1%) |
| Lactococcus lactis | 1 (0.0%) | 0(0.0%) 0(0.0%) | 2(1.0%) 0(0.0\%) | 1 (0.0%) |
| Leifsonia aquatica | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Leptotrichia trevisanii | 0 (0.0%) | 1 (0.4%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Leuconostoc lactis | 1 (0.0%) | $1(0.470) \\ 0(0.0\%)$ | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| | 1(0.0%) 1(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Listeria monocytogenes | () | $0 (0.0\%) \\ 0 (0.0\%)$ | 0(0.0%) 0(0.0%) | 1(0.0%) 2(0.0%) |
| Lodderomyces elongisporus | 2(0.0%) | · / | (/ | · · · · |
| Meyerozyma species Mianaka atamiana la ati aum | 4 (0.1%) | 0 (0.0%) 0 (0.0%) | 0 (0.0%) | 4 (0.1%) |
| Microbacterium lacticum | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Microbacterium paraoxydans | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Microbacterium testaceum | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Microbacterium species | 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Micrococcus luteus | 36 (0.6%) | 2(0.8%) | 5(2.5%) | 43 (0.6%) |
| Micrococcus species | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1%) |
| Mogibacterium species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Moraxella catarrhalis | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Moraxella nonliquefaciens | 1 (0.0%) | 1 (0.4%) | 0 (0.0%) | 2(0.0%) |
| Moraxella osloensis | 4 (0.1%) | 0 (0.0%) | 1 (0.5%) | 5(0.1%) |
| Moraxella species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Morganella morganii | 16 (0.3%) | 0 (0.0%) | 0 (0.0%) | 16 (0.2%) |
| Neisseria flavescens | 3(0.0%) | 0 (0.0%) | 0 (0.0%) | 3 (0.0%) |
| Neisseria sicca | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Neisseria subflava | 1 (0.0%) | 0~(0.0%) | $0 \ (0.0\%)$ | 1 (0.0%) |
| Paenibacillus species | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0%) |
| Paenibacillus urinalis | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1 (0.0%) |
| Pantoea agglomerans group | 3(0.0%) | 0~(0.0%) | 0~(0.0%) | 3~(0.0%) |

| | Not tested $(N=6260)$ | Negative $(N=245)$ | Positive (N=201) | Total $(N=6706)$ |
|--|-------------------------|--------------------------|--------------------------|----------------------|
| Pantoea dispersa | 2 (0.0%) | 0 (0.0%) | 0 (0.0%) | 2 (0.0% |
| Pantoea septica | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Pantoea species | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | $\frac{2}{2}(0.0\%)$ |
| Parabacteroides distasonis | $\frac{2}{3}(0.0\%)$ | 0(0.0%) | 0(0.0%) 0(0.0%) | $\frac{2}{3}(0.0\%)$ |
| Parvimonas micra | 6 (0.1%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 6 (0.0%) |
| Pediococcus pentosaceus | 3 (0.0%) | 0 (0.0%) | 0 (0.0%) | 3 (0.0%) |
| Peptoniphilus asaccharolyticus | 5(0.0%) 5(0.1%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 5(0.0%) 5(0.1%) |
| Peptoniphilus harei | 2(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 2 (0.0%) |
| Peptostreptococcus anaerobius | 2(0.0%) 2(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 2(0.0%) 2(0.0%) |
| Prevotella bivia | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Prevotella buccae | 1(0.0%) 1(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) 1 (0.0%) |
| Prevotella denticola | 3(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 3 (0.0%) |
| Prevotella melaninogenica | 3(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 3 (0.0%) |
| Prevotella nigrescens | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Prevotella species | 1(0.0%) 1(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) 1 (0.0%) |
| Propionibacterium avidum | 2(0.0%) | 0(0.0%) 0(0.0%) | 1 (0.5%) | 3 (0.0%) |
| Propionibacterium granulosum | 1 (0.0%) | 0(0.0%) 0(0.0%) | 0 (0.0%) | 1 (0.0%) |
| Propionibacterium species | 2(0.0%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 2 (0.0%) |
| Proteus mirabilis | 102(0.070) 102(1.6%) | 4(1.6%) | 0(0.0%) 0(0.0%) | 2 (0.070 |
| 1 Toteus mituonis | 102(1.070) | 4(1.070) | 0 (0.070) | (1.6%) |
| Proteus penneri | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Proteus vulgaris group | 1(0.0%) 1(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 0(0.0%) 0(0.0%) | 1(0.0%) 1(0.0%) |
| Proteus | 6 (0.1%) | 1 (0.4%) | 0(0.0%) 0(0.0%) | 7 (0.0%) |
| Providencia rettgeri | 6 (0.1%) | 0 (0.0%) | 0(0.0%) 0(0.0%) | 6 (0.1%) |
| Providencia stuartii | 6 (0.1%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 6 (0.1%) |
| Pseudomonas aeruginosa | 147 (2.3%) | 14(5.7%) | 0(0.0%) 0(0.0%) | 161 |
| i seuvonionius veruginosu | 147(2.570) | 14(0.170) | 0 (0.070) | (2.4%) |
| Pseudomonas aeruginosa, mucoid* | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1% |
| Pseudomonas putida | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Pseudomonas stutzeri | 0 (0.0%) | 2(0.8%) | 0 (0.0%) | 2(0.0%) |
| Psychrobacter species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| Rhodococcus species | 1(0.0%) 1(0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Rhodotorula mucilaginosa | 7 (0.1%) | 0 (0.0%) | 0 (0.0%) | 7 (0.1% |
| Roseomonas mucosa | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Roseomonas species | 1(0.0%) 1(0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0% |
| Rothia aeria | 0 (0.0%) | 0 (0.0%) 0 (0.0%) | 1 (0.5%) | 1(0.0%) |
| Rothia dentocariosa | 2(0.0%) | 0(0.0%) 0(0.0%) | 0 (0.0%) | 2 (0.0%) |
| Rothia mucilaginosa | $\frac{2}{8}(0.1\%)$ | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 8 (0.1% |
| Ruminococcus gnavus | 6 (0.1%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 6 (0.1%) |
| Salmonella enterica | 1 (0.0%) | 0 (0.0%) 0 (0.0%) | 0(0.0%) 0(0.0%) | 1 (0.0%) |
| Salmonella typhi | 9 (0.1%) | 0(0.0%) 0(0.0%) | 0(0.0%) 0(0.0%) | 9 (0.1%) |
| Salmonella | 13 (0.2%) | 0(0.0%) 0(0.0%) | 2(1.0%) | 15 (0.2%) |
| Salmonella serotype Paratyphi A | $\frac{13}{2}(0.2\%)$ | 0 (0.0%) 0 (0.0%) | 2(1.0%) 0(0.0%) | 2 (0.0% |
| Salmonella serotype Typhi | 4 (0.1%) | 0(0.0%) 0(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | $\frac{2}{4}(0.0\%)$ |
| Saimonena serotype Typin Serratia liquefaciens | 4(0.1%) 4(0.1%) | 0(0.0%) 0(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 4(0.1) 4(0.1) |
| | 4(0.1%) 1(0.0%) | 0(0.0%) 0(0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 4 (0.1%) 1 (0.0%) |
| Serratia liquefaciens complex Serratia marcescens | 1(0.0%) 57(0.9%) | 1 (0.0%) | $0 (0.0\%) \\ 0 (0.0\%)$ | 1(0.0%) 58 (0.9%) |
| | . , | · · · · | | |
| Sphingomonas paucimobilis | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Sphingomonas species | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0% |
| Staphylococcus aureus | 105~(1.7%) | 4(1.6%) | 1 (0.5%) | 110 |
| | | | | (1.6%) |

| | Not tested (N=6260) | Negative (N=245) | Positive (N=201) | $\begin{array}{c} \text{Total} \\ (N=6706) \end{array}$ |
|---|------------------------|---------------------|---------------------|---|
| Staphylococcus aureus, Methicillin Resistant | 352~(5.6%) | 9~(3.7%) | 4 (2.0%) | 365 (5.4%) |
| $Staphylococcus \ aureus,$ Methicillin Susceptible | 548 (8.8%) | $21 \ (8.6\%)$ | 6 (3.0%) | (5.4%) 575 (8.6%) |
| Staphylococcus auricularis | 9(0.1%) | 0 (0.0%) | 1 (0.5%) | $10 \ (0.1\%)$ |
| Staphylococcus capitis | 139 (2.2%) | 7(2.9%) | 11 (5.5%) | 157 (2.3%) |
| Staphylococcus caprae | 5(0.1%) | 0 (0.0%) | 0 (0.0%) | 5(0.1%) |
| Staphylococcus cohnii | 6 (0.1%) | 0 (0.0%) | 0 (0.0%) | 6 (0.1%) |
| Staphylococcus epidermidis | 633 (10.1%) | 24 (9.8%) | 39~(19.4%) | 696 (10.4%) |
| Staphylococcus epidermidis, Methicillin Susceptible | 15~(0.2%) | $0 \ (0.0\%)$ | 0 (0.0%) | (10.4%) 15 (0.2%) |
| Staphylococcus epidermidis, Methicillin | 24~(0.4%) | 0 (0.0%) | 0 (0.0%) | 24~(0.4%) |
| Resistant | | | | |
| Staphylococcus equorum | 2(0.0%) | 0 (0.0%) | 0 (0.0%) | 2(0.0%) |
| Staphylococcus haemolyticus | 59(0.9%) | 1(0.4%) | 8(4.0%) | 68(1.0%) |
| Staphylococcus hominis | 209 (3.3%) | 7(2.9%) | 28 (13.9%) | 244 $(3.6%)$ |
| Staphylococcus lugdunensis | 22 (0.4%) | 0 (0.0%) | 1 (0.5%) | 23(0.3%) |
| Staphylococcus pasteuri | 1(0.0%) | 1(0.4%) | 0(0.0%) | 2(0.0%) |
| Staphylococcus pettenkoferi | 33(0.5%) | 1(0.4%) | 0(0.0%) | 34(0.5%) |
| Staphylococcus pseudintermedius | 3(0.0%) | 2(0.8%) | 0(0.0%) | 5(0.1%) |
| Staphylococcus saccharolyticus | 6(0.1%) | 0(0.0%) | 0(0.0%) | 6(0.1%) |
| Staphylococcus saprophyticus | 4(0.1%) | 2(0.8%) | 0(0.0%) | 6(0.1%) |
| Staphylococcus simulans | 6(0.1%) | 0(0.0%) | 0(0.0%) | 6(0.1%) |
| Staphylococcus species | 3(0.0%) | 0(0.0%) | 0(0.0%) | 3(0.0%) |
| Staphylococcus warneri | 12 (0.2%) | 2(0.8%) | 5(2.5%) | 19 (0.3%) |
| Staphylococcus, Coagulase-negative | 171 (2.7%) | 3 (1.2%) | 8 (4.0%) | $182 \\ (2.7\%)$ |
| Staphylococcus, Coagulase-negative, Methicillin | 41~(0.7%) | 4~(1.6%) | 7~(3.5%) | 52~(0.8%) |
| Resistant | | | | |
| <i>Staphylococcus</i> , Coagulase-negative, Methicillin Susceptible | 39~(0.6%) | 4(1.6%) | $10 \ (5.0\%)$ | 53~(0.8%) |
| $Stenotrophomonas\ maltophilia$ | 19~(0.3%) | 0~(0.0%) | 0~(0.0%) | 19 (0.3%) |
| Streptococci | 1 (0.0%) | 0~(0.0%) | 0~(0.0%) | 1~(0.0%) |
| Streptococci, Viridans Group | 34~(0.5%) | 2 (0.8%) | 2(1.0%) | 38~(0.6%) |
| Streptococcus anginosus | 30~(0.5%) | 4(1.6%) | 2(1.0%) | 36~(0.5%) |
| $Streptococcus \ constellatus$ | 18~(0.3%) | 0~(0.0%) | 0~(0.0%) | $18 \ (0.3\%)$ |
| $Streptococcus \ constellatus \ subsp \ constellatus$ | 7~(0.1%) | 0~(0.0%) | 0~(0.0%) | 7~(0.1%) |
| Streptococcus cristatus | 10~(0.2%) | 0~(0.0%) | 0~(0.0%) | $10 \ (0.1\%)$ |
| Streptococcus dysgalactiae (Group C or G) | 9~(0.1%) | 4~(1.6%) | 0~(0.0%) | 13~(0.2%) |
| Streptococcus dysgalactiae subsp equisimilis | 8~(0.1%) | 0~(0.0%) | 0~(0.0%) | 8~(0.1%) |
| $Streptococcus \ gallolyticus$ | 25~(0.4%) | 1 (0.4%) | 0~(0.0%) | 26~(0.4%) |
| $Streptococcus \ gordonii$ | 38~(0.6%) | 0~(0.0%) | 0~(0.0%) | 38~(0.6%) |
| Streptococcus infantarius | 1 (0.0%) | 0 (0.0%) | 0 (0.0%) | 1 (0.0%) |
| $Streptococcus \ intermedius$ | 1 (0.0%) | 0~(0.0%) | 1 (0.5%) | 2 (0.0%) |
| $Streptococcus \ lutetiensis \ (S. bovis)$ | 5~(0.1%) | 0~(0.0%) | 0 (0.0%) | 5(0.1%) |
| Streptococcus mitis group | 15~(0.2%) | 0~(0.0%) | 0~(0.0%) | 15~(0.2%) |
| Streptococcus mitis oralis group | 47 (0.8%) | 0~(0.0%) | 0 (0.0%) | 47 (0.7%) |
| Streptococcus mitis | 25~(0.4%) | 1 (0.4%) | 0~(0.0%) | 26~(0.4%) |

| | Not tested $(N=6260)$ | Negative (N=245) | Positive (N=201) | Total $(N=6706)$ |
|--|-----------------------|---------------------|---------------------|------------------|
| Streptococcus mutans | 7 (0.1%) | 0 (0.0%) | 0 (0.0%) | 7 (0.1%) |
| Streptococcus parasanguinis | 26(0.4%) | 1(0.4%) | 0(0.0%) | 27(0.4%) |
| Streptococcus pasteurianus (S.bovis) | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Streptococcus pneumoniae | 92(1.5%) | 6(2.4%) | 0(0.0%) | 98 (1.5%) |
| Streptococcus pseudoporcinus | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Streptococcus salivarius group | 4 (0.1%) | 0(0.0%) | 0(0.0%) | 4(0.1%) |
| Streptococcus salivarius | 19(0.3%) | 1(0.4%) | 0(0.0%) | 20(0.3%) |
| Streptococcus sanguinis | 27(0.4%) | 0(0.0%) | 1(0.5%) | 28 (0.4%) |
| Streptococcus species | 88 (1.4%) | 5(2.0%) | 2(1.0%) | 95(1.4%) |
| Streptococcus, Beta Hemolytic, Group G | 9(0.1%) | 0(0.0%) | 0(0.0%) | 9(0.1%) |
| Tsukamurella species | 3(0.0%) | 0(0.0%) | 0(0.0%) | 3(0.0%) |
| Veillonella parvula | 3(0.0%) | 0(0.0%) | 0(0.0%) | 3(0.0%) |
| Veillonella species | 1(0.0%) | 2(0.8%) | 0(0.0%) | 3(0.0%) |
| Weissella confusa | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Weissella viridescens | 1(0.0%) | 0(0.0%) | 0(0.0%) | 1(0.0%) |
| Anaerobic Gram Positive Rod | 5(0.1%) | 0(0.0%) | 0(0.0%) | 5(0.1%) |
| Yeast | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |
| Yeast, not Candida albicans | 2(0.0%) | 0(0.0%) | 0(0.0%) | 2(0.0%) |

| | Not tested $(N=1672)$ | Negative $(N=112)$ | Positive $(N=75)$ | Total (N= 1859) | p value |
|----------------|-----------------------|--------------------|-------------------|--------------------|-------------|
| Incubation day | | | | | 0.002^{1} |
| 1 | 993~(59.4%) | 65~(58.0%) | 26 (34.7%) | 1084~(58.3%) | |
| 2 | 487 (29.1%) | 32(28.6%) | 37(49.3%) | 556(29.9%) | |
| 3 | 113(6.8%) | 7(6.2%) | 10 (13.3%) | 130 (7.0%) | |
| 4 | 48(2.9%) | 6(5.4%) | 1(1.3%) | 55(3.0%) | |
| 5 | 31~(1.9%) | 2(1.8%) | 1(1.3%) | 34(1.8%) | |

Supplementary Table 4: Proportion of positive cultures identified by gram stain at days 1 to 5 of culture

1. Pearson's Chi-squared test

| ID | SARS-CoV-2 Status | Time to Gram stain | Organism Name | LikelyContaminant |
|-------|-------------------|--------------------|--|-------------------|
| 08770 | Not tested | 5.0 days | Escherichia coli | No |
| 10753 | Not tested | 4.2 days | Burkholderia multivorans | No |
| 11446 | Not tested | 4.2 days | Candida glabrata | No |
| 12500 | Not tested | 4.3 days | Klebsiella pneumoniae | No |
| 14732 | Not tested | 4.4 days | Escherichia coli | No |
| 14883 | Not tested | 4.7 days | Propionibacterium granulosum | No |
| 15480 | Not tested | 4.7 days | Burkholderia multivorans | No |
| 15750 | Not tested | 4.9 days | Gordonia aichiensis | No |
| 16262 | Not tested | 4.1 days | Klebsiella pneumoniae | No |
| 16516 | Not tested | 4.3 days | Staphylococcus aureus, Methicillin Susceptible | No |
| 18586 | Not tested | 4.4 days | Fusobacterium nucleatum | No |
| 18665 | Not tested | 4.8 days | Proteus mirabilis | No |
| 18993 | Not tested | 4.0 days | Enterococcus faecalis | No |
| 19101 | Not tested | 4.9 days | Proteus mirabilis | No |
| 19834 | Not tested | 4.7 days | Staphylococcus aureus, Methicillin Susceptible | No |
| 20852 | Not tested | 4.2 days | Staphylococcus aureus, Methicillin Susceptible | No |
| 20900 | Not tested | 4.5 days | Candida albicans | No |
| 21446 | Not tested | 4.6 days | Haemophilus parainfluenzae | No |
| 08825 | Not tested | 4.4 days | Anaerobic Gram Positive Rod | Yes |
| 09160 | Not tested | 4.2 days | Staphylococcus epidermidis | Yes |
| 09309 | Not tested | 4.2 days | Staphylococcus hominis | Yes |
| 09536 | Not tested | 5.0 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 11928 | Not tested | 4.4 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 12590 | Not tested | 4.1 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 13366 | Not tested | 4.5 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 14694 | Not tested | 4.6 days | Corynebacterium afermentans | Yes |
| 15565 | Not tested | 4.3 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 16048 | Not tested | 4.8 days | Staphylococcus pettenkoferi | Yes |
| 17171 | Not tested | 4.4 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 18105 | Not tested | 4.5 days | Cutibacterium (Propionibacterium) acnes | Yes |
| 18638 | Not tested | 4.3 days | Staphylococcus lugdunensis | Yes |
| 19887 | Not tested | 4.8 days | Staphylococcus pettenkoferi | Yes |
| 19887 | Not tested | 4.8 days | Corynebacterium | Yes |
| 22589 | Negative | 4.1 days | Proteus mirabilis | No |
| 23268 | Negative | 4.7 days | Candida glabrata | No |
| 23021 | Positive | 4.0 days | Candida albicans | No |

Supplementary Table 5: Organisms identified after 4 days of incubation