

Fusing a Bayesian case velocity model with random forest for predicting COVID-19 in the U.S.: Supplemental material

Gregory L. Watson¹, Di Xiong¹, Lu Zhang¹, Joseph A. Zoller¹, John Shamshoian¹, Phillip Sundin¹, Teresa Bufford¹, Anne W. Rimoin², Marc A. Suchard^{1,3}, Christina M. Ramirez¹

1 Department of Biostatistics, Fielding School of Public Health, University of California, Los Angeles, California, United States of America

2 Department of Epidemiology, Fielding School of Public Health, University of California, Los Angeles, California, United States of America

3 Departments of Biomathematics and Human Genetics, David Geffen School of Medicine, University of California, Los Angeles, California, United States of America

Parameter Values and Prior Distributions

Table 1 lists parameters for the various components of the model and gives their values or distributions. The ρ parameter governs flow out of the I compartment of the SIRD model. Its inverse, ρ^{-1} , is interpreted as the expected amount of time until an infected individual recovers or dies.

Table 1. Parameter Values & Distributions

Model	Parameter	Value/Distribution
SIRD	ρ^{-1}	$N(14, 1)$
Mixed Model	μ_α	$N(-1, 1)$
Mixed Model	μ_β	$N(0, 0.1)$
Mixed Model	μ_γ	$N(0, 1)$
Mixed Model	μ_δ	$N(-0.05, 0.1)$
Mixed Model	μ_η	$N(0, 1)$
Mixed Model	μ_ζ	$N(0, 0.1)$
Mixed Model	σ_α^2	1
Mixed Model	σ_β^2	0.1
Mixed Model	σ_γ^2	1
Mixed Model	σ_δ^2	0.1
Mixed Model	σ_η^2	1
Mixed Model	σ_ζ^2	0.1
Mixed Model	n.chains	3
Mixed Model	n.iter	500,000
Mixed Model	n.burnin	10,000
Mixed Model	n.thin	500
Random Forest	ntree	500
Random Forest	mtry	9
Random Forest	nodesize	5

Values or prior distributions for the parameters of each component of the model.