

## Delirium Incidence, Duration and Severity in Critically Ill Patients with COVID-19

**Supplementary Table 1.** Laboratory Values and Hemodynamic Characteristics of Patients Admitted with Novel Coronavirus to the Intensive Care Unit<sup>a</sup>

Variable	Total (n=144)	Delirium Positive (n=106)	Delirium Negative (n=38)	P value
<b>All ICU Patients</b>				
Sodium mmol/L, median (IQR)	137.0 (135.0-139.5)	137.0 (135.0-139.0)	137.0 (135.0-140.0)	0.888
Hematocrit %, median (IQR)	36.7 (32.0-40.9)	36.0 (30.0-40.0)	38.1 (34.4-42.0)	0.088
Platelet count x10 <sup>3</sup> , median (IQR)	216.0 (152.0-256.0)	213.0 (145.0-252.0)	227.0 (185.0-278.0)	0.208
Creatinine mg/dl, median (IQR)	1.1 (0.8-1.8)	1.1 (0.8-1.9)	1.1 (0.8-1.7)	0.585
Mean Arterial Pressure mmHg, median (IQR)	100.0 (88.0-112.0)	100.0 (87.0-112.0)	100.5 (92.0-115.0)	0.691


<sup>a</sup>Data presented is from laboratory values, arterial blood gas results, vitals, and other components of the electronic medical record (first 24 hours of intensive care unit admission).

**Supplementary Table 2.** Delirium rates and clinical outcomes for patients admitted to the intensive care unit during the 2009-1010 Influenza Pandemic.

Outcomes	2010 (n=383)
<b>Delirium Incidence and Subtypes</b>	
Delirium incidence* (%)	22.7
Hypoactive delirium** (%)	19.3
Hyperactive delirium** (%)	3.4
<b>Coma and Level of Consciousness</b>	
Coma*** (%)	24.0
Level of consciousness at ICU admission by average RASS score – median IQR	-0.3 (-2, 0)
<b>Diagnoses****</b>	
Acute Respiratory failure (%)	26.6
Acute renal failure (%)	29.5
Trauma (%)	15.9
Hypoalbuminemia	26.6
<b>Clinical Outcomes</b>	
Underwent invasive mechanical ventilation (%)	36.0
Hospital mortality (%)	3.4
<b>Medication Orders*****</b>	
Benzodiazepines (%)	20.4
Opioids (%)	66.3

n: Unique patient admissions. \*Patients with delirium by positive CAM-ICU at any time during the follow up period (at first assessment or during subsequent screenings in the intensive care unit); \*\* Delirium subtypes determined based on first positive delirium screening. \*\*\*Coma defined by RASS score -4 or -5. \*\*\*\*Diagnoses extracted from the medical record. \*\*\*\*\*Medication orders indicate the percentage of patients with active orders. CAM-ICU: Confusion Assessment Method for the Intensive Care Unit. RASS: Richmond Agitation-Sedation Scale.

**Supplementary Figure 1.** Flow Diagram of Patients with COVID-19 Screened For Inclusion in the Cohort



**Supplementary Figure 2.** Daily Rates of Patient's Discharge Status As Assessed Up to First 14 Days of Intensive Care Unit Stay (n=144)

