

An MRI-based, data-driven model of cortical laminar connectivity

Ittai Shamir ¹ and Yaniv Assaf ^{1,2}

¹ Department of Neurobiology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel

² Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

* Correspondence:

Ittai Shamir

ittaisha@mail.tau.ac.il

ORCID: 0000-0003-4028-5154

Electronic Supplementary Material

1. Summary of reviewed articles

Following is a summary table of all 51 articles included in our systematic review (table 1):

	First Author/s	Year	Journal	Human/ Animal	Method	Model Y/N	Region/s
1	Hubel & Wiesel	1962	Journal of Physiology	Cat	Recordings	N	Visual cortex
2	Herkenham	1980	Science	Rat	Microphotographs (also mention histology, electrophysiology, tracing)	N	primary somatosensory
3	Douglas et al.	1989	Neural Computation	Cat (generalized)	Intracellular recordings, histological staining (horseradish)	Y	whole-cortex (canonical), based on visual cortex
4	Felleman & Van Essen	1991	Cerebral Cortex	Primate (macaque)	Histology	Y	multiple cortical regions (occipital, temporal, parietal)
5	Douglas & Martin	1991	Journal of Physiology	Cat	Recording of electrical stimulation	Y	Visual cortex
6	Mumford	1991	Biological Cybernetics	Cat	Electrical recording	N	V1 and primary motor
7	Mountcastle	1997	Brain	Primate (monkey, macaque) mainly, also cat	Electrical recording	Y	general
8	Nelson	2002	Neuron	Cat, rodent (mouse), primate, human	Recording, photomicrograph, histology, photostimulation	N	canonical circuit, visual cortex
9	Buxhoeveden & Casanova	2002	Brain	Cat, rodent (mouse, rat, rabbit), primate (rhesus monkey), human	Histology (Nissl, Golgi staining), metabolic labelling, electrical recording	Y	intrinsic pathways
10	DeFelipe et al.	2002	Journal of Neurocytology	Multiple	Multiple	N	multiple
11	Raizada & Grossberg	2003	Cerebral Cortex	Macaque mainly, also cat	Multiple methods (review of functional recorded connection into a single model- LAMINART +simulation)	Y	Visual cortex
12	Binzegger et al.	2004	The Journal of Neuroscience	Cat	In-vivo staining, in vivo recordings	Y	Visual Cortex

Electronic Supplementary Material

13	Douglas & Martin	2004	Annual Review of Neuroscience	Human, cat, rat, monkey (multiple-review)	Multiple (review)- staining (Golgi), electrophysiology, degeneration, retrograde traces (horseradish, peroxidase, fast blue), ESEM	Y	Visual Cortex, generalized to entire cortex
14	Shipp	2005	Philosophical transactions of the royal society	Animal, some human (multiple-review)	Multiple- imaging, histology, recording. Etc.	Y	General (agranular, granular and dysgranular regions connectivity)
15	Hirsch & Martinez	2006	Current Opinion in Neurobiology	Cat, also monkey	Recordings of stimuli	Y	Visual cortex
16	Douglas & Martin	2007	Neuron	Cat, primate	Staining, electrophysiology, etc.	Y	Canonical circuit, visual Cortex
17	Lubke & Feldmeyer	2007	Brain Structure and Function	Rodent mainly, also human and primate	Electrical recording, infrared microscopy, optical probing, photostimulation, ESEM	Y	Simplified model
18	Helmstaedter et al.	2007	Brain Research Reviews	Rat	Electrical recordings, optical reconstruction	Y	Somatosensory, general
19	Thomson & Lamy	2007	Frontiers in Neuroscience	Cat, mice ('green mice')	Cellular recordings (intra/whole-cell), anatomy, imaging, (histology), stereological, analysiss, immunocytochemical, cellular markers, genomics, immunofluorescence	Y	General
20	Shipp	2007	Current Biology	Primate	Electric recording, etc.	Y	Nonprimary visual area
21	Helmstaedter et al.	2008	The Journal of Neuroscience	Rat	Electrical recording, histology	Y	Somatosensory
22	Weiler et al.	2008	Nature Neuroscience	Mouse	Electrical recording	Y	Sensory cortex
23	Binzegger et al.	2009	Neural Networks	Cat	Horseradish labelling	Y	Visual cortex
24	Da Costa & Martin	2010	Frontiers in Neuroanatomy	Multiple (cat mainly, also primates and rodents)	Multiple	Y	Intra-regional
25	Thomson	2010	Frontiers in Neuroanatomy	Mice, rat, primate, cat, human	Electrophysiological, dye (histology), retrograde labelling	Y	Primary sensory
26	Meyer et al.	2010	Cerebral Cortex	Rat	Staining (immunofluorescence), single-cell recording	Y	Somatosensory
27	Meyer et al.	2010	Cerebral Cortex	Rat	Staining (immunofluorescence), single-cell recording	N	Somatosensory
28	Fishell & Rudy	2011	Annual Review of Neuroscience	Human, rodent, cat, primate (multiple)	Multiple methods (historical review focused on inhibition)	N	General
29	Shepherd	2011	Frontiers in Neuroanatomy	Turtle, human, also cat, primate, opossum (multiple-review)	Electrical recording, histology, genetics, etc. (multiple- review)	Y	General
30	Feldmeyer	2012	Frontiers in Neuroanatomy	Rodents (rat and mouse)	Electrophysiological, dye (histology), anterograde and retrograde tracing, cage glutamate release, light activation, etc.	Y	somatosensory (barrel) cortex

Electronic Supplementary Material

31	Du et al.	2012	PLoS Computational Biology	Multiple- review as the basis for the model	Multiple- review +simulation	Y	Visual cortex, generalized
32	Bastos et al.	2012	Neuron	Mammal, rodents (mice,rats)	Multiple: fMRI, recordings, optogenetics, photostimulation, computational model	Y	Visual cortex
33	Kennedy & Dehay	2012	Progress in Brain Research	Mainly human, also primate, carnivores, rodents, insectivores, reptiles, amphibians	Multiple	Y	General
34	Estrada-Sanchez & Rebec	2013	Frontiers in Neural Circuits	Multiple	Multiple	N	Visual cortex
35	Harris & Mrsic-Flogel	2013	Nature	Rodents	Optical methods, electrical methods, optogenetics	Y	Sensory cortex
36	Beul & Hilgetag	2014	Frontiers in Neuroanatomy	Mainly rodent (also discuss human, cat)	Multiple, including histology (anatomical and electrophysiological approaches)	Y	Cortico-cortical
37	Kiernan & Rajakumar	2014	Lippincott Williams & Wilkins	Human	Histology (Nissl, Golgi), ESEM, immunohistochemical	Y	Intracortical
38	Markov et al.	2014	The Journal of Comparative Neurology, Research in Systems Neuroscience	Macaque	Retrograde tracing + model	Y	Visual cortex
39	Dhruv	2015	Nature Neuroscience	Mouse, human	Recording, optogenetics	Y	Canonical
40	Lodato & Arlotta	2015	Annual Review of Developmental Biology	Mouse, rat	Histology (Golgi staining, etc.), electrical recording, tracing, optogenetics	Y	Barrel cortex
41	Bosman & Aboitiz	2015	Frontiers in Neuroscience	Animals- mammals (cats, rodents, primates) and sauropsids (birds and reptiles) (review of many studies)	Multiple methods- histology, tracing, electrical recording (review of many studies)	Y	Canonical circuit, Visual Cortex
42	Jiang et al.	2015	Science	Mice	Tissue slicing, whole-cell recording, morphological reconstruction	Y	General / canonical
43	Markram et al.	2015	Cell	Rat	Digital reconstruction from experimental data recordings, followed by simulations	Y	Somatosensory
44	Ramaswamy et al.	2015	Frontiers in Neural Circuits	Rat	Digital reconstruction from experimental data recordings, followed by simulations (data includes many methods, database of papers)	Y	Somatosensory
45	Wang & Kennedy	2016	, Current Opinion in Neurobiology	Monkey (macaque), rodent (mouse), (generalized model for mammals)	Multiple: recordings,	N	General
46	Guy & Staiger	2017	Frontiers in Neuroanatomy	Mouse	Optogenetics	N	Somatosensory
47	Mercer & Thomson	2017	Frontiers in Neuroanatomy	Primate, rat, human, cat (review)	Multiple (review)	N	General

48	Hawkins & Ahmad	2017	Frontiers in Neural Circuits	Mouse	Multiple (electrical recording, etc.)- focus on simulation	Y	General
49	Rockland	2017	NeuroImage	Multiple (review)- mainly primates (NHP), also rodents, cats	Multiple methods (review)- anterograde/retrograde tracing, MRI	Y	General
50	Roelfsema & Holtmaat	2018	Nature Reviews Neuroscience	Primate, Rodent (human)	Stimulation, simulation (deep learning)	Y	Intracortical (local)
51	Larkum et al.	2018	Frontiers in Neuroanatomy	Human, rodents	fMRI (high resolution)	Y	General

Table 1 A summary of all 51 records included in the systematic review

2. Summary of region-specific reported connections

Following is a summary of reported region-specific cortical laminar connections, organized by origin and termination in the three-layer grouping of infragranular, granular and supragranular layers:

2.1 Visual cortex

For a summary of reported cortical laminar connections in the visual cortex see figure 1 (below).




Fig. 1 Summary of all reported cortical laminar connections in the ‘visual cortex’:

- a- In connectivity matrix format, where: [row, column]=[connection origin, connection termination] and connections are averaged per origin (per row)
- b- Presented as distributions of connection terminations that originate in each of the following: subcortex (left), infragranular (mid left), granular (mid right), supragranular (right)

2.2 Sensory cortex

For a summary of reported cortical laminar connections in the sensory cortex see figure 2 (below).




Fig. 2 Summary of all reported cortical laminar connections in the ‘sensory cortex’:

- a- In connectivity matrix format, where: [row, column]=[connection origin, connection termination] and connections are averaged per origin (per row)

- b- Presented as distributions of connection terminations that originate in each of the following: subcortex (left), infragranular (mid left), granular (mid right), supragranular (right)

2.3 General

For a summary of reported cortical laminar connections in the ‘general’ cortical regions see figure 3 (below).




Fig. 3 Summary of all reported cortical laminar connections in ‘general’ regions:

- a- In connectivity matrix format, where: [row, column]=[connection origin, connection termination] and connections are averaged per origin (per row)
- b- Presented as distributions of connection terminations that originate in each of the following: subcortex (left), infragranular (mid left), granular (mid right), supragranular (right)

2.4 Somatosensory cortex

For a summary of reported cortical laminar connections in the somatosensory cortex see figure 4 (below).




Fig. 4 Summary of all reported cortical laminar connections in ‘somatosensory cortex’:

- a- In connectivity matrix format, where: [row, column]=[connection origin, connection termination] and connections are averaged per origin (per row)
- b- Presented as distributions of connection terminations that originate in each of the following: subcortex (left), infragranular (mid left), granular (mid right), supragranular (right)

2.5 Other

For a summary of reported cortical laminar connections in ‘other’ cortical regions see figure 5 (below).




Fig. 5 Summary of all reported cortical laminar connections in ‘other’ regions:

- a- In connectivity matrix format, where: [row, column]=[connection origin, connection termination] and connections are averaged per origin (per row)
- b- Presented as distributions of connection terminations that originate in each of the following: subcortex (left), infragranular (mid left), granular (mid right), supragranular (right)

*Too little region-specific reported data regarding local cortical laminar connectivity in motor and auditory cortices.

3. Explanation of origin of rules for horizontal laminar connections

Horizontally oriented laminar connections are extrinsic nonlocal connections between laminar components of different cortical regions. Our model's rules for horizontal connections were extracted from a general schematic model of directed connections (see figure 6).




Fig. 6 Schematic model for horizontal laminar connections, based on the granularity of the connecting regions. The model is adapted from a similar model, which has been widely reported and studied since the 1980s (Beul and Hilgetatg 2015, adapted from von Economo 2009 and from Barbas and Rempel-Clover 1997; a similar model also appears in Shipp 2005, adapted from Barbas and Rempel-Clover 1997 and Barabs 1986)

An initial set of rules of directed laminar horizontal connectivity was extracted from the abovementioned schematic model, based on the granularity indices and laminar compositions of the regions of origin and termination (see table 2). The rules of directed connectivity were then expanded from multiple laminar regions of origin/termination to single laminar groups, using the corresponding laminar composition (see table 3, rules: C.1.1, C.1.2, C.4.1, C.4.2, D.3.1, D.3.2, D.4.1, D.4.2).

After the rules of directed connectivity were expanded, they were organized into undirected categories according to the connecting regions (see table 4, organized by color). Subsequently, the probabilities within each category were summed, resulting in the final undirected model of horizontally oriented connections (see table 5).

a	Number:	Cortex granularity:
	1	Agranular
	2	Slightly granular
	3	Increasing granule cell presence
	4	Increasing granule cell presence
	5	Increasing granule cell presence
6	Granular	

b	Termination:	1	2	3	4	5	6
	Origin:						
	1	E, local	C			A	
	2	D	E, local	C			
	3	D		E, local	C		
	4	D			E, local	C	
5	D				E, local	C	
6	B	D			E, local		

c	Rule:	Subcategory:	Origin:	Termination:	Origin		Termination		Strength probability:
					Layer composition:	Prob.:	Layer composition:	Prob.:	
A	A	1	6	P(IG,low)	1	P(SG,high)	1	$\frac{P(IG,low)}{P(IG,low) + P(SG,high)}$	
B	B	6	1	P(SG,high)	1	P(IG,low)	1	$\frac{P(SG,high)}{P(IG,low) + P(SG,high)}$	
	C.1			P(SG,low)	(1/3)	P(IG,high)+P(G,high)	(1/3)	$\left(\frac{1}{3}\right)^2 \cdot P_{low2high}$	
	C.2			P(SG,low)	(1/3)	P(SG,high)	(2/3)	$\left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right) \cdot P_{low2high}$	
	C.3			P(IG,low)	(2/3)	P(SG,high)	(2/3)	$\left(\frac{2}{3}\right)^2 \cdot P_{low2high}$	
C	C.4	X	>X	P(IG,low)	(2/3)	P(IG,high)+P(G,high)	(1/3)	$\left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right) \cdot P_{low2high}$	
	D.1			P(SG,high)	(2/3)	P(IG,low)	(2/3)	$\left(\frac{2}{3}\right)^2 \cdot P_{high2low}$	
	D.2			P(SG,high)	(2/3)	P(SG,low)	(1/3)	$\left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right) \cdot P_{high2low}$	
	D.3			P(IG,high)+P(G,high)	(1/3)	P(SG,low)	(1/3)	$\left(\frac{1}{3}\right)^2 \cdot P_{high2low}$	
D	D.4	>X	X	P(IG,high)+P(G,high)	(1/3)	P(IG,low)	(2/3)	$\left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right) \cdot P_{high2low}$	
	E.1			P(SG,1)	1	P(SG,2)	1	$\frac{P(SG,1)}{P(SG,1) + P(SG,2)}$	
	E.2			P(SG,2)	1	P(SG,1)	1	$\frac{P(SG,2)}{P(SG,1) + P(SG,2)}$	
E	E.2	X	X	P(SG,2)	1	P(SG,1)	1	$\frac{P(SG,2)}{P(SG,1) + P(SG,2)}$	

Table 2 Initial set of rules of directed horizontal connectivity, expanded from figure 6:

- a- Cortical granularity indices, ranging from agranular (1) to granular (6);
 - b- Rules of directed laminar connectivity (A-E), applied according to granularity of connecting regions;
 - c- Full description of all rules of directed connectivity, estimated according to their laminar content.
- Final laminar connection strengths are calculated by multiplying tractography connection strengths by laminar strength probabilities (last column);

Where: P(L,R)- composition of laminar group L (SG- supragranular, G- granular , IG- infragranular) in region R (high/low represents relative granularity index, or 1/2 when they are equal)

Rule:	Subcategory:	Origin:	Termination:	Origin		Termination		Strength probability:			
				Layer composition:	Prob.:	Layer composition:	Prob.:				
A	A	1	6	P(IG,low)	1	P(SG,high)	1	$\frac{P(IG,low)}{P(IG,low) + P(SG,high)}$			
B	B	6	1	P(SG,high)	1	P(IG,low)	1	$\frac{P(SG,high)}{P(IG,low) + P(SG,high)}$			
C	C.1	X	>X	P(SG,low)	(1/3)	P(IG,high)+P(G,high)	(1/3)	$S1 = (\frac{1}{3})^2 \cdot P_{low2high}$			
	C.1.1					P(IG,high)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S1$			
	C.1.2					P(G,high)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S1$			
	C.2			P(SG,low)	(1/3)	P(SG,high)	(2/3)	$(\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{low2high}$			
	C.3			P(IG,low)	(2/3)	P(SG,high)	(2/3)	$(\frac{2}{3})^2 \cdot P_{low2high}$			
	C.4			P(IG,low)	(2/3)	P(IG,high)+P(G,high)	(1/3)	$S2 = (\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{low2high}$			
	C.4.1					P(IG,high)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S2$			
	C.4.2					P(G,high)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S2$			
	D			D.1	X	X	P(SG,high)	(2/3)	P(IG,low)	(2/3)	$(\frac{2}{3})^2 \cdot P_{high2low}$
	D.2			P(SG,high)			(2/3)	P(SG,low)	(1/3)	$(\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{high2low}$	
D.3	P(IG,high)+P(G,high)	(1/3)	P(SG,low)	(1/3)			$S3 = (\frac{1}{3})^2 \cdot P_{high2low}$				
D.3.1			P(IG,high)				$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S3$				
D.3.2			P(G,high)				$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S3$				
D.4	P(IG,high)+P(G,high)	(1/3)	P(IG,low)	(2/3)			$S4 = (\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{high2low}$				
D.4.1			P(IG,high)				$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S4$				
D.4.2			P(G,high)				$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S4$				
E	E.1	X	X	P(SG,1)			1	P(SG,2)	1	$\frac{P(SG,1)}{P(SG,1) + P(SG,2)}$	
E.2							P(SG,2)	1	P(SG,1)	1	$\frac{P(SG,2)}{P(SG,1) + P(SG,2)}$

Table 3 Expanded set of rules of directed horizontal connectivity, expanded to single lamina groups from table 2c: Full description of all rules of directed connectivity, estimated according to their lamina content. Final lamina connection strengths are calculated by multiplying tractography connection strengths by lamina strength probabilities (last column);

Where: P(L,R)- composition of lamina group L (SG- supragranular, G- granular, IG- infragranular) in region R (high/low represents relative granularity index, or 1/2 when they are equal);

Additional values:

$$P_{low} = \frac{2}{3} \cdot P(IG,low) + \frac{1}{3} \cdot P(SG,low) ; P_{high} = \frac{2}{3} \cdot P(SG,high) + \frac{1}{3} \cdot (P(IG,high) + P(G,high)) ;$$

$$P_{low2high} = \frac{P_{low}}{P_{low} + P_{high}} ; P_{high2low} = \frac{P_{high}}{P_{low} + P_{high}}$$

$$S1 = (\frac{1}{3})^2 \cdot P_{low2high} ; S2 = (\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{low2high} ; S3 = (\frac{1}{3})^2 \cdot P_{high2low} ; S4 = (\frac{1}{3}) \cdot (\frac{2}{3}) \cdot P_{high2low}$$

Rule:	Subcategory:	Connecting regions:		Region 1		Region 2		Strength probability:
				Layer composition:	Prob.:	Layer composition:	Prob.:	
A	A	1	6	P(IG,low)	1	P(SG,high)	1	1
B	B.1	X	>X	P(SG,low)		P(IG,high)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S1$
				P(IG,high)		P(SG,low)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S3$
	B.2			P(SG,low)		P(G,high)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S1$
				P(G,high)		P(SG,low)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S3$
	B.3			P(SG,low)	(1/3)	P(SG,high)	(2/3)	$\frac{1}{3} \cdot \frac{2}{3} \cdot P_{low2high}$
				P(SG,high)	(2/3)	P(SG,low)	(1/3)	$\frac{1}{3} \cdot \frac{2}{3} \cdot P_{high2low}$
	B.4			P(IG,low)	(2/3)	P(SG,high)	(2/3)	$\frac{2}{3}^2 \cdot P_{low2high}$
				P(SG,high)	(2/3)	P(IG,low)	(2/3)	$\frac{2}{3}^2 \cdot P_{high2low}$
	B.5			P(IG,low)		P(IG,high)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S2$
				P(IG,high)		P(IG,low)		$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot S4$
	B.6			P(IG,low)		P(G,high)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S2$
				P(G,high)		P(IG,low)		$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot S4$
B	C	X	>X	P(G,high)		P(IG,low)		
C	C	X	X	P(SG,1)	1	P(SG,2)	1	1

Table 4 Categorized rules of undirected horizontal connectivity (derived from table 3):

Full description of all rules of undirected connectivity, estimated according to their laminar content.

Final laminar connection strengths are calculated by multiplying tractography connection strengths by laminar strength probabilities (last column);

Where: P(L,R)- composition of laminar group L (SG- supragranular, G- granular , IG- infragranular) in region R (high/low represents relative granularity index, or 1/2 when they are equal);

Additional values as described for table 3

a	Number:	Cortex granularity:
	1	Agranular
	2	Slightly granular
	3	Increasing granule cell presence
	4	Increasing granule cell presence
	5	Increasing granule cell presence
6	Granular	

b	Termination:	1	2	3	4	5	6
	Origin:						
	1	C, local	B				A
	2	B	C, local	B			
	3	B		C, local	B		
	4	B			C, local	B	
5	B				C, local	B	
6	A	B				C, local	

c	Rule:	Subcategory:	Connecting regions:		Region 1	Region 2	Strength probability:
					Layer composition:	Layer composition:	
A	A	1	6	P(IG,low)	P(SG,high)	1	
B	B.1	X	>X	P(IG,low)	P(IG,high)	$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot \left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right)$	
	B.2			P(IG,low)	P(G,high)	$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot \left(\frac{1}{3}\right) \cdot \left(\frac{2}{3}\right)$	
	B.3			P(IG,low)	P(SG,high)	$\left(\frac{2}{3}\right)^2$	
	B.4			P(SG,low)	P(IG,high)	$\frac{P(IG,high)}{P(IG,high)+P(G,high)} \cdot \left(\frac{1}{3}\right)^2$	
	B.5			P(SG,low)	P(G,high)	$\frac{P(G,high)}{P(IG,high)+P(G,high)} \cdot \left(\frac{1}{3}\right)^2$	
	B.6			X	>X	P(SG,low)	P(SG,high)
C	C	X	X	P(XG,1)	P(XG,2)	1	

Table 5 Final undirected model of horizontally oriented connections (derived from table 4):

- a- Cortical granularity indices, ranging from agranular (1) to granular (6);
 - b- Rules of directed laminar connectivity (A-E), applied according to granularity of connecting regions;
 - c- Full description of all rules of undirected connectivity, estimated according to their laminar content.
- Final laminar connection strengths are calculated by multiplying tractography connection strengths by laminar strength probabilities (last column);

Where: P(L,R)- composition of laminar group L (SG- supragranular, G- granular , IG- infragranular) in region R (high/low represents relative granularity index, or 1/2 when they are equal)