

A modest increase in ¹¹C-PK11195-PET TSPO binding in depression is not associated with serum C-reactive protein or body mass index

Julia J. Schubert et al.

Supplemental material




Figure S1. A. Group differences in mean ¹¹C-PK11195 binding potential measurements in anterior cingulate cortex (ACC), prefrontal cortex (PFC), and insular cortex (INS) regions between healthy controls (HC), treatment resistant depression, and untreated depression. **B.** Group differences in mean ¹¹C-PK11195 binding potential measurements in ACC, PFC, and INS regions between healthy controls, depressed cases without suicidal thoughts, and depressed cases with suicidal thoughts. Error bars represent standard error.

Table S1. One-tailed t-statistic results comparing regional TSPO binding potentials (BP_{ND}) in anterior cingulate cortex (ACC), prefrontal cortex (PFC), and insular cortex (INS) brain regions between groupings of depressed cases (DC) (all DC, high C-reactive protein (CRP), low CRP) and healthy controls (HC).

Region		t-statistic	P-Value	Mean±SD	
				DC	HC
ACC	All DC	t(74)=1.996	0.025	1.171±0.036	
	High CRP	t(43)=1.236	0.112	1.168±0.035	1.152±0.046
	Low CRP	t(54)=1.958	0.028	1.174±0.037	
PFC	All DC	t(74)=1.114	0.134	1.204±0.035	
	High CRP	t(43)=0.435	0.333	1.199±0.030	1.194±0.040
	Low CRP	t(54)=1.284	0.102	1.207±0.038	
INS	All DC	t(74)=1.484	0.071	1.200±0.044	
	High CRP	t(43)=1.396	0.085	1.202±0.045	1.184±0.042
	Low CRP	t(54)=1.215	0.115	1.198±0.043	

Table S2. Analysis of variance (ANOVA) results comparing regional TSPO binding potentials (BP_{ND}) in anterior cingulate cortex (ACC), prefrontal cortex (PFC), and insular cortex (INS) brain regions between groupings of depressed cases (DC) (all DC, high C-reactive protein (CRP), low CRP) and healthy controls (HC).

Region		F-statistic	P-Value
Controlled for age			
All regions	All DC	F(1,73)=3.421	0.068
	High CRP	F(1,42)=1.413	0.241
	Low CRP	F(1,53)=3.024	0.088
ACC	All DC	F(1,73)=4.880	0.030
	High CRP	F(1,42)=1.559	0.219
	Low CRP	F(1,53)=4.892	0.031
PFC	All DC	F(1,73)=1.156	0.286
	High CRP	F(1,42)=0.191	0.665
	Low CRP	F(1,53)=1.364	0.248
INS	All DC	F(1,73)=2.567	0.113
	High CRP	F(1,42)=1.929	0.172
	Low CRP	F(1,53)=1.769	0.189
Controlled for age and sex			
All regions	All DC	F(1,72)=5.590	0.021
	High CRP	F(1,41)=3.752	0.060
	Low CRP	F(1,52)=4.102	0.048
ACC	All DC	F(1,72)=6.599	0.012
	High CRP	F(1,41)=3.376	0.073
	Low CRP	F(1,52)=5.720	0.020
PFC	All DC	F(1,72)=2.589	0.112
	High CRP	F(1,41)=0.999	0.323
	Low CRP	F(1,52)=2.261	0.139
INS	All DC	F(1,72)=3.973	0.050
	High CRP	F(1,41)=4.551	0.039
	Low CRP	F(1,52)=2.365	0.130
Controlled for age, sex, and BMI			
All regions	All DC	F(1,71)=6.971	0.010
	High CRP	F(1,40)=7.224	0.010
	Low CRP	F(1,51)=5.708	0.021
ACC	All DC	F(1,71)=9.067	0.004
	High CRP	F(1,40)=5.741	0.021
	Low CRP	F(1,51)=8.815	0.005
PFC	All DC	F(1,71)=3.858	0.053
	High CRP	F(1,40)=3.707	0.061
	Low CRP	F(1,51)=3.460	0.069
INS	All DC	F(1,71)=3.848	0.054
	High CRP	F(1,40)=6.678	0.014
	Low CRP	F(1,51)=2.628	0.111

Table S3. Correlation between regional TSPO binding potentials and experimental variables in anterior cingulate cortex (ACC), prefrontal cortex (PFC), and insular cortex (INS) regions in depressed cases (DC) and healthy controls (HC)

Experimental Variable		Region, r Value (P Value)*		
Variable only		ACC	PFC	INS
CRP	All DC	-0.047(0.745)	-0.073(0.610)	0.053(0.710)
	High CRP	-0.127(0.593)	-0.102(0.670)	-0.050(0.833)
	Low CRP	0.034(0.856)	0.064(0.734)	0.031(0.867)
HDRS	All DC	-0.009(0.951)	-0.035(0.812)	0.080(0.582)
	High CRP	0.486(0.035)*	0.223(0.359)	0.501(0.029)*
	Low CRP	-0.247(0.180)	-0.088(0.638)	-0.186(0.317)
Controlled for age				
CRP	All DC	-0.171(0.241)	-0.184(0.207)	-0.035(0.812)
	High CRP	-0.276(0.252)	-0.173(0.478)	-0.218(0.371)
	Low CRP	-0.024(0.901)	0.098(0.605)	0.044(0.816)
HDRS	All DC	-0.052(0.723)	-0.047(0.746)	0.041(0.781)
	High CRP	0.294(0.236)	0.112(0.658)	0.434(0.072)
	Low CRP	-0.233(0.216)	-0.110(0.564)	-0.219(0.245)
Controlled for age and sex				
CRP	All DC	-0.142(0.334)	-0.127(0.391)	-0.002(0.988)
	High CRP	-0.193(0.458)	-0.194(0.456)	-0.124(0.634)
	Low CRP	0.022(0.911)	0.098(0.611)	0.043(0.827)
HDRS	All DC	-0.077(0.605)	-0.103(0.487)	0.018(0.901)
	High CRP	0.279(0.278)	0.086(0.743)	0.427(0.087)
	Low CRP	-0.258(0.177)	-0.195(0.310)	-0.247(0.197)
Controlled for age, sex, and BMI				
CRP	All DC	-0.155(0.298)	-0.125(0.403)	-0.042(0.778)
	High CRP	-0.213(0.428)	-0.193(0.475)	-0.124(0.646)
	Low CRP	0.064(0.747)	0.072(0.714)	0.012(0.950)
HDRS	All DC	-0.077(0.607)	-0.104(0.486)	0.022(0.884)
	High CRP	0.354(0.178)	0.073(0.788)	0.432(0.095)
	Low CRP	-0.269(0.167)	-0.190(0.333)	-0.242(0.215)

Table S4. Correlation between regional TSPO binding potential (ACC: anterior cingulate cortex; PFC: prefrontal cortex; INS: insular cortex) and C-reactive protein (CRP), Hamilton depression rating scale (HDRS), and body mass index (BMI) in depressed cases.

	PFC	INS	CRP	HDRS	BMI
ACC	r=0.483 P<0.001	r=0.757 P<0.001	r=-0.047 P=0.745	r=-0.009 P=0.951	r=-0.19 P=0.895
	PFC	r=0.456 P=0.001	r=-0.073 P=0.610	r=-0.035 P=0.812	r=0.112 P=0.434
		INS	r=0.053 P=0.710	r=0.080 P=0.582	r=0.044 P=0.760
			CRP	r=0.302 P=0.033	r=0.462 P=0.001
				HDRS	r=0.012 P=0.935

Table S5. Correlation between regional TSPO binding potential (ACC: anterior cingulate cortex; PFC: prefrontal cortex; INS: insular cortex) and C-reactive protein (CRP), Hamilton depression rating scale (HDRS), and body mass index (BMI) in healthy controls.

	PFC	INS	CRP	HDRS	BMI
ACC	r=0.755 P<0.001	r=0.796 P<0.001	r=-0.162 P=0.438	r=0.204 P=0.327	r=-0.045 P=0.831
	PFC	r=0.548 P=0.005	r=0.271 P=0.190	r=0.001 P=0.995	r=0.003 P=0.990
		INS	r=0.243 P=0.241	r=0.401 P=0.047	r=0.059 P=0.780
			CRP	r=0.322 P=0.117	r=0.323 P=0.116
				HDRS	r=-0.026 P=0.903