

**Analysis of the time evolution of SARS-CoV-2 lethality rate in Italy:
Evidence of an unaltered virus potency**

Nicole Balasco¹, Vincenzo d'Alessandro², Giovanni Smaldone³, Luigi Vitagliano^{1,*}

¹Institute of Biostructures and Bioimaging, CNR, Naples, Italy.

²Department of Electrical Engineering and Information Technology, University Federico II, Naples, Italy.

³IRCCS SDN, Naples, Italy.

*Correspondence should be addressed to L.V. via Mezzocannone 16, 80134, Naples, Italy e-mail: luigi.vitagliano@unina.it

Supplementary Material




Figure S1. Percentage of positive tests over total tests from W_8 to W_{13} . Data are counted by omitting multiple tests made on the same person. The data used to generate the graph are reported in **Table S4** (individual Cases/Tests).

Table S1. Cases/deaths/tests daily collected from the Reports of the Italian National Institute of Health (ISS).

Date	Daily Cases	Daily Deaths	Daily Tests
20-Feb	0	0	
21-Feb	17	0	
22-Feb	47	1	
23-Feb	90	2	
24-Feb	72	4	4324
25-Feb	94	4	4299
26-Feb	147	1	964
27-Feb	185	5	2427
28-Feb	234	4	3681
29-Feb	239	8	2966
01-Mar	573	12	2466
02-Mar	335	11	2218
03-Mar	466	27	2511
04-Mar	587	28	3981
05-Mar	769	41	2525
06-Mar	778	49	3997
07-Mar	1247	36	5703
08-Mar	1492	133	7875
09-Mar	1797	97	3889
10-Mar	1577	168	6935
11-Mar	1713	196	12393
12-Mar	2651	189	12857
13-Mar	2547	250	11477
14-Mar	3497	175	11682
15-Mar	3590	368	15729
16-Mar	3385	349	13063
17-Mar	3374	345	10695
18-Mar	4207	475	16884
19-Mar	5322	427	17236
20-Mar	5986	627	24109
21-Mar	6557	793	26336
22-Mar	5560	651	25180
23-Mar	4790	601	17066
24-Mar	5249	743	21496
25-Mar	5210	683	27481
26-Mar	6153	712	36615
27-Mar	5909	919	33019
28-Mar	5974	889	35447
29-Mar	5217	756	24504
30-Mar	4050	812	23329
31-Mar	4053	837	29609
01-Apr	4782	727	34455

02-Apr	4668	760	39809
03-Apr	4585	766	38617
04-Apr	4805	681	37375
05-Apr	4316	525	34237
06-Apr	3599	636	30271
07-Apr	3039	604	33713
08-Apr	3836	542	51680
09-Apr	4204	610	46244
10-Apr	3951	570	53495
11-Apr	4694	619	56609
12-Apr	4092	431	46720
13-Apr	3153	566	36717
14-Apr	2972	602	26779
15-Apr	2667	578	43715
16-Apr	3786	525	60999
17-Apr	3493	575	65705
18-Apr	3491	482	61725
19-Apr	3047	433	50708
20-Apr	2256	454	41483
21-Apr	2729	570	52126
22-Apr	3370	401	63101
23-Apr	2646	464	66658
24-Apr	3021	420	62447
25-Apr	2357	415	65387
26-Apr	2324	260	49916
27-Apr	1739	333	32003
28-Apr	2019	382	57272
29-Apr	2086	323	63827
30-Apr	1872	285	68456
01-May	1965	269	74208
02-May	1900	474	55412
03-May	1389	174	44935
04-May	1221	195	37631
05-May	1075	236	55263
06-May	1444	369	64263
07-May	1401	274	70359
08-May	1327	243	63775
09-May	1083	194	69171
10-May	802	165	51678
11-May	744	179	40740
12-May	1402	172	67003
13-May	888	195	61973
14-May	992	262	71876
15-May	789	242	68176
16-May	875	153	69179

17-May	675	145	60101
18-May	451	99	36406
19-May	813	162	63158
20-May	665	161	67195
21-May	642	156	71679
22-May	652	130	75380
23-May	669	119	72410
24-May	531	50	55824
25-May	300	92	35241
26-May	397	78	57674
27-May	584	117	67324
28-May	593	70	75893
29-May	516	87	72135
30-May	416	111	69342
31-May	333	75	54118

Table S2. Week definition with starting and ending date.

Week	Starting Date	Ending Date
W ₀	24-Feb	01-Mar
W ₁	02-Mar	08-Mar
W ₂	09-Mar	15-Mar
W ₃	16-Mar	22-Mar
W ₄	23-Mar	29-Mar
W ₅	30-Mar	05-Apr
W ₆	06-Apr	12-Apr
W ₇	13-Apr	19-Apr
W ₈	20-Apr	26-Apr
W ₉	27-Apr	03-May
W ₁₀	04-May	10-May
W ₁₁	11-May	17-May
W ₁₂	18-May	24-May
W ₁₃	25-May	31-May

Table S3. Cases and deaths *per* week. The normalization was performed by dividing the actual values by the maximum of each ensemble.

Week	Average number of Cases	Average number of Deaths	Normalized number of Cases	Normalized number of Deaths
W ₀	221	5	0.0401	0.0072
W ₁	811	46	0.1474	0.0613
W ₂	2482	206	0.4512	0.2721
W ₃	4913	524	0.8932	0.6915
W ₄	5500	758	1.0000	1.0000
W ₅	4466	730	0.8119	0.9632
W ₆	3916	573	0.7120	0.7566
W ₇	3230	537	0.5872	0.7092
W ₈	2672	426	0.4858	0.5627
W ₉	1853	320	0.3369	0.4224
W ₁₀	1193	239	0.2169	0.3160
W ₁₁	909	193	0.1653	0.2542
W ₁₂	632	125	0.1149	0.1654
W ₁₃	448	90	0.0815	0.1188

Table S4. Cases/tests ratios and WLR values of the different shift schemes (0WS, 1WS, and 2WS). Cases/tests ratios and 0WS WLR values were calculated considering data derived from the same week. 1WS and 2WS WLR values were computed by dividing the deaths of a given week (W_i) by the cases of one week (W_{i-1}) or two weeks (W_{i-2}) before, respectively. Individual cases/tests were counted by omitting multiple tests made on the same person (Data were retrieved from <https://lab.gedidigital.it/gedi-visual/2020/coronavirus-i-contagi-in-italia>).

Week	Cases/Tests	Individual Cases/Tests	0WS WLR	1WS WLR	2WS WLR
W ₁	19.69		5.73		
W ₂	23.17		8.31	25.43	
W ₃	25.76		10.66	21.11	64.63
W ₄	19.68		13.77	15.42	30.53
W ₅	13.17		16.34	13.27	14.85
W ₆	8.60		14.63	12.83	10.42
W ₇	6.53		16.63	13.72	12.03
W ₈	4.66	7.23	15.95	13.20	10.88
W ₉	3.27	5.39	17.27	11.98	9.91
W ₁₀	2.03	4.46	20.06	12.92	8.96
W ₁₁	1.45	2.49	21.18	16.14	10.39
W ₁₂	1.00	1.67	19.83	13.78	10.50
W ₁₃	0.727	1.34	20.07	14.24	9.90

Table S5. Ratio of cases (or deaths) between consecutive weeks. Values were computed by dividing the cases (or deaths) of a given week (W_i) by the cases (or deaths) of one week (W_{i-1}) before. For example, the ratio #2 is calculated by dividing the Cases (or Deaths) of W_2 for those of W_1 .

Ratio	Cases	Deaths
2	3.062	4.440
3	1.980	2.541
4	1.120	1.446
5	0.812	0.963
6	0.877	0.785
7	0.825	0.937
8	0.827	0.793
9	0.693	0.751
10	0.644	0.748
11	0.762	0.804
12	0.695	0.651
13	0.710	0.718