

Supplementary Information

High levels of SARS-CoV-2 specific T-cells with restricted functionality in patients with severe course of COVID-19


David Schub¹, Verena Klemis¹, Sophie Schneitler², Janine Mihm³, Philipp M. Lepper⁴, Heinrike Wilkens⁴, Robert Bals⁴, Hermann Eichler⁵, Barbara C. Gärtner², Sören L. Becker², Urban Sester³, Martina Sester^{1,*}, and Tina Schmidt¹

¹Department of Transplant and Infection Immunology, ²Institute of Medical Microbiology and Hygiene, ³Department of Internal Medicine IV, ⁴Department of Internal Medicine V, ⁵Institute of Clinical Hemostaseology and Transfusion Medicine, Saarland University, 66421 Homburg, Germany

*Correspondence: martina.sester@uks.eu

Supplementary figures

Supplementary figure S1




Supplementary figure S1: Gating strategies for quantitation of lymphocyte populations. (A)

To quantify lymphocyte subpopulations, lymphocytes were gated within all events according to SSC-A and FSC-A. Exclusion of doublets was performed using high and area signal of FSC. Among single lymphocytes, B cells and T cells were identified according to expression of CD19 and CD3. T cells were further subdivided into helper T cells (CD4) and cytotoxic T cells (CD8). NK cells were determined by CD16 and/or CD56 expressing cells among single lymphocytes neither expressing CD19 nor CD3. **(B)** Among single lymphocytes, B cells were identified as CD19+CD3- and T cells were gated according to expression of CD19 and CD3. According to IgD-

and CD27-expression, B cells were subdivided in naïve (IgD+CD27-), non-switched memory (IgD+CD27+) or switched memory B cells (IgD-CD27+). Plasmablasts were identified among switched memory B cells by additional staining of CD38. As plasmablasts may also be extended in size and granularity compared with non-plasmablast, lymphocyte gates (FSC and SSC) were enlarged for quantitation of these cells. **(C)** CD3+ T cells were identified among single lymphocytes and further subdivided in CD4 and CD8 T cells. Both subpopulations were analyzed towards expression of Ki67 (percentage of positive cells) and CTLA-4 (MFI of CD4 or CD8 T cells). **(D)** CD4 and CD8 T cells were identified among total T cells (CD3) and analyzed towards expression of PD-1 (PD1 MFI). CTLA-4, cytotoxic T lymphocyte antigen 4; FSC, forward scatter; PD-1, programmed cell death 1; SSC, side scatter.


Supplementary figure S2



Supplementary figure S2: Gating strategy for analysis of cytokine and CTLA-4 expression of antigen-specific T cells. (A) Lymphocytes were gated among total events according to size (FSC) and granularity (SSC) followed by exclusion of doublets using high and area signal of FSC. (B) To ensure accurate gating of lymphocytes, CD4/CD8 backgating was used which allows for an highlighted visualization of this population. (C) Among single lymphocytes, CD4 T cells

(fine) were identified by exclusion of CD4-negative and CD8-positive cells. **(D)** Similarly, CD8 T cells (fine) were identified among single lymphocytes by exclusion of CD8-negative and CD4-positive cells. To determine the percentage of antigen-specific cells, CD69+IFN γ + T cells were identified among CD4 **(E)** and CD8 T cells **(F)**. CD69+IFN γ + CD4 or CD8 T cells were subdivided according to expression of TNF α and IL-2. Among CD4 or CD8 T cells not coexpressing CD69 and IFN γ , „AND/OR-gating“ of CD69+TNF α + and CD69+IL2+ CD4 or CD8 T cells allowed for identification of cells with single or combined expression of TNF α and IL-2. In addition expression of CTLA-4 (MFI) of CD69+IFN γ + CD4 and CD8 T cells was analyzed. IFN, interferon; IL, interleukin; TNF, tumor necrosis factor.

Supplementary figure S3



Supplementary Figure S3: Gating strategy for analysis of Ki67-, Granzyme B- and PD-1-expression of antigen-specific T cells. Among CD4 (A) and CD8 T cells (B) (fine gating s. supplementary figure S2), antigen-specific cells were identified by co-expression of CD69 and IFN γ . These cells were analyzed towards their expression of Ki67 and/or Granzyme B (percentage of positive cells) as well as expression of PD-1 (MFI). IFN, interferon; PD-1, programmed cell death 1.

Supplementary Table

Supplementary Table S1: Antibodies for flow cytometric analysis

Antigen	conjugate	clone	isotype	reactivity
CD3	APC, PerCP	SK7	IgG1 k	mouse anti-human
CD4	PE-Cy7, APC-H7	SK3	IgG1 k	mouse anti-human
CD8	V500, PerCP	RPA-T8, SK1	IgG1 k	mouse anti-human
CD16	PE	3G8	IgG1 k	mouse anti-human
CD19	FITC	HIB19	IgG1 k	mouse anti-human
CD27	APC	L128	IgG1 k	mouse anti-human
CD38	PE	HB7	IgG1 k	mouse anti-human
CD56	PE	B159	IgG1 k	mouse anti-human
CD69	PE-Cy7	L78	IgG1 k	mouse anti-human
CD152 (CTLA-4)	PE, APC	BNI3	IgG2a k	mouse anti-human
CD279 (PD-1)	APC, PE	MIH4	IgG1 k	mouse anti-human
Granzyme B	V450	GB11	IgG1 k	mouse anti-human
IFN γ	FITC	4S.B3	IgG1 k	mouse anti-human
IgD	PE-Cy7	IA6-2	IgG2a k	mouse anti-human
IL-2	PE	MQ1-17H12	IgG2a k	rat anti-human
Ki67	AlexaFluor 647	B56	IgG1 k	mouse anti-human
TNF α	V450	MAB11	IgG1 k	mouse anti-human

All antibodies were purchased from BD.