

Supplementary Material

Supplementary Tables

Supplementary Table 1. Sample characteristics. Values are reported as number and percentage. Ethnicity, smoking, disease status and medications were compared using Pearson's χ^2 test, as implemented in the *prop.test* function (*stats* R package, v 3.6.0). "Users tested" refers to individuals self-reporting a positive or negative swab test result. "Symptomatic untested users" refers to individuals who reported at least one of the 16 collected symptoms, did not believe that they had already had COVID-19 when registering with the app, and had not been tested for SARS-CoV-2. "Classic symptoms" refer to those included in the NHS guidelines (*i.e.*, fever, persistent cough, and/or anosmia), whose presence would require isolation and testing for SARS-CoV-2 infection.

	All users	Users tested for SARS-CoV-2				Symptomatic untested users			P
		All	Positive	Negative	P	All	With classical symptoms	Without classic symptoms	
N	336,847	27,157	2,021	25,136		54,652	17,371	37,281	
Ethnicity					4.72x10 ⁻⁶				<2e-16
Asian	6,503 (1.9%)	646 (2.4%)	70 (3.5%)	576 (2.3%)		1,029 (1.9%)	422 (2.4%)	607 (1.6%)	
Black	2,105 (0.6%)	232 (0.9%)	34 (1.7%)	198 (0.8%)		370 (0.7%)	143 (0.8%)	227 (0.6%)	
Chinese	1,029 (0.3%)	87 (0.3%)	6 (0.3%)	81 (0.3%)		134 (0.2%)	43 (0.2%)	91 (0.2%)	
Middle East	1,204 (0.4%)	114 (0.4%)	13 (0.6%)	101 (0.4%)		187 (0.3%)	84 (0.5%)	103 (0.3%)	
Mixed	6,974 (2.1%)	526 (1.9%)	35 (1.7%)	491 (2.0%)		1,295 (2.4%)	508 (2.9%)	787 (2.1%)	
White	316,567 (94.0%)	25,341 (93.3%)	1,840 (91.0%)	23,501 (93.5%)		51,188 (93.7%)	16,021 (92.2%)	35,167 (94.3%)	
N/A	2,465 (0.7%)	211 (0.8%)	23 (1.1%)	188 (0.7%)		449 (0.8%)	150 (0.9%)	299 (0.8%)	
Smoking status					0.04				<2e-16
Never	241,093 (71.6%)	18,499 (68.1%)	1,396 (69.1%)	17,103 (68.0%)		37,932 (69.4%)	11,752 (67.7%)	26,180 (70.2%)	
Ex	62,563 (18.6%)	5,655 (20.8%)	436 (21.6%)	5,219 (20.8%)		10,726 (19.6%)	3,193 (18.4%)	7,533 (20.2%)	
Current	33,176 (9.8%)	3,002 (11.1%)	189 (9.4%)	2,813 (11.2%)		5,994 (11.0%)	2,426 (14.0%)	3,568 (9.6%)	
Has diabetes (%)	12,884 (3.8%)	1,119 (4.4%)	113 (5.6%)	1,078 (4.3%)	7.04x10 ⁻³	2,109 (3.9%)	704 (4.1%)	1,405 (3.8%)	0.11
Has heart disease (%)	10,052 (3.0%)	936 (3.4%)	78 (3.9%)	858 (3.4%)	0.32	1,663 (3.0%)	529 (3.0%)	1,134 (3.0%)	1.00

Has lung disease (%)	31,489 (9.3%)	3,150 (11.6%)	264 (13.1%)	2,886 (11.5%)	3.58×10^{-2}	6,912 (12.6%)	2,603 (15.0%)	4,309 (11.6%)	3.71×10^{-29}
Has kidney disease (%)	2,598 (0.8%)	293 (1.1%)	29 (1.4%)	264 (1.1%)	0.13	527 (1.0%)	169 (1.0%)	358 (1.0%)	0.93
Has cancer (%)	4,456 (1.3%)	543 (2.0%)	34 (1.7%)	509 (2.0%)	0.33	660 (1.2%)	210 (1.2%)	450 (1.2%)	1.00
Corticosteroids (%)	24,583 (7.3%)	2,491 (9.2%)	190 (9.4%)	2,301 (9.2%)	0.74	5,601 (10.2%)	2,028 (11.7%)	3,573 (9.6%)	6.95×10^{-14}
Immunosuppressants (%)	12,006 (3.6%)	1,138 (4.2%)	75 (3.7%)	1,063 (4.2%)	0.29	2,415 (4.4%)	910 (5.2%)	1,505 (4.0%)	2.26×10^{-10}
Blood pressure medications (%)	44,061 (13%)	3,593 (13.2%)	273 (13.5%)	3,320 (13.2%)	0.73	6,272 (11.5%)	1,852 (10.7%)	4,420 (11.9%)	4.80×10^{-5}

Supplementary Table 2. Sensitivity analysis for multivariate logistic regression. “All tested” refers to individuals self-reporting a positive or negative swab test result. “All tested, no healthcare workers” refers to individuals self-reporting a positive or negative swab test result but not declaring to be healthcare workers. “Symptomatic untested users” refers to individuals who reported at least one of the 16 collected symptoms, did not believe that they had already had COVID-19 when registering with the app, and had not been tested for SARS-CoV-2.

	Model	N	OR	95% CI	P value
	All tested	26,945	1.65	1.37-1.99	1.30x10 ⁻⁷
	All tested, no healthcare workers	19,516	1.52	1.15-2.02	3.72x10 ⁻³
Body rash	Only males tested	10,598	1.52	1.04-2.21	0.03
	Only females tested	16,347	1.71	1.38-2.12	9.59x10 ⁻⁷
	Symptomatic untested users	54,203	1.46	1.35-1.58	2.92x10 ⁻²⁰
	All tested	26,945	1.73	1.32-2.27	7.25x10 ⁻⁵
	All tested, no healthcare workers	19,516	1.83	1.25-2.66	1.69x10 ⁻³
Acral rash	Only males tested	10,598	1.15	0.66-1.99	0.63
	Only females tested	16,347	2.05	1.49-2.80	8.14x10 ⁻⁶
	Symptomatic untested users	54,203	1.08	0.96-1.22	0.22