

Supplemental Tables

Room	Days between admission and sampling	Days between first COVID-19 positive and sampling	Days between most recent COVID-19 positive and sampling	Notes
Room 7A	24	24	1	
Room 7B	7	>10	0 (same day)	transfer from other facility
Room 5A	3	3	3	
Room 5B	2	2	2	
Room 5C	4	3,2	3,2	2 patients in this room
Room 5D	6	6	6	

Table S1. Metadata for rooms sampled in this study.

PCR Reaction Mix	
Nuclease Free Water	6.1 µL
Invitrogen 2X Master Mix	12.5 µL
MgSO4 (50mM)	0.4 µL
Primer/Probe Mix (IDT)* (Primers 10uM, Probe 5uM)	0.5 µL
SuperScript III Platinum Taq	0.5 µL
extracted sample RNA, nuclease free water or positive control	5.0 µL
*E gene target primers and probe:	
5'/56-FAM/ACACTAAGCC/ZEN/ATCCTTACTGCGCTTCG/3AIBkFG/-3'	Probe
5'-ATATTGCAGCAGTACGCACACA-3'	Primer 1
5'-ACAGGTACGTTAATAGTTAATAGCGT-3'	Primer 2

Table S2. PCR Reaction Mix used for all PCR of SARS-CoV-2 throughout the study.

Process	Reagents	Conditions
Fixing	2% paraformaldehyde in 0.1M Sorenson's phosphate buffer (pH 7.2)	24 hours at 4 °C
Washing	0.1M Sorenson's phosphate buffer (pH 7.2)	3 times
Post Fix	1% aqueous solution of osmium tetroxide	30 minutes
Dehydration series	50,70, 90, 95 and 100% ethanol	
Embedding	50:50 Embed 812 Resin and propylene oxide	Overnight
Final Embedding	Fresh Embed 812 Resin	2 hours at room temperature followed by polymerization at 65 °C for 24 hours
Sectioning	100 nm sections stained with 2% Uranyl Acetate followed by Reynolds Lead Citrate	Cut with Leica UC6 Ultracut ultramicrotome

Table S3. Sample preparation for transmission electron microscopy.

Room	Sample Type	Airborne Viral Concentration TCID50/cm ³ of Air	Standard Deviation	Day 1 TCID50/mL	Day 1 Standard Deviation	Final Day TCID50/mL	Final Day Standard Deviation	Percent Increase in TCID50/mL in Cell Culture	Significance of Calculated Increase (P-value)
Room 7A	BC251 <1 µm	5.50	4.98	0.07	0.04	0.08	0.05	23.78	0.32
Room 7A	BC251 1-4 µm	0.45	0.50	0.04	0.00	0.08	0.03	87.49	0.08
Room 7A	BC251 4.1 µm	1.57	1.48	0.05	0.02	0.10	0.07	81.17	0.24
Room 7B	BC251 <1 µm	5.94	6.79	0.07	0.02	0.10	0.04	44.57	0.03
Room 7B	BC251 1-4 µm	0.40	0.35	0.07	0.07	0.03	0.02	17.30	0.39
Room 7B	BC251 4.1 µm	2.45	2.62	0.05	0.01	0.09	0.04	82.25	0.13
Room 5A	BC251 <1 µm	7.41	2.03	0.05	0.01	0.08	0.01	66.04	0.03
Room 5A	BC251 1-4 µm	0.36	0.32	0.03	0.03	0.06	0.03	89.99	0.18
Room 5A	BC251 4.1 µm	1.34	1.17	0.07	0.01	0.06	0.02	-23.53	NA
Room 5B	BC251 <1 µm	3.73	3.23	0.09	0.05	0.14	0.03	45.47	0.13
Room 5B	BC251 1-4 µm	0.26	0.23	0.08	0.01	0.06	0.02	-26.55	NA
Room 5B	BC251 4.1 µm	1.57	2.71	0.28	0.03	0.09	0.04	-68.55	NA
Room 5C	BC251 <1 µm	4.10	2.44	0.06	0.02	0.22	0.02	260.65	0.01
Room 5C	BC251 1-4 µm	0.58	0.02	0.05	0.05	0.09	0.03	79.25	0.09
Room 5C	BC251 4.1 µm	1.97	3.41	0.08	0.01	0.07	0.01	-4.77	NA
Room 5D	BC251 <1 µm	1.18	2.05	0.04	0.02	0.00	0.00	-100.00	NA
Room 5D	BC251 1-4 µm	0.65	0.25	0.06	0.04	0.07	0.07	10.65	0.45
Room 5D	BC251 4.1 µm	4.00	2.17	0.09	0.02	0.09	0.05	-0.56	NA

Table S4. Complete data set from Figure 1, indicating viral concentration derived from RT-PCR of raw samples, RT-PCR-derived viral concentrations from cell culture experiments and calculations on percent increase in viral titre and statistical significance, (p) as determined by a one-tailed Student's t-test.

Room	Small Mode			Large Mode		
	M	σ_g	d_{pg}	M	σ_g	d_{pg}
Room 5A	3.5E-04	1.3E+00	6.4E-01	4.4E-03	1.9E+00	4.2E+00
Room 5B	3.7E-04	1.3E+00	6.8E-01	4.5E-03	1.7E+00	3.8E+00
Room 5C	3.9E-04	1.2E+00	6.7E-01	6.8E-03	2.3E+00	4.3E+00
Room 5D	2.4E-04	1.2E+00	6.7E-01	3.1E-03	3.0E+00	3.9E+00
Room 7A	2.7E-05	1.3E+00	6.9E-01	2.1E-03	2.5E+00	1.1E+01
Room 7B	2.9E-04	1.3E+00	8.0E-01	5.8E-03	1.6E+00	2.3E+00

Table S5. Log-normal parameters calculated for the large and small modes shown in Figure 4.

	Location	PCR (TCID50/cm ³)	Measured (µg/cm ³)	Small Mode (µg/cm ³)	Large Mode (µg/cm ³)
<1 µm	Room 5A	7.41E+00	7.95E-03	4.79E-03	3.11E-03
	Room 5B	3.73E+00	6.28E-03	3.05E-03	3.17E-03
	Room 5C	4.10E+00	4.33E-03	3.63E-03	6.45E-04
	Room 5D	1.18E+00	4.61E-03	4.09E-03	5.00E-04
	Room 7A	5.50E+00	4.00E-04	3.05E-04	1.34E-04
	Room 7B	5.94E+00	6.90E-03	3.16E-03	3.45E-03
1 - 4 µm	Room 5A	3.57E-01	4.36E-02	8.50E-05	3.96E-02
	Room 5B	2.61E-01	1.90E-02	1.53E-05	1.68E-02
	Room 5C	5.80E-01	2.88E-02	2.19E-04	2.71E-02
	Room 5D	6.47E-01	3.76E-02	2.65E-04	3.54E-02
	Room 7A	4.53E-01	5.25E-03	2.30E-05	4.51E-03
	Room 7B	4.00E-01	3.76E-02	2.65E-04	3.54E-02
>4.1 µm	Room 5A	1.34E+00	4.59E-02	1.03E-23	3.72E-02
	Room 5B	1.57E+00	1.58E-02	1.39E-32	1.29E-02
	Room 5C	1.97E+00	3.03E-02	9.44E-15	2.86E-02
	Room 5D	4.00E+00	3.13E-02	6.93E-16	2.80E-02
	Room 7A	1.67E+00	1.29E-02	9.29E-17	1.08E-02
	Room 7B	2.74E+00	3.13E-02	6.93E-16	2.80E-02

Table S6. RT-PCR derived virus concentration and recovered mass concentrations for each stage of the BC251 sampler in each room included in this study.

SARS-CoV-2 Detected in Aerosol by PCR and Cell Culture		
	rRT-PCR	Cell Culture
>4.1 µm	6:6 (100%)	0:6 (0%)
1-4 µm	6:6 (100%)	2:6 (33%)*
< 1 µm	6:6 (100%)	3:6 (50%)

Table S7. Number and percentage of aerosol samples where SARS-CoV-2 was detected by rRT-PCR and where replication was observed in cell culture. Samples from the 1-4 µm stage did reach the level of statistical significance ($0.05 < P < 0.10$).