

*Supplemental Table 1: Univariable logistic regression results with in-hospital mortality as the outcome*

<b>Variable</b>	<b>Univariable Odds Ratio (95% CI)</b>	<b>P-value</b>
Age (Continuous)	1.06 (1.05-1.06)	1.83E-64
Age <55	Ref	Ref
Age 55-64	2.43 (1.78-3.31)	2.19E-08
Age 65-74	4.1 (3.05-5.47)	3.77E-21
Age >=75	8.25 (6.22-10.9)	1.82E-49
Male	1.02 (0.868-1.19)	0.831
Race: Non-Hispanic White	Ref	Ref
Race: Non-Hispanic Black	0.733 (0.589-0.913)	0.00561
Race: Hispanic	0.606 (0.486-0.755)	8.31E-06
Race: Asian	0.769 (0.518-1.14)	0.192
Race: Other	0.85 (0.66-1.1)	0.211
Hypertension	1.29 (1.1-1.52)	0.00174
Diabetes	1.3 (1.09-1.56)	0.00339
Coronary Artery Disease	1.87 (1.5-2.32)	1.87E-08
Heart Failure	1.62 (1.22-2.15)	0.000929
Atrial Fibrillation	2.28 (1.7-3.04)	2.47E-08
Chronic Kidney Disease	1.5 (1.19-1.88)	0.000503
COPD/Asthma	0.941 (0.71-1.25)	0.67
Obesity	0.825 (0.613-1.11)	0.205
Cancer	1.22 (0.904-1.66)	0.19
Initial temperature	0.91 (0.867-0.954)	0.000116
Initial heart rate	1 (0.997-1)	0.654
Initial systolic blood pressure	0.998 (0.995-1)	0.3
Initial respiratory rate	1.07 (1.06-1.09)	1.60E-21
Initial hypoxia (O <sub>2</sub> <92%)	2.57 (2.16-3.05)	3.73E-27
Manhattan facility	0.483 (0.412-0.566)	3.35E-19
Current or former smoker	1.12 (0.94-1.34)	0.202


*Supplemental Table 2: Multivariable logistic regression results for inflammatory markers as predictors of in-hospital mortality, adjusting for age, comorbidities, and initial oxygen saturation, stratified by sex. The unstratified model includes sex. Interaction P-values indicate the significance of the interaction term between sex and the lab value in a multivariable model incorporating the interaction. Lab values were standardized prior to stratification and regression; odds ratios represent the change in odds associated with a one-standard-deviation increase in the lab value.*

<b>Lab test</b>	<b>Male Odds Ratio (95% CI)</b>	<b>Female Odds Ratio Ratio (95% CI)</b>	<b>Unstratified Odds ratio (95% CI)</b>	<b>Interaction P-value</b>
<b>WBC</b>	1.213 (0.979-1.5)	1.751 (1.26-2.44)	1.33 (1.11-1.59)	0.0522
<b>ALBUMIN</b>	0.668 (0.524-0.85)	0.834 (0.645-1.08)	0.754 (0.634-0.896)	0.475
<b>DDIMER</b>	1.176 (0.965-1.43)	1.465 (1.07-2)	1.214 (1.03-1.43)	0.24
<b>FERRITIN</b>	0.947 (0.776-1.16)	1.66 (1.2-2.29)	1.1 (0.944-1.28)	0.00241
<b>PROCALCITONIN</b>	2.34 (1.19-4.61)	1.046 (0.85-1.29)	1.297 (1.06-1.59)	0.0335
<b>LDH</b>	1.26 (1.01-1.57)	2.504 (1.66-3.79)	1.534 (1.25-1.89)	0.00317
<b>CRP</b>	1.66 (1.32-2.09)	1.246 (0.974-1.59)	1.451 (1.23-1.71)	0.224
<b>IL1B</b>	1.035 (0.769-1.39)	0.957 (0.672-1.36)	0.982 (0.776-1.24)	0.719
<b>IL6</b>	1.774 (1.26-2.49)	1.324 (0.971-1.8)	1.496 (1.21-1.85)	0.177
<b>IL8</b>	3.706 (1.31-10.6)	6.234 (1.22-31.7)	3.857 (1.65-9.01)	0.876
<b>TNFa</b>	0.881 (0.502-1.55)	2.252 (0.72-7.05)	1.073 (0.885-1.3)	0.204


*Supplemental Table 3: Multivariable logistic regression results for inflammatory markers as predictors of in-hospital mortality, adjusting for age, sex, comorbidities, stratified by initial oxygen saturation. The unstratified model includes initial oxygen saturation. Interaction P-values indicate the significance of the interaction term between sex and the lab value in a multivariable model incorporating the interaction. Lab values were standardized prior to stratification and regression; odds ratios represent the change in odds associated with a one-standard-deviation increase in the lab value.*





<b>Lab Test</b>	<b>Normoxic Odds Ratios (95% CI)</b>	<b>Hypoxic Odds Ratios (95% CI)</b>	<b>Unstratified Odds Ratios (95% CI)</b>	<b>Interaction P-value</b>
<b>WBC</b>	1.392 (1.13-1.71)	1.255 (0.911-1.73)	1.33 (1.11-1.59)	0.463
<b>ALBUMIN</b>	0.682 (0.555-0.837)	0.927 (0.656-1.31)	0.754 (0.634-0.896)	0.095
<b>DDIMER</b>	1.106 (0.891-1.37)	1.543 (1.12-2.12)	1.214 (1.03-1.43)	0.169
<b>FERRITIN</b>	1.458 (1.19-1.78)	0.702 (0.48-1.03)	1.1 (0.944-1.28)	0.000256
<b>PROCALCITONIN</b>	1.513 (1.09-2.1)	1.165 (0.94-1.44)	1.297 (1.06-1.59)	0.145
<b>LDH</b>	2.094 (1.6-2.75)	1.089 (0.896-1.32)	1.534 (1.25-1.89)	7.62E-05
<b>CRP</b>	1.709 (1.37-2.13)	1.033 (0.768-1.39)	1.451 (1.23-1.71)	0.00481
<b>IL1B</b>	0.969 (0.759-1.24)	1.052 (0.408-2.71)	0.982 (0.776-1.24)	0.979
<b>IL6</b>	2.024 (1.43-2.86)	1.177 (0.866-1.6)	1.496 (1.21-1.85)	0.0134
<b>IL8</b>	3.706 (1.37-9.95)	5.989 (1.45-24.6)	3.857 (1.65-9.01)	0.786
<b>TNFa</b>	0.942 (0.627-1.42)	2.746 (0.648-11.7)	1.073 (0.885-1.3)	0.449


Supplemental Figure 1: Flow diagram of included patients



Supplemental Figure 2: Baseline lab values among patients admitted to The Mount Sinai Hospital, by sex. (A) White blood cells. (B) Procalcitonin. (C) lactate dehydrogenase. (D) C-reactive protein. (E) Ferritin, absolute values. (F) Ferritin, relative to upper limit of normal. (G) Interleukin-8. (H) Interleukin-6. (I) Tumor necrosis factor alpha. (J) Interleukin-1 beta. Dotted lines represent reference ranges. Female-specific reference ranges are in red while male-specific reference ranges are in blue. Significance levels are indicated by ns:  $p > 0.05$ , \*\*:  $p \leq 0.01$ , \*\*\*:  $p \leq 0.001$ ; \*\*\*\*:  $p \leq 0.0001$ .



**E****F****G****H**

**I****J**