

SUPPLEMENTAL DATA

Table S1. Job Role Classifier, Odds Ratios and 95% Confidence Intervals (in descending order)

Role	Seroprevalence	Reference Set	Odds Ratio (95% CI)
Housekeeper	8.03% (11/137)	2.46% (241/9800)	3.17 (1.59-5.71)
Radiologist	7.27% (4/55)	2.51% (248/9882)	3.09 (0.93-7.69)
Medical Assistant	3.09% (21/680)	2.50% (231/9257)	1.41 (0.86-2.18)
Pharmacist	3.25% (5/154)	2.52% (247/9783)	1.34 (0.47-2.96)
Social Work	3.4% (8/235)	2.51% (244/9702)	1.29 (0.58-2.47)
Clerical, administrative	3.15% (25/793)	2.48% (227/9144)	1.27 (0.81-1.90)
Physician	3.12% (32/1026)	2.47% (220/8911)	1.21 (0.81-1.75)
Management, director	3.11% (8/257)	2.52% (244/9680)	1.14 (0.51-2.19)
Surgeon	2.66% (8/301)	2.53% (244/9636)	0.97 (0.43-1.89)
Nursing	2.34% (82/3504)	2.64% (170/6433)	0.90 (0.68-1.18)
Technician	1.98% (25/1260)	2.62% (227/8677)	0.77 (0.49-1.15)
Respiratory	2.06% (2/97)	2.54% (250/9840)	0.74 (0.12-2.37)
Therapist	1.78% (11/617)	2.59% (241/9320)	0.69 (0.35-1.21)
Food Service	1.77% (2/113)	2.54% (250/9824)	0.64 (0.11-2.04)
Researcher	1.45% (1/69)	2.54% (251/9868)	0.62 (0.03-2.82)
NP, PA	1.49% (6/402)	2.58% (246/9535)	0.58 (0.23-1.20)
Facilities, engineer	1.16% (1/86)	2.55% (251/9851)	0.40 (0.02-1.83)
Security	0.00% (0/41)	2.55% (252/9896)	0.00 (0.00-36.01)
Anesthesiologist	0.00% (0/110)	2.56% (252/9827)	0.00 (0.00-0.26)

Notes:

- Only one role classifier assigned per participant.
- Social Work includes case managers and chaplaincy.
- Pharmacist includes pharmacy technicians.
- Medical Assistant includes scribes.
- Physician includes all except surgeon, anesthesiologist, radiologist.
- Nursing includes nurse assistants.
- Respiratory (non-physician) indicates respiratory therapists.
- Therapist includes all other therapists (e.g. PT, OT, Speech, Dietician, Massage, etc . . .).
- NP, PA = nurse practitioner, physician assistant.
- Facilities includes information systems/technology.

Table S2. Workplace Classifier, Odds Ratios and 95% Confidence Intervals (in descending order)

Workplace	Seroprevalence	Reference Set	Odds Ratio (95% CI)
Research	3.09% (3/97)	2.48% (292/11757)	1.30 (0.32-3.49)
ICU	2.77% (22/795)	2.47% (273/11059)	1.16 (0.73-1.76)
Pharmacy	2.69% (6/223)	2.48% (289/11631)	1.11 (0.43-2.30)
OR	2.77% (36/1299)	2.45% (259/10555)	1.07 (0.74-1.51)
Desk job	2.74% (25/913)	2.47% (270/10941)	1.07 (0.68-1.58)
Inpatient wards	2.51% (55/2195)	2.48% (240/9659)	1.04 (0.77-1.39)
Radiology	2.51% (9/358)	2.49% (286/11496)	1.00 (0.47-1.84)
L&D	2.44% (15/614)	2.49% (280/11240)	0.99 (0.56-1.62)
Outpatient clinic	2.47% (70/2836)	2.50% (225/9018)	0.98 (0.74-1.28)
Behavioral Health	2.38% (6/252)	2.49% (289/11602)	0.96 (0.37-1.98)
Emergency	1.95% (17/870)	2.53% (278/10984)	0.81 (0.47-1.29)
Cardiology	1.71% (6/350)	2.51% (289/11504)	0.70 (0.27-1.44)
Rehab	1.66% (4/241)	2.51% (291/11613)	0.65 (0.20-1.55)
Pathology	1.55% (7/453)	2.53% (288/11401)	0.63 (0.27-1.25)
Pulmonary	1.48% (2/135)	2.50% (293/11719)	0.56 (0.09-1.76)
Hospital facilities	1.16% (1/86)	2.50% (294/11768)	0.40 (0.02-1.85)

Notes:

- Participants could designate up to three workplace settings of potential exposure, but only one role (e.g. job role = portable X-ray tech; workplace = ICU, ER, Inpatient Wards).
- The ICU classifier includes all intensive care units, aside from neonatal.
- The OR classifier includes operating rooms, recovery units and all interventional procedure suites (e.g. endoscopy, bronchoscopy, interventional radiology, etc . .).
- The L&D classifier includes labor & delivery, fetal-maternal, neonatal intensive care, etc . . .
- The Rehab classifier includes wound care and hyperbaric units.
- Hospital facilities applies to engineering, security, information systems/technology.

Figure S1. Providence Oregon Cohort Asymptomatic HCW Serosurveillance by Hospital Test Site

