## **Supplementary Material**

## Methods

**Table S1.** Age distribution and links to websites of the cohorts included in the training sample.

Sample	Source	Min	Max	Mean	SD
	https://camcan-archive.mrc-				
CAMCAN	cbu.cam.ac.uk/dataaccess/	18	87	54.26	18.36
DLBS	http://fcon_1000.projects.nitrc.org/	20.57	88.14	53.72	20.24
ds000222	https://openfmri.org/	23	73	47.73	20.01
HCP	https://www.humanconnectome.org/	22	37	27.87	3.59
	http://brain-development.org/ixi-				
IXI	dataset/32/	20.07	86.2	46.68	16.37
OASIS	http://www.oasis-brains.org/	18	90	48.23	24.68
SALD	http://fcon_1000.projects.nitrc.org/	20	80	45.3	17.41
STROKEMRI	Local sample (L.T.W.)	20	92	63.94	14.43

## Results

**Table S2.** Pearson correlation between estimated brain age and chronological age within the training sample, and the calculated mean absolute error (MAE) and root mean squared error (RMSE) for the different models.

Models	r	rmse	MAE		
Fullbrain	.93	7.57	5.76		
Occipital	.76	12.94	1.05		
Temporal	.85	1.49	8.08		
Frontal	.88	9.55	7.19		
Parietal	.85	1.43	7.92		
Cingulate	.78	12.61	9.79		
Insula	.79	12.22	9.49		
Subcortical	.90	8.54	6.46		

**Table S3.** Differences in regional brain age estimates between AAS-users with >2 years of AAS use and WLC adjusted for age and age2. Mean values are corrected brain age gap and are presented as estimated marginal means

	WLC (n	=102)	AAS users (n=123)					
Brain Age Gap	Mean	SE	Mean	SE	F	Sig.	FDR	P Eta S
Fullbrain	-1.96	.69	1.55	.62	14.282	<.001	<.001	.061
Occipital	-1.50	.89	1.37	.81	5.613	.019	.022	.025
Temporal	-1.36	.78	1.16	.71	5.729	.018	.029	.025
Frontal	-2.26	.88	1.76	.80	11.358	.001	.004	.049
Parietal	-1.27	.77	1.22	.70	5.676	.018	.024	.025
Cingulate	-1.63	.90	1.43	.82	6.343	.012	.024	.028
Insula	-1.55	.73	1.17	.66	7.63	.006	.016	.033
Subcortical	-1.07	.80	.93	.73	3.447	.065	.065	.015

**Table S4.** Differences in regional brain age estimates between AAS-users and weightlifting controls when adjusting for age, age2, general cognitive abilities, drug and alcohol use and depression. Mean values are corrected brain age gap presented as estimated marginal means. Note that this model is based upon a smaller sample size as the clinical variables were not available from all participants.

	WLC (n=89)		AAS users (n=89)					
	]							
Brain Age Gap	Mean	SE	Mean	SE	F	Sig.	FDR	P Eta S
Fullbrain	-2.07	.79	1.64	.79	9.363	.003	.024	.052
Occipital	91	1.09	.10	1.09	.358	.550	.550	.002
Temporal	-1.31	.95	.58	.95	1.673	.198	.396	.01
Frontal	-2.17	1.03	1.11	1.03	4.235	.041	.109	.024
Parietal	-1.07	.87	24	.87	.382	.537	.610	.002
Cingulate	-1.73	1.05	.24	1.05	1.474	.226	.362	.009
Insula	-1.15	.85	.02	.85	.803	.372	.614	.005
Subcortical	-1.06	.95	2.05	.95	4.550	.034	.136	.026