

Supplementary material

Onset, duration, and persistence of taste and smell changes and other COVID-19 symptoms: longitudinal study in Israeli patients

Hadar Klein^{1#}, Kim Asseo^{1#}, Noam Karni^{2#}, Yuval Benjamini³, Ran Nir-Paz⁴, Mordechai Muszkat², Sarah Israel^{2§}, Masha Y. Niv^{1§*}

#equal contributions as first authors

§equal contributions as senior authors

*corresponding author

¹The Institute of Biochemistry, Food and Nutrition, The Hebrew University, Rehovot, Israel

²Department of Medicine, Hadassah University Hospital, Mt. Scopus Campus, Jerusalem, Israel

³Department of Statistics, The Hebrew University, Mt. Scopus Campus, Jerusalem, Israel

⁴Department of Clinical Microbiology and Infectious Diseases, Hadassah-Hebrew University medical center

27 symptoms:		
≥50%	<50%	
<u>Taste change</u>	<u>Productive cough</u>	Bad taste in mouth
<u>Smell change</u>	<u>Runny nose</u>	Voice change
<u>Fever</u>	<u>Sore throat</u>	Eyes burn
<u>Dry cough</u>	Chemesthesis change	Dizziness
<u>Muscle aches</u>	<u>Diarrhea</u>	Vision changes
<u>Headache</u>	Chest pain	Coated tongue
<u>Lack of appetite</u>	Abdominal pain	Ears pressure
	<u>Breath difficulty</u>	Lacrimation
	<u>Vomit/nausea</u>	Hearing change
	<u>Fatigue</u>	Eyes discharge

Figure S1. 27 symptoms of the questionnaire. Symptoms with prevalence higher than 50% are presented with orange background, and those with prevalence lower than 50% are presents with light blue background. Official CDC symptoms are bolded. Symptoms that were included in the final analysis are underlined.




Figure S2. Distributions of symptoms' order of appearance. Y axis shows number of patients (n) and X-axis is order of appearance of the specified symptom (first symptom to appear or followed other symptoms).




Figure S3. Persistent symptoms after recovery (13 days post initiation of symptoms). Y axis list symptoms and X axis shows prevalence of symptoms' persistent post recovery.




Figure S4. Correlation between taste change severity and duration (n=76). Y axis shows taste change duration (days) and X axis shows taste change severity.