

Supplementary Material

Table S1. Difference-in Difference (DiD) regression results (Michigan)

| | AS-ME | | | Influenza Vaccine | | Spacer | | | Nebulizer | | |
|----------------------|-------------------------------|-------------------------|------------------------|--------------------------------|-------------------------|---------------------------------|------------------------|-------------------------|-------------------------------|-------------------------|------------------------|
| | Utilization Cost ⁺ | ED Visits ⁺⁺ | IP Visits ⁺ | Utilization Cost ⁺⁺ | PCP Visits ⁺ | Utilization Cost ⁺⁺⁺ | ED Visits ⁺ | IP Visits ⁺⁺ | Utilization Cost ⁺ | ED Visits ⁺⁺ | IP Visits ⁺ |
| | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. |
| (Intercept) | 1.991*** | 0.101*** | 0.008*** | 0.888*** | 0.236*** | 1.677*** | 0.106*** | 0.004** | 0.924*** | 0.015*** | 0.001 |
| Urban | 1.352*** | 0.176*** | 0.017*** | 0.695*** | 0.082*** | 1.129*** | 0.189*** | | 0.716*** | 0.049*** | 0.010*** |
| Suburban | 0.412*** | 0.037*** | | 0.176*** | 0.041*** | 0.334*** | 0.033*** | | 0.254*** | 0.008* | 0.004*** |
| Age 9 - 17 | -0.463*** | -0.074*** | -0.008*** | -0.122*** | | -0.148*** | -0.076*** | -0.006*** | | -0.008** | -0.003* |
| ASM | | | | 0.245*** | 0.192*** | -0.236*** | | | 0.171*** | | |
| AMR | | | -0.010** | -0.228*** | -0.099*** | | | | | | |
| IP visits | | 0.438*** | | | 0.130*** | | | | | | |
| PCP visits | | 0.009*** | | | | | | 0.003*** | | | |
| ED visits | | | | | -0.027*** | | | 0.075*** | | | |
| Non-asthma ED visits | | | | | | | | 0.006*** | 0.439*** | | |
| % with flu | | -0.025*** | -0.004* | | | | | | 0.595*** | -0.012*** | -0.003* |
| % with spacers | | | 0.011*** | | 0.096*** | | | | | | 0.006*** |
| % with nebulizers | | | 0.045*** | 1.700*** | 0.185*** | 2.545*** | | | | | |
| Year | 0.026 | -0.011*** | -0.001 | -0.131*** | -0.116*** | 0.013 | -0.019*** | -0.004** | -0.116* | -0.007* | -0.002 |
| Intervention 2010 | 2.332*** | 0.312*** | 0.029*** | 0.446*** | 0.121*** | 0.426*** | -0.001 | 0.007** | 1.763*** | 0.263*** | 0.030*** |
| DiD Estimator = Year | | | | | | | | | | | |
| * Intervention 2010 | -1.039*** | -0.134*** | -0.022** | -0.160** | -0.035*** | -0.171* | -0.015 | -0.010** | -0.120*** | -0.208*** | -0.016*** |

Notes: * p -value ≤ 0.05 , ** p -value ≤ 0.01 , *** p-value ≤ 0.001 , + $0.05 \leq R^2 < 0.1$, ++ $0.1 \leq R^2 < 0.2$, +++ $0.2 \leq R^2 < 0.3$

Abbreviations: AMR – Asthma medication ratio, ASM – Appropriate medications for people with asthma, AS-ME – Asthma self-management education, ED – Emergency department, IP – Inpatient (hospitalization), PCP – Primary care physician

Table S2. Difference-in Difference (DiD) regression results (New York)

| | AS-ME | | | | Influenza Vaccine | | |
|--|-------------------------------|--------------------------|--------------------------|------------------------|-------------------------------|-------------------------|-------------------------|
| | Utilization Cost ⁺ | Med. Cost ⁺⁺⁺ | ED Visits ⁺⁺⁺ | IP Visits ⁺ | Utilization Cost ⁺ | Med. Cost ⁺⁺ | PCP Visits ⁺ |
| | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. |
| (Intercept) | 0.751 ^{***} | 4.268 ^{***} | 0.967 ^{***} | 0.037 ^{***} | 1.863 ^{***} | 1.243 ^{***} | 0.149 ^{***} |
| Age 9 to 17 | -0.065 ^{***} | 0.647 ^{***} | -0.079 ^{***} | -0.018 ^{***} | | 0.213 ^{***} | -0.046 ^{***} |
| ASM | -0.291 ^{***} | | -0.582 ^{***} | 0.014 ^{***} | 2.531 ^{***} | 2.899 ^{***} | 0.207 ^{***} |
| AMR | -0.193 ^{***} | 2.464 ^{***} | -0.265 ^{***} | -0.026 ^{***} | -0.317 ^{***} | -0.199 ^{***} | -0.115 ^{***} |
| IP | | | | | | | 0.050 ^{***} |
| PCP | | | | | | | |
| ED | | | | 0.088 ^{***} | | | -0.014 ^{***} |
| Medication Cost | | | | | | | 0.0001 ^{***} |
| Non-asthma ED visits | 0.228 ^{***} | -0.085 ^{***} | | | 0.126 ^{***} | | 0.001 ^{***} |
| % with flu | 0.142 ^{***} | | -0.030 ^{***} | | | | |
| % with spacers | -0.169 ^{***} | | 0.096 ^{***} | 0.015 ^{***} | | | 0.070 ^{***} |
| % with nebulizers | | | | | 0.370 ^{***} | 0.355 ^{***} | 0.226 ^{***} |
| Year | -0.072 ^{***} | -0.368 ^{***} | -0.001 | -0.016 ^{***} | -0.070 ^{***} | -0.041 [*] | 0.031 ^{***} |
| Intervention 2010 | 0.166 ^{***} | 0.536 ^{***} | 0.100 ^{***} | 0.046 ^{***} | 0.364 ^{***} | 0.335 ^{***} | 0.114 ^{***} |
| DiD Estimator = Year * Intervention 2010 | -0.185 ^{***} | -0.163 ^{**} | -0.060 ^{***} | -0.035 ^{***} | -0.179 ^{***} | -0.170 ^{***} | -0.060 ^{***} |

Table S2. Difference-in Difference (DiD) regression results (New York) (continued)

| | Spacer | | | | Nebulizer | | | |
|----------------------|-------------------------------|---------------------------|--------------------------|-------------------------|-------------------------------|---------------------------|--------------------------|------------------------|
| | Utilization Cost ⁺ | Med. Cost ⁺⁺⁺⁺ | ED Visits ⁺⁺⁺ | IP Visits ⁺⁺ | Utilization Cost ⁺ | Med. Cost ⁺⁺⁺⁺ | ED Visits ⁺⁺⁺ | IP Visits ⁺ |
| | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. | Coef. |
| (Intercept) | 0.728*** | 2.408*** | 0.680*** | 0.035*** | 1.258*** | 2.254*** | 0.916*** | 0.020*** |
| Age 9 to 17 | | 0.255*** | -0.131*** | -0.018*** | -0.066*** | 0.349*** | -0.029*** | -0.012*** |
| ASM | -0.318*** | 0.373*** | | 0.014*** | -0.783*** | 3.731*** | -0.810*** | 0.015*** |
| AMR | -0.179*** | | -0.612*** | -0.026*** | -0.248*** | 0.410*** | | -0.022*** |
| IP | | | | | | | | |
| PCP | | 0.042*** | | | | | | |
| ED | | 0.029*** | | 0.087*** | | | | 0.128*** |
| Medication Cost | | | | | | | | |
| Non-asthma ED visits | 0.243*** | | | -0.002*** | | | | -0.002* |
| % with flu | 0.155*** | | -0.033*** | | 0.146*** | | -0.026*** | |
| % with spacers | | | | | -0.155*** | | | 0.025*** |
| % with nebulizers | | | | | | | | |
| Year | -0.071*** | -0.449*** | -0.004# | -0.015*** | -0.060*** | -0.421*** | -0.005* | -0.009*** |
| Intervention 2010 | 0.175*** | 0.032 | 0.053*** | 0.042*** | 0.193*** | 0.112*** | 0.188*** | 0.026*** |
| DiD Estimator = Year | | | | | | | | |
| * Intervention 2010 | -0.112** | -0.070# | -0.021* | -0.020*** | -0.115*** | -0.057* | -0.124*** | -0.008* |

Notes: # p ≤ 0.1, * p-value ≤ 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001, + 0.05 ≤ R² < 0.1, ++ 0.1 ≤ R² < 0.2, +++ 0.2 ≤ R² < 0.3, ++++ 0.3 ≤ R² < 0.4.

Abbreviations: AMR – Asthma medication ratio, ASM – Appropriate medications for people with asthma, AS-ME – Asthma self-management education, ED – Emergency department, IP – Inpatient (hospitalization), PCP – Primary care physician




Figure S1. Cost trendlines for influenza vaccine




Figure S2. Probability of utilization trendlines for influenza vaccine

Abbreviations: AS-ME – Asthma self-management education, ED – Emergency department, IP – Inpatient (hospitalization), PCP – Primary care physician




Figure S3. Cost trendlines for spacer




Figure S4. Probability of utilization trendlines for spacer

Abbreviations: AS-ME – Asthma self-management education, ED – Emergency department, IP – Inpatient (hospitalization), PCP – Primary care physician




Figure S5. Cost trendlines for nebulizer




Figure S6. Probability of utilization trendlines for nebulizer

Abbreviations: AS-ME – Asthma self-management education, ED – Emergency department, IP – Inpatient (hospitalization), PCP – Primary care physician